

Table 19.11  
Non-Hodgkin Lymphoma

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Survival Time																				
1-year	71.0	73.3	74.1	74.7	75.1	75.8	76.8	78.7	79.1	78.1	80.1	81.4	81.0	82.1	82.3	80.9	81.8	84.5	85.1	84.7
2-year	61.1	63.7	65.6	67.2	66.3	68.7	70.1	72.0	73.2	72.6	75.8	76.9	76.6	76.7	78.4	77.3	76.9	80.1	80.8	
3-year	55.9	58.9	61.1	62.8	61.4	64.9	66.2	68.5	69.9	69.1	72.8	74.4	73.7	74.6	76.7	75.5	74.2	77.0		
4-year	51.9	55.2	57.6	59.5	58.8	61.8	63.4	65.4	68.0	67.4	71.1	72.5	72.4	73.0	74.9	73.8	72.4			
5-year	48.2	51.3	54.3	56.7	56.3	59.0	60.8	63.0	65.7	66.4	68.9	70.9	71.3	71.5	73.2	72.2				
6-year	44.6	48.4	51.7	53.9	53.8	58.0	58.4	60.5	63.3	64.3	68.0	69.6	69.9	69.9	72.1					
7-year	41.7	45.8	49.0	51.1	51.8	56.0	56.4	58.7	61.3	62.5	66.4	67.3	68.6	68.8						
8-year	39.2	43.7	46.3	49.1	50.4	54.1	54.0	57.4	59.4	61.4	65.0	66.7	67.0							
9-year	37.4	41.6	44.3	47.6	49.1	52.9	52.9	55.9	57.6	60.1	63.3	65.2								
10-year	36.0	39.8	42.4	45.8	47.8	51.3	51.7	54.4	55.9	58.5	62.3									
11-year	34.2	38.5	40.9	43.8	46.6	50.1	50.1	52.9	54.4	56.8										
12-year	32.8	36.7	39.0	42.5	45.7	49.2	49.3	52.3	52.5											
13-year	32.1	35.2	37.5	41.1	44.6	48.3	48.3	51.7												
14-year	30.9	34.0	36.5	40.0	43.9	47.6	46.7													
15-year	29.6	32.5	34.9	39.1	43.2	46.7														
16-year	28.7	31.5	34.2	38.5	42.9															
17-year	28.1	30.6	33.6	37.7																
18-year	27.3	29.7	32.7																	
19-year	26.4	28.7	32.0																	
20-year	25.3	27.9	31.1																	

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).