

Table 19.16
Non-Hodgkin Lymphoma

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years,
Lifetime Risk of Being Diagnosed with Cancer Given Alive and Cancer-Free at Current Age, and
Lifetime Risk of Dying from Cancer Given Alive at Current Age
Both Sexes, 2009-2011 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer
		+10 yrs	+20 yrs	+30 yrs	Ever				+10 yrs	+20 yrs	+30 yrs	Ever	
All Races	0	0.01	0.02	0.05	2.12	0.78	Asian/ Pacific Islander	0	0.01	0.02	0.04	1.83	0.74
	10	0.01	0.04	0.10	2.14	0.79		10	0.01	0.03	0.07	1.84	0.75
	20	0.03	0.09	0.21	2.13	0.79		20	0.02	0.06	0.13	1.83	0.75
	30	0.06	0.18	0.42	2.12	0.79		30	0.04	0.11	0.28	1.81	0.75
	40	0.12	0.36	0.82	2.09	0.79		40	0.07	0.24	0.57	1.78	0.75
	50	0.25	0.72	1.38	2.02	0.80		50	0.17	0.50	1.00	1.73	0.75
	60	0.50	1.20	1.75	1.88	0.80		60	0.34	0.86	1.41	1.61	0.74
	70	0.80	1.42	-	1.57	0.76		70	0.56	1.15	-	1.36	0.69
80	0.85	-	-	1.05	0.63	80	0.71	-	-	0.97	0.57		
White	0	0.01	0.02	0.05	2.22	0.82	American Indian/ Alaska Native ^a	0	0.01	0.02	0.04	1.33	0.50
	10	0.02	0.04	0.10	2.24	0.82		10	0.01	0.03	0.07	1.34	0.50
	20	0.03	0.09	0.21	2.23	0.83		20	0.02	0.06	0.13	1.34	0.50
	30	0.06	0.18	0.43	2.22	0.83		30	0.05	0.11	0.25	1.34	0.51
	40	0.13	0.37	0.86	2.19	0.83		40	0.07	0.21	0.53	1.32	0.52
	50	0.26	0.75	1.45	2.12	0.84		50	0.15	0.49	0.93	1.31	0.53
	60	0.52	1.27	1.84	1.97	0.84		60	0.36	0.83	1.15	1.24	0.55
	70	0.85	1.50	-	1.64	0.80		70	0.54	0.90	-	1.00	0.51
80	0.89	-	-	1.09	0.66	80	0.50	-	-	0.63	0.33		
Black	0	0.01	0.02	0.05	1.28	0.45	Hispanic ^b	0	0.01	0.02	0.04	2.23	0.77
	10	0.01	0.05	0.12	1.30	0.46		10	0.01	0.04	0.09	2.25	0.78
	20	0.04	0.11	0.23	1.29	0.46		20	0.03	0.07	0.18	2.24	0.78
	30	0.07	0.20	0.40	1.28	0.46		30	0.05	0.15	0.37	2.23	0.78
	40	0.13	0.34	0.65	1.23	0.46		40	0.11	0.33	0.76	2.20	0.78
	50	0.22	0.54	0.88	1.14	0.45		50	0.23	0.67	1.34	2.14	0.78
	60	0.36	0.73	0.95	1.02	0.43		60	0.46	1.17	1.79	2.00	0.76
	70	0.45	0.72	-	0.80	0.39		70	0.78	1.47	-	1.69	0.72
80	0.40	-	-	0.51	0.31	80	0.88	-	-	1.16	0.58		

Devcan 6.8.0, August 2014, National Cancer Institute (<http://surveillance.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic could not be calculated.

A percent of 0.00 represents a value that is below 0.005.