

Table 19.17
Non-Hodgkin Lymphoma

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years,
Lifetime Risk of Being Diagnosed with Cancer Given Alive and Cancer-Free at Current Age, and
Lifetime Risk of Dying from Cancer Given Alive at Current Age
Males, 2009-2011 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer
		+10 yrs	+20 yrs	+30 yrs	Ever				+10 yrs	+20 yrs	+30 yrs	Ever	
All Races	0	0.01	0.03	0.06	2.36	0.87	Asian/ Pacific Islander	0	0.01	0.03	0.05	2.06	0.83
	10	0.02	0.05	0.13	2.38	0.88		10	0.02	0.04	0.08	2.06	0.84
	20	0.04	0.11	0.25	2.37	0.89		20	0.02	0.06	0.14	2.05	0.84
	30	0.07	0.22	0.49	2.37	0.89		30	0.04	0.12	0.30	2.04	0.84
	40	0.15	0.43	0.94	2.33	0.90		40	0.08	0.27	0.66	2.02	0.84
	50	0.29	0.82	1.57	2.26	0.91		50	0.19	0.58	1.20	1.97	0.84
	60	0.58	1.38	1.99	2.12	0.91		60	0.41	1.06	1.65	1.85	0.83
	70	0.94	1.65	-	1.81	0.87		70	0.70	1.35	-	1.57	0.78
80	1.02	-	-	1.26	0.75	80	0.82	-	-	1.10	0.67		
White	0	0.01	0.03	0.06	2.46	0.92	American Indian/ Alaska Native ^a	0	0.02	0.03	0.06	1.36	0.52
	10	0.02	0.06	0.13	2.48	0.93		10	0.02	0.04	0.08	1.36	0.52
	20	0.04	0.11	0.25	2.47	0.93		20	0.02	0.07	0.13	1.35	0.53
	30	0.07	0.22	0.50	2.46	0.94		30	0.05	0.10	0.21	1.36	0.53
	40	0.15	0.43	0.98	2.43	0.94		40	0.06	0.16	0.52	1.35	0.55
	50	0.30	0.86	1.65	2.36	0.95		50	0.11	0.48	1.04	1.35	0.57
	60	0.60	1.45	2.08	2.22	0.95		60	0.40	1.01	1.34	1.35	0.58
	70	0.99	1.73	-	1.89	0.92		70	0.72	1.11	-	1.12	0.51
80	1.07	-	-	1.30	0.78	80	0.56	-	-	0.58	0.24		
Black	0	0.01	0.03	0.07	1.41	0.51	Hispanic ^b	0	0.01	0.02	0.05	2.39	0.82
	10	0.02	0.06	0.15	1.43	0.52		10	0.02	0.05	0.10	2.41	0.82
	20	0.04	0.14	0.29	1.43	0.52		20	0.03	0.09	0.20	2.40	0.83
	30	0.09	0.26	0.51	1.41	0.52		30	0.06	0.17	0.42	2.39	0.83
	40	0.17	0.43	0.77	1.35	0.52		40	0.12	0.36	0.84	2.37	0.83
	50	0.27	0.63	0.98	1.24	0.51		50	0.25	0.75	1.49	2.31	0.83
	60	0.40	0.80	1.02	1.10	0.49		60	0.52	1.31	1.98	2.18	0.82
	70	0.50	0.78	-	0.88	0.46		70	0.90	1.65	-	1.87	0.77
80	0.46	-	-	0.62	0.37	80	1.01	-	-	1.31	0.61		

Devcan 6.8.0, August 2014, National Cancer Institute (<http://surveillance.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic could not be calculated.

A percent of 0.00 represents a value that is below 0.005.