

Table 19.25  
Non-Hodgkin LymphomaEstimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2011  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 <sup>e</sup>	0 to <36 <sup>e</sup>	>=36 <sup>g</sup>	Complete <sup>h</sup>	
Race	Sex											
All Races <sup>b</sup>	Both Sexes	203,992	138,548	82,649	46,499	26,613	14,965	478,424	522,113	8,806	530,919	
	Males	109,031	71,416	42,880	24,393	14,018	8,043	251,155	274,206	4,630	278,836	
	Females	94,961	67,132	39,769	22,106	12,595	6,922	227,269	247,907	4,176	252,083	
White <sup>b</sup>	Both Sexes	176,587	121,718	73,137	41,335	23,890	13,662	418,719	458,521	9,171	467,692	
	Males	94,865	62,935	38,025	21,688	12,655	7,313	220,566	241,606	5,180	246,786	
	Females	81,722	58,783	35,112	19,647	11,235	6,349	198,153	216,915	3,991	220,906	
Black <sup>b</sup>	Both Sexes	16,393	11,194	5,958	3,101	1,826	801	37,209	39,725	715	40,440	
	Males	8,305	5,551	3,045	1,601	926	456	18,751	20,065	299	20,364	
	Females	8,088	5,643	2,913	1,500	900	345	18,458	19,660	416	20,076	
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	5,757	3,689	2,092	+	+	+	12,532	+	+	+	
	Males	3,044	1,904	1,117	+	+	+	6,533	+	+	+	
	Females	2,713	1,785	975	+	+	+	5,999	+	+	+	
Hispanic <sup>d</sup>	Both Sexes	15,678	8,999	5,398	+	+	+	32,329	+	+	+	
	Males	8,083	4,633	2,885	+	+	+	16,751	+	+	+	
	Females	7,595	4,366	2,513	+	+	+	15,578	+	+	+	

Estimated prevalence percent<sup>a</sup> on January 1, 2011, of the SEER<sup>c</sup> population diagnosed in the previous 19 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)								Age-Adjusted <sup>f</sup>		
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
Race	Sex											
All Races <sup>c</sup>	Both Sexes	0.1349%	0.0032%	0.0117%	0.0244%	0.0455%	0.0974%	0.1924%	0.3624%	0.5763%	0.6536%	0.1311%
	Males	0.1439%	0.0039%	0.0156%	0.0294%	0.0537%	0.1144%	0.2225%	0.4110%	0.6473%	0.7399%	0.1504%
	Females	0.1260%	0.0025%	0.0077%	0.0191%	0.0373%	0.0806%	0.1637%	0.3184%	0.5192%	0.6031%	0.1143%
White <sup>c</sup>	Both Sexes	0.1516%	0.0033%	0.0121%	0.0257%	0.0477%	0.1031%	0.2075%	0.3920%	0.6328%	0.6985%	0.1412%
	Males	0.1608%	0.0042%	0.0164%	0.0311%	0.0564%	0.1210%	0.2388%	0.4395%	0.7091%	0.7841%	0.1612%
	Females	0.1423%	0.0024%	0.0076%	0.0200%	0.0385%	0.0847%	0.1766%	0.3477%	0.5699%	0.6479%	0.1232%
Black <sup>c</sup>	Both Sexes	0.0835%	0.0024%	0.0121%	0.0239%	0.0472%	0.0935%	0.1607%	0.2708%	0.3385%	0.3185%	0.0953%
	Males	0.0896%	0.0024%	0.0154%	0.0302%	0.0554%	0.1105%	0.1894%	0.3229%	0.3191%	0.3524%	0.1073%
	Females	0.0779%	0.0025%	0.0086%	0.0180%	0.0401%	0.0785%	0.1358%	0.2297%	0.3520%	0.3023%	0.0852%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0828%	0.0036%	0.0090%	0.0174%	0.0299%	0.0604%	0.1147%	0.2145%	0.3575%	0.4777%	0.0835%
	Males	0.0895%	0.0039%	0.0115%	0.0199%	0.0347%	0.0680%	0.1318%	0.2465%	0.4247%	0.5732%	0.0977%
	Females	0.0768%	0.0032%	0.0064%	0.0148%	0.0257%	0.0537%	0.1001%	0.1880%	0.3059%	0.4190%	0.0724%
Hispanic <sup>d</sup>	Both Sexes	0.0646%	0.0032%	0.0089%	0.0184%	0.0328%	0.0729%	0.1482%	0.2770%	0.4412%	0.5261%	0.1014%
	Males	0.0657%	0.0040%	0.0115%	0.0216%	0.0390%	0.0810%	0.1665%	0.2893%	0.4502%	0.5719%	0.1098%
	Females	0.0635%	0.0025%	0.0061%	0.0148%	0.0262%	0.0645%	0.1307%	0.2664%	0.4345%	0.4989%	0.0939%

<sup>a</sup> US 2011 cancer prevalence counts are based on 2011 cancer prevalence proportions from the SEER registries and 1/1/2011 US population estimates based on the average of 2010 and 2011 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 36 years for 1975-2011 SEER 9 data; 19 years for 1992-2011 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 36 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <36 and >=36. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.