

Table 23.7
Cancer of the Prostate (Invasive)

SEER Incidence^a and U.S. Death^b Rates, Age-Adjusted and Age-Specific Rates, by Race

	SEER Incidence ^a			U.S. Mortality ^b		
	All Races Males	White Males	Black Males	All Races Males	White Males	Black Males
<u>Age-Adjusted Rates, 2007-2011</u>						
Age at Diagnosis/Death:						
All ages	147.8	139.9	223.9	22.3	20.6	48.9
Under 65	57.9	54.2	101.2	1.7	1.5	4.4
65 and over	769.1	732.2	1,071.9	164.9	153.0	356.7
All ages (IARC world std) ^c	102.3	97.0	159.4	9.7	8.8	22.0
<u>Age-Specific Rates, 2007-2011</u>						
Age at Diagnosis/Death:						
<1	-	-	-	-	-	-
1-4	-	-	-	-	-	-
5-9	-	-	-	-	-	-
10-14	-	-	-	-	-	-
15-19	-	-	-	-	-	-
20-24	-	-	-	-	-	-
25-29	-	-	-	-	-	-
30-34	0.1	-	-	-	-	-
35-39	1.2	1.0	2.9	0.0	-	-
40-44	10.5	8.9	25.7	0.2	0.1	0.4
45-49	47.3	41.0	106.5	0.9	0.7	2.3
50-54	145.9	134.7	269.1	3.2	2.7	8.1
55-59	321.6	303.5	547.3	8.9	7.5	22.7
60-64	552.9	527.2	900.8	20.5	17.7	52.6
65-69	835.2	800.8	1,246.4	41.8	36.6	104.5
70-74	895.0	859.0	1,215.1	80.2	71.1	198.8
75-79	806.4	763.2	1,060.6	144.6	132.1	329.4
80-84	603.9	564.3	804.7	257.0	239.8	567.3
85+	490.6	460.1	720.4	539.4	514.3	1,042.2

^a SEER 18 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^c Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

- Rates are per 100,000 and are age-adjusted to the IARC world standard population.

- Statistic not shown. Rate based on less than 16 cases for the time interval.