Table 32.11

Age-Adjusted SEER Incidence^a, by Year and Sex
By Adapted Classification Scheme for Tumors of Adolescents and Young Adults^b
All Races, Ages 35-39

| | 1975-2011 | | | | 2002-2011 | | |
|---|-----------|-------|-------------------|-----------|-----------|-------|--|
| Both Sexes | No. | Rate | AAPC | No. | Rate | AAPC | |
| | 01 005 | 100 5 | 0.01 | 0.7.010 | 100.0 | 0.01 | |
| All Sites Combined | 91,805 | 132.7 | 0.8* | 27,810 | 138.0 | 0.9* | |
| 1. Leukemias | 2,583 | 3.7 | 0.2 | 777 | 3.9 | 0.2 | |
| 2. Lymphomas | 7,726 | 11.2 | 1.6* | 2,250 | 11.2 | -1.4* | |
| 3. CNS, intracranial, intraspinal | 2,736 | 4.0 | -0.2 | 758 | 3.8 | -0.2 | |
| 4. Osseous & chondromatous neoplasms | 456 | 0.7 | 1.4* | 154 | 0.8 | 1.4* | |
| 5. Soft tissue sarcomas | 5,935 | 8.6 | 1.1 | 1,025 | 5.1 | -1.0 | |
| 6. Germ cell and trophoblastic neoplasms | 4,060 | 5.9 | 1.7* | 1,311 | 6.5 | 0.3 | |
| 7. Melanoma and skin carcinomas | 10,003 | 14.5 | 1.6* | 3,148 | 15.6 | 0.4* | |
| 8. Carcinomas (excluding skin) | 56,654 | 81.9 | 0.5 | 17,877 | 88.7 | 1.4* | |
| 9. Miscellaneous specified neoplasms, NOS | 1,178 | 1.7 | 1.1* | 391 | 1.9 | 1.1* | |
| 10. Unspecified malignant neoplasms | 474 | 0.7 | -0.2 | 119 | 0.6 | -0.2 | |
| | 1975-2011 | | | 2002-2011 | | | |
| Males | No. | Rate | AAPC° | No. | Rate | AAPC | |
| | | | | | | | |
| All Sites Combined | 34,583 | 100.3 | 0.6 | 9,688 | 96.2 | 0.1 | |
| 1. Leukemias | 1,500 | 4.4 | 0.1 | 443 | 4.4 | 0.1 | |
| 2. Lymphomas | 5,044 | 14.6 | 1.8* | 1,373 | 13.6 | -0.7 | |
| 3. CNS, intracranial, intraspinal | 1,604 | 4.7 | -0.1 | 439 | 4.4 | -0.1 | |
| 4. Osseous & chondromatous neoplasms | 262 | 0.8 | 0.8 | 88 | 0.9 | 0.8 | |
| 5. Soft tissue sarcomas | 4,683 | 13.6 | 2.6 | 674 | 6.7 | -1.3 | |
| 6. Germ cell and trophoblastic neoplasms | 3,851 | 11.2 | 1.8* | 1,245 | 12.4 | 0.3 | |
| 7. Melanoma and skin carcinomas | 4,362 | 12.7 | 0.7 | 1,253 | 12.4 | -0.6 | |
| 8. Carcinomas (excluding skin) | 12,614 | 36.6 | 0.2 | 3,977 | 39.5 | 1.4* | |
| 9. Miscellaneous specified neoplasms, NOS | 471 | 1.4 | 0.5 | 153 | 1.5 | 0.5 | |
| 10. Unspecified malignant neoplasms | 192 | 0.6 | -0.7 | 43 | 0.4 | -0.7 | |
| | 1975-2011 | | , | 2002-2011 | | | |
| Females | No. | Rate | AAPC ^c | No. | Rate | AAPC° | |
| | | | | | | | |
| All Sites Combined | 57,222 | 164.9 | 0.7* | 18,122 | 179.7 | 1.2* | |
| 1. Leukemias | 1,083 | 3.1 | 0.9 | 334 | 3.3 | 2.4* | |
| 2. Lymphomas | 2,682 | 7.7 | 1.4* | 877 | 8.7 | 1.4* | |
| 3. CNS, intracranial, intraspinal | 1,132 | 3.3 | -0.3 | 319 | 3.2 | -0.3 | |
| 4. Osseous & chondromatous neoplasms | 194 | 0.6 | _ | 66 | 0.7 | _ | |
| 5. Soft tissue sarcomas | 1,252 | 3.6 | -0.6 | 351 | 3.5 | -0.6 | |
| 6. Germ cell and trophoblastic neoplasms | 209 | 0.6 | -0.1 | 66 | 0.7 | -0.1 | |
| 7. Melanoma and skin carcinomas | 5,641 | 16.3 | 2.1* | 1,895 | 18.8 | 1.8* | |
| 8. Carcinomas (excluding skin) | 44,040 | 126.9 | 0.6 | 13,900 | 137.8 | 1.4* | |
| 9. Miscellaneous specified neoplasms, NOS | 707 | 2.0 | 1.4* | 238 | 2.4 | 1.4* | |
| 10. Unspecified malignant neoplasms | 282 | 0.8 | -0.1 | 76 | 0.8 | -0.1 | |
| | 202 | | ٠.ـ | , 0 | 0.0 | · · - | |

Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

* The AAPC is significantly different from zero (p<.05).

Rates are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta).

Adapted from: RD Barr, EJ Holowaty, JM Birch. Classification Scheme for tumors diagnosed in adolescents and young adults. Cancer. 106:7; 1425-1430. (Restricted to malignant tumors and updated based on ICD-0-3 definitions).

 $^{^{\}rm c}$ $\,$ The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint over the time range of 1975-2011.

⁻ Statistic could not be calculated. Count and rate based on less than 16 cases for the time interval. AAPC could not be calculated due to insufficient data.