

Table 1.22
U.S. Prevalence Counts, Invasive Cancers Only, January 1, 2012^a
Using Different Tumor Inclusion Criteria^b

Site	Sex	5-Year Limited Duration			37-year Limited Duration	
		1st Invasive Tumor Ever ^c	1st Per Site in Previous 37 Years ^d	1st Per Site in Previous 5 Years ^e	1st Invasive Tumor Ever ^c	1st Per Site in Previous 37 Years ^d
All Sites	Both Sexes	4,660,390	4,751,475	5,238,866	13,405,317	13,624,510
	Male	2,395,057	2,430,935	2,669,928	6,387,558	6,462,521
	Female	2,265,333	2,320,540	2,568,938	7,017,759	7,161,989
Oral Cavity & Pharynx	Both Sexes	108,762	126,337	131,109	282,380	312,792
	Male	76,856	88,637	91,609	188,848	207,488
	Female	31,906	37,700	39,500	93,532	105,304
Esophagus	Both Sexes	21,344	26,589	26,680	35,667	42,388
	Male	16,607	20,581	20,639	27,750	32,814
	Female	4,737	6,008	6,041	7,917	9,574
Stomach	Both Sexes	37,369	45,065	45,492	75,640	87,222
	Male	22,014	27,032	27,269	43,300	50,402
	Female	15,355	18,033	18,223	32,340	36,820
Colon & Rectum	Both Sexes	398,737	458,984	468,251	1,149,814	1,279,479
	Male	204,170	234,705	239,034	573,302	635,108
	Female	194,567	224,279	229,217	576,512	644,371
Liver & Intrahepatic Bile Duct	Both Sexes	36,620	41,939	41,976	50,271	56,613
	Male	27,039	30,599	30,620	35,990	40,166
	Female	9,581	11,340	11,356	14,281	16,447
Pancreas	Both Sexes	34,042	41,747	41,771	45,432	54,507
	Male	17,628	21,807	21,819	22,666	27,478
	Female	16,414	19,940	19,952	22,766	27,029
Larynx	Both Sexes	31,457	37,206	37,786	87,479	98,015
	Male	25,542	30,220	30,728	70,303	78,592
	Female	5,915	6,986	7,058	17,176	19,423
Lung & Bronchus	Both Sexes	233,827	301,827	311,522	402,326	498,404
	Male	107,351	140,977	144,670	180,530	225,061
	Female	126,476	160,850	166,852	221,796	273,343
Melanoma of the Skin	Both Sexes	285,569	326,073	342,024	962,438	1,041,311
	Male	153,364	179,735	189,728	478,737	525,778
	Female	132,205	146,338	152,296	483,701	515,533
Breast	Female	872,541	944,632	1,005,580	2,928,546	3,104,561
Cervix	Female	39,323	41,561	41,655	214,008	220,136
Corpus & Uterus, NOS	Female	183,434	205,521	205,655	598,766	653,998
Ovary ^f	Female	59,878	68,736	68,817	177,706	199,493

^a U.S. 2012 cancer prevalence counts are based on 2012 cancer prevalence proportions from the SEER 9 registries and 1/1/2012 U.S. population estimates based on the average of 2011 and 2012 population estimates from the U.S. Bureau of the Census.

^b Prevalence estimates are ambiguous for those with multiple cancers, unless the tumor inclusion criteria are understood. Depending on the application, different inclusion criteria may be appropriate. This table provides three different methods of tumor inclusion:

^c (c) First invasive tumor ever

^d (d) First invasive tumor for each cancer site diagnosed during the previous 37 years (1975-2011)

^e (e) First invasive tumor for each cancer site diagnosed during the previous 5 years (2007-2011)

For definitions (d) and (e) all sites is treated as a separate cancer "site".

Consider a woman who had three invasive cancers: Melanoma in 1981; Breast cancer in 2007; Melanoma in 2008.

In method (c) the melanoma is the woman's first cancer, and is counted for the melanoma and all sites 37-year limited duration prevalence. For 5-year limited duration prevalence, the woman is not counted at all since her first cancer occurred more than 5 years prior to 1/1/2012.

In method (d) the 1981 melanoma is counted for the melanoma and all sites 37-year limited duration prevalence. The 2007 breast cancer is counted for the breast 5-year and 37-year limited duration prevalence.

In method (e) the 2007 breast cancer is counted for the breast cancer and all sites 5-year limited duration prevalence. The 2008 melanoma is counted for 5-year limited duration prevalence for melanoma.

^f Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

Table 1.22 - continued
 U.S. Prevalence Counts, Invasive Cancers Only, January 1, 2012^a
 Using Different Tumor Inclusion Criteria^b

Site	Sex	5-Year Limited Duration			37-year Limited Duration	
		1st Invasive Tumor Ever ^c	1st Per Site in Previous 37 Years ^d	1st Per Site in Previous 5 Years ^e	1st Invasive Tumor Ever ^c	1st Per Site in Previous 37 Years ^d
Prostate	Male	1,044,262	1,128,141	1,128,224	2,794,570	2,976,626
Testis	Male	42,435	43,354	44,013	218,903	221,980
Urinary Bladder	Both Sexes	201,191	253,322	258,716	567,531	664,041
	Male	153,198	193,786	198,124	424,112	496,868
	Female	47,993	59,536	60,592	143,419	167,173
Kidney & Renal Pelvis	Both Sexes	154,353	188,076	190,370	365,660	427,078
	Male	94,867	117,507	119,055	217,927	257,221
	Female	59,486	70,569	71,315	147,733	169,857
Brain & Nervous System	Both Sexes	45,479	48,676	49,045	136,100	140,756
	Male	24,643	26,423	26,633	72,270	74,764
	Female	20,836	22,253	22,412	63,830	65,992
Thyroid	Both Sexes	185,714	207,444	208,227	563,884	604,543
	Male	40,645	47,892	48,044	121,802	134,236
	Female	145,069	159,552	160,183	442,082	470,307
Hodgkin Lymphoma	Both Sexes	38,167	40,722	40,777	176,990	181,967
	Male	20,781	22,226	22,270	91,797	94,359
	Female	17,386	18,496	18,507	85,193	87,608
Non-Hodgkin Lymphoma	Both Sexes	205,597	242,575	245,938	541,393	606,972
	Male	110,512	131,592	133,324	284,672	319,941
	Female	95,085	110,983	112,614	256,721	287,031
Myeloma	Both Sexes	56,508	67,424	67,605	89,416	103,463
	Male	30,245	36,895	36,980	47,990	56,338
	Female	26,263	30,529	30,625	41,426	47,125
Leukemia	Both Sexes	120,252	139,986	140,426	312,923	345,422
	Male	70,027	82,564	82,799	177,810	197,506
	Female	50,225	57,422	57,627	135,113	147,916
Acute Lymphocytic Leukemia	Both Sexes	17,081	17,590	17,590	71,898	72,647
	Male	9,480	9,693	9,693	39,375	39,666
	Female	7,601	7,897	7,897	32,523	32,981
Childhood (Ages 0-19)	Both Sexes	65,893	65,980	66,533	326,769	327,259
	Male	34,508	34,555	34,858	167,928	168,153
	Female	31,385	31,425	31,675	158,841	159,106
Kaposi Sarcoma	Both Sexes	7,384	7,975	7,975	26,300	27,684
	Male	6,836	7,367	7,367	24,829	26,079
	Female	548	608	608	1,471	1,605
Mesothelioma	Both Sexes	3,247	4,116	4,116	4,957	5,962
	Male	2,341	2,982	2,982	3,042	3,748
	Female	906	1,134	1,134	1,915	2,214

^a U.S. 2012 cancer prevalence counts are based on 2012 cancer prevalence proportions from the SEER 9 registries and 1/1/2012 U.S. population estimates based on the average of 2011 and 2012 population estimates from the U.S. Bureau of the Census.

^b Prevalence estimates are ambiguous for those with multiple cancers, unless the tumor inclusion criteria are understood. Depending on the application, different inclusion criteria may be appropriate. This table provides three different methods of tumor inclusion:

^c (c) First invasive tumor ever
^d (d) First invasive tumor for each cancer site diagnosed during the previous 37 years (1975-2011)
^e (e) First invasive tumor for each cancer site diagnosed during the previous 5 years (2007-2011)
 For definitions (d) and (e) all sites is treated as a separate cancer "site".

Consider a woman who had three invasive cancers: Melanoma in 1981; Breast cancer in 2007; Melanoma in 2008.

In method (c) the melanoma is the woman's first cancer, and is counted for the melanoma and all sites 37-year limited duration prevalence. For 5-year limited duration prevalence, the woman is not counted at all since her first cancer occurred more than 5 years prior to 1/1/2012. In method (d) the 1981 melanoma is counted for the melanoma and all sites 37-year limited duration prevalence. The 2007 breast cancer is counted for the breast 5-year and 37-year limited duration prevalence. In method (e) the 2007 breast cancer is counted for the breast cancer and all sites 5-year limited duration prevalence. The 2008 melanoma is counted for 5-year limited duration prevalence for melanoma.

Table 1.23
U.S. Complete Prevalence Counts, Invasive Cancers Only, January 1, 2012^a
By Age at Prevalence

Site/Sex	Age at Prevalence								
	All Ages ^c	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
All Sites									
Males	6,464,529	19,057	42,522	87,325	156,133	356,474	907,855	1,732,881	3,162,283
Females	7,311,722	17,006	36,354	89,937	229,430	636,759	1,353,262	1,832,064	3,116,909
Oral Cavity & Pharynx									
Males	193,779	73	416	1,506	3,443	14,067	46,864	62,089	65,320
Females	97,329	96	569	1,562	3,831	8,690	18,524	24,531	39,526
Esophagus									
Males	27,808	0	0	34	126	993	4,575	10,168	11,912
Females	7,973	0	0	11	11	207	1,315	2,059	4,369
Stomach									
Males	43,975	0	46	105	596	2,524	7,100	12,229	21,374
Females	32,854	3	23	172	691	2,440	4,745	7,157	17,623
Colon & Rectum									
Males	579,901	11	93	1,215	5,831	26,335	84,353	142,161	319,901
Females	589,028	0	105	1,463	5,773	23,768	72,770	115,560	369,590
Liver & Intrahep									
Males	36,063	560	460	615	605	1,716	11,082	13,605	7,421
Females	14,671	438	522	404	506	925	3,227	4,037	4,612
Pancreas									
Males	22,783	34	18	152	297	1,722	4,835	7,158	8,566
Females	22,919	0	33	237	479	1,550	4,120	6,272	10,228
Larynx									
Males	71,414	0	0	91	244	1,657	10,356	21,554	37,512
Females	17,438	0	0	47	128	806	3,474	4,783	8,200
Lung & Bronchus									
Males	184,230	34	75	504	1,243	5,258	24,519	55,147	97,451
Females	224,578	34	92	414	1,489	6,994	30,650	60,573	124,332
Melanoma of the Skin									
Males	489,614	45	685	5,156	17,606	45,367	99,463	137,601	183,691
Females	506,973	102	812	10,505	34,549	73,225	119,346	119,937	148,497

^a U.S. 2012 cancer prevalence counts are based on 2012 cancer prevalence proportions from the SEER 9 registries (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta) and 1/1/2012 U.S. population estimates based on the average of 2011 and 2012 population estimates from the U.S. Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^b Cases diagnosed more than 37 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^c Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.

Table 1.23 - continued
 U.S. Complete Prevalence Counts, Invasive Cancers Only, January 1, 2012^a
 By Age at Prevalence

Site/Sex	Age at Prevalence								
	All Ages ^c	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
Breast									
Males	15,499	0	0	11	134	699	1,855	4,533	8,267
Females	2,975,314	11	35	3,037	35,604	214,136	559,716	824,053	1,338,721
Cervix									
Females	249,512	0	68	2,062	15,420	41,029	59,636	58,895	72,402
Corpus & Uterus, NOS									
Females	621,612	11	23	614	5,890	24,821	91,757	174,495	324,000
Ovary ^d									
Females	192,446	77	1,080	3,491	6,874	18,099	42,498	51,208	69,119
Prostate									
Males	2,795,592	34	47	102	293	20,286	240,483	807,991	1,726,357
Urinary Bladder									
Males	430,485	56	117	541	2,355	10,502	43,269	104,139	269,506
Females	146,918	45	34	208	1,046	4,028	13,688	31,217	96,652
Kidney & Renal Pelvis									
Males	223,683	1,445	2,452	2,765	4,922	19,349	43,876	64,800	84,074
Females	152,242	1,519	2,612	2,882	4,794	12,658	26,698	37,929	63,150
Hodgkin Lymphoma									
Males	98,082	237	2,413	9,323	16,360	23,948	22,943	15,053	7,806
Females	91,544	39	1,969	9,432	16,470	22,866	20,748	12,271	7,749
Non-Hodgkin Lymphoma									
Males	289,004	794	3,674	8,038	13,729	29,671	55,734	74,542	102,823
Females	260,621	475	1,812	4,755	8,981	21,174	43,182	63,503	116,739
Myeloma									
Males	48,074	0	6	121	521	3,285	8,685	16,099	19,356
Females	41,584	0	0	28	392	2,271	7,446	12,033	19,413
Leukemia									
Males	180,231	6,723	13,446	13,043	11,701	14,648	25,247	36,927	58,495
Females	138,158	5,979	10,128	12,020	10,156	10,667	16,951	24,384	47,873
Acute Lymphocytic Leuk									
Males	41,055	5,618	11,450	9,948	6,685	4,256	1,672	861	564
Females	34,121	5,033	8,635	8,663	5,668	3,311	1,318	916	579

^a U.S. 2012 cancer prevalence counts are based on 2012 cancer prevalence proportions from the SEER 9 registries (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta) and 1/1/2012 U.S. population estimates based on the average of 2011 and 2012 population estimates from the U.S. Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^b Cases diagnosed more than 37 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^c Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.

Table 2.21
All Cancer Sites (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2012
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <20 ^e	0 to <37 ^e	>=37 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	4,660,390	3,220,696	2,297,725	1,454,626	861,141	505,137	12,037,276	13,405,317	370,934	13,776,251
	Males	2,395,057	1,636,053	1,105,506	654,928	315,984	163,447	5,951,188	6,387,558	76,971	6,464,529
	Females	2,265,333	1,584,643	1,192,219	799,698	545,157	341,690	6,086,088	7,017,759	293,963	7,311,722
White ^b	Both Sexes	3,913,280	2,780,192	2,013,548	1,287,998	779,233	458,563	10,359,308	11,601,795	324,151	11,925,946
	Males	2,003,404	1,405,324	962,849	578,882	289,312	150,214	5,096,308	5,497,887	72,025	5,569,912
	Females	1,909,876	1,374,868	1,050,699	709,116	489,921	308,349	5,263,000	6,103,908	252,126	6,356,034
Black ^b	Both Sexes	487,189	296,165	190,445	110,608	52,052	29,576	1,109,887	1,189,165	22,527	1,211,692
	Males	261,563	162,329	101,649	53,718	17,259	7,999	588,391	609,432	1,907	611,339
	Females	225,626	133,836	88,796	56,890	34,793	21,577	521,496	579,733	20,620	600,353
Asian/ Pacific Islander ^c	Both Sexes	147,701	90,366	58,598	+	+	+	330,690	+	+	+
	Males	62,926	37,188	22,375	+	+	+	134,340	+	+	+
	Females	84,775	53,178	36,223	+	+	+	196,350	+	+	+
Hispanic ^d	Both Sexes	327,526	204,575	131,420	+	+	+	742,005	+	+	+
	Males	155,663	95,961	59,027	+	+	+	342,644	+	+	+
	Females	171,863	108,614	72,393	+	+	+	399,361	+	+	+

Estimated prevalence percent^a on January 1, 2012, of the SEER^c population diagnosed in the previous 20 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted ^f	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	3.3439%	0.0867%	0.1870%	0.3323%	0.7825%	1.9172%	4.3781%	9.4329%	15.9290%	17.8982%	3.2158%
	Males	3.3276%	0.0907%	0.1973%	0.3215%	0.6175%	1.3317%	3.5697%	10.0706%	20.3717%	25.5905%	3.5875%
	Females	3.3597%	0.0826%	0.1762%	0.3436%	0.9482%	2.4979%	5.1523%	8.8564%	12.3280%	13.3496%	3.0004%
White ^c	Both Sexes	3.6851%	0.0946%	0.2067%	0.3762%	0.8788%	2.0546%	4.5798%	9.8605%	16.7896%	18.5683%	3.3871%
	Males	3.6345%	0.0977%	0.2199%	0.3648%	0.7023%	1.4504%	3.7052%	10.3218%	21.0546%	26.0188%	3.7143%
	Females	3.7355%	0.0914%	0.1929%	0.3884%	1.0646%	2.6790%	5.4452%	9.4296%	13.2452%	14.1132%	3.2085%
Black ^c	Both Sexes	2.5129%	0.0631%	0.1280%	0.2137%	0.5347%	1.5799%	4.1368%	9.8340%	15.9616%	16.0571%	3.0533%
	Males	2.7904%	0.0677%	0.1266%	0.2103%	0.3830%	1.1080%	4.0489%	12.8776%	24.1031%	27.6383%	4.0581%
	Females	2.2601%	0.0582%	0.1294%	0.2170%	0.6672%	1.9962%	4.2134%	7.4455%	10.2639%	10.4330%	2.4333%
Asian/ Pacific Islander ^c	Both Sexes	2.1407%	0.0676%	0.1360%	0.2033%	0.4844%	1.3780%	3.1398%	5.8799%	9.8282%	12.7380%	2.1271%
	Males	1.8597%	0.0748%	0.1430%	0.1837%	0.3286%	0.7281%	1.9350%	5.2145%	11.7196%	17.9383%	2.1145%
	Females	2.3981%	0.0600%	0.1287%	0.2229%	0.6243%	1.9531%	4.1756%	6.4300%	8.3655%	9.5615%	2.1951%
Hispanic ^d	Both Sexes	1.4540%	0.0814%	0.1769%	0.2627%	0.5666%	1.3533%	3.1080%	6.6182%	11.5132%	13.4762%	2.3397%
	Males	1.3163%	0.0864%	0.1864%	0.2626%	0.4401%	0.8479%	2.2370%	6.7085%	15.3104%	20.1082%	2.6149%
	Females	1.5950%	0.0763%	0.1670%	0.2628%	0.7018%	1.8807%	3.9425%	6.5406%	8.7002%	9.5071%	2.2195%

^a US 2012 cancer prevalence counts are based on 2012 cancer prevalence proportions from the SEER registries and 1/1/2012 US population estimates based on the average of 2011 and 2012 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 37 years for 1975-2012 SEER 9 data; 20 years for 1992-2012 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 37 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <37 and >=37. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 3.21
Brain and Other Nervous System (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2012
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <20 ^e	0 to <37 ^e	>=37 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	45,479	26,552	20,519	16,125	12,531	7,759	114,403	136,100	12,718	148,818
	Males	24,643	14,047	10,387	8,833	6,511	4,276	61,047	72,270	5,786	78,056
	Females	20,836	12,505	10,132	7,292	6,020	3,483	53,356	63,830	6,932	70,762
White ^b	Both Sexes	39,324	23,138	18,018	14,010	11,024	6,962	99,442	118,867	4,142	123,009
	Males	21,480	12,479	9,147	7,710	5,655	3,801	53,485	63,520	2,624 ⁱ	66,144
	Females	17,844	10,659	8,871	6,300	5,369	3,161	45,957	55,347	1,518 ⁱ	56,865
Black ^b	Both Sexes	3,696	2,064	1,607	1,273	1,024	573	9,154	10,816	+	+
	Males	1,760	897	854	653	582	357	4,486	5,342	+	+
	Females	1,936	1,167	753	620	442	216	4,668	5,474	+	+
Asian/ Pacific Islander ^c	Both Sexes	1,487	812	532	+	+	+	3,303	+	+	+
	Males	803	374	248	+	+	+	1,668	+	+	+
	Females	684	438	284	+	+	+	1,635	+	+	+
Hispanic ^d	Both Sexes	4,903	3,063	2,275	+	+	+	11,684	+	+	+
	Males	2,674	1,632	1,176	+	+	+	6,187	+	+	+
	Females	2,229	1,431	1,099	+	+	+	5,497	+	+	+

Estimated prevalence percent^a on January 1, 2012, of the SEER^c population diagnosed in the previous 20 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted ^f	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0318%	0.0146%	0.0328%	0.0331%	0.0318%	0.0370%	0.0395%	0.0393%	0.0308%	0.0177%	0.0314%
	Males	0.0344%	0.0152%	0.0338%	0.0348%	0.0350%	0.0404%	0.0433%	0.0459%	0.0309%	0.0214%	0.0338%
	Females	0.0292%	0.0140%	0.0316%	0.0312%	0.0286%	0.0336%	0.0359%	0.0333%	0.0307%	0.0156%	0.0289%
White ^c	Both Sexes	0.0360%	0.0164%	0.0366%	0.0377%	0.0365%	0.0423%	0.0454%	0.0440%	0.0342%	0.0174%	0.0355%
	Males	0.0391%	0.0169%	0.0383%	0.0398%	0.0393%	0.0466%	0.0492%	0.0513%	0.0353%	0.0205%	0.0383%
	Females	0.0330%	0.0160%	0.0348%	0.0355%	0.0334%	0.0378%	0.0417%	0.0371%	0.0333%	0.0156%	0.0327%
Black ^c	Both Sexes	0.0203%	0.0094%	0.0237%	0.0234%	0.0197%	0.0222%	0.0210%	0.0278%	0.0172%	0.0161%	0.0201%
	Males	0.0203%	0.0097%	0.0225%	0.0239%	0.0207%	0.0211%	0.0233%	0.0289%	0.0101%	0.0185%	0.0200%
	Females	0.0203%	0.0091%	0.0249%	0.0230%	0.0188%	0.0231%	0.0189%	0.0269%	0.0222%	0.0149%	0.0202%
Asian/ Pacific Islander ^c	Both Sexes	0.0185%	0.0096%	0.0202%	0.0187%	0.0189%	0.0213%	0.0212%	0.0189%	0.0203%	0.0169%	0.0184%
	Males	0.0196%	0.0101%	0.0206%	0.0189%	0.0223%	0.0210%	0.0226%	0.0217%	0.0174%	0.0240%	0.0195%
	Females	0.0176%	0.0090%	0.0198%	0.0184%	0.0157%	0.0216%	0.0199%	0.0167%	0.0225%	0.0125%	0.0175%
Hispanic ^d	Both Sexes	0.0225%	0.0112%	0.0252%	0.0198%	0.0207%	0.0287%	0.0327%	0.0331%	0.0247%	0.0227%	0.0237%
	Males	0.0235%	0.0115%	0.0260%	0.0191%	0.0214%	0.0312%	0.0354%	0.0397%	0.0219%	0.0309%	0.0251%
	Females	0.0215%	0.0108%	0.0243%	0.0205%	0.0200%	0.0262%	0.0301%	0.0275%	0.0268%	0.0179%	0.0224%

^a US 2012 cancer prevalence counts are based on 2012 cancer prevalence proportions from the SEER registries and 1/1/2012 US population estimates based on the average of 2011 and 2012 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 37 years for 1975-2012 SEER 9 data; 20 years for 1992-2012 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 37 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <37 and >=37. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 4.24
Cancer of the Breast (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2012
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <20 ^e	0 to <37 ^e	>=37 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	878,837	683,310	557,489	364,603	237,790	129,315	2,592,173	2,943,928	46,885	2,990,813
	Males	6,296	3,718	2,470	1,306	799	439	14,190	15,382	117	15,499
	Females	872,541	679,592	555,019	363,297	236,991	128,876	2,577,983	2,928,546	46,768	2,975,314
White ^b	Both Sexes	737,273	592,066	492,299	324,516	214,831	116,791	2,242,861	2,561,799	41,301	2,603,100
	Males	5,175	3,215	2,089	1,155	689	394	11,988	13,049	0 ⁱ	13,049
	Females	732,098	588,851	490,210	323,361	214,142	116,397	2,230,873	2,548,750	41,301	2,590,051
Black ^b	Both Sexes	93,078	60,161	42,176	27,086	15,700	8,741	230,141	252,947	+	+
	Males	915	388	281	97	110	36	1,729	1,851	+	+
	Females	92,163	59,773	41,895	26,989	15,590	8,705	228,412	251,096	3,636	254,732
Asian/ Pacific Islander ^c	Both Sexes	35,590	24,237	17,511	+	+	+	87,126	+	+	+
	Males	145	98	39	+	+	+	322	+	+	+
	Females	35,445	24,139	17,472	+	+	+	86,804	+	+	+
Hispanic ^d	Both Sexes	61,983	42,773	29,054	+	+	+	151,046	+	+	+
	Males	296	179	78	+	+	+	580	+	+	+
	Females	61,687	42,594	28,976	+	+	+	150,466	+	+	+

Estimated prevalence percent^a on January 1, 2012, of the SEER^c population diagnosed in the previous 20 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted ^f	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.7268%	-	0.0001%	0.0067%	0.0852%	0.4759%	1.2094%	2.2244%	3.0789%	3.5556%	0.6832%
	Males	0.0080%	-	-	-	0.0006%	0.0030%	0.0084%	0.0266%	0.0478%	0.0683%	0.0086%
	Females	1.4276%	-	0.0003%	0.0136%	0.1702%	0.9449%	2.3596%	4.2110%	5.5356%	5.6177%	1.2548%
White ^c	Both Sexes	0.7971%	-	0.0001%	0.0065%	0.0835%	0.4748%	1.2416%	2.3391%	3.2974%	3.7883%	0.7149%
	Males	0.0086%	-	-	-	0.0005%	0.0025%	0.0077%	0.0277%	0.0508%	0.0733%	0.0088%
	Females	1.5829%	-	0.0003%	0.0133%	0.1709%	0.9628%	2.4625%	4.4982%	5.9954%	6.0098%	1.3295%
Black ^c	Both Sexes	0.5266%	-	-	0.0076%	0.0988%	0.4898%	1.1052%	1.9877%	2.6319%	3.0335%	0.6170%
	Males	0.0078%	-	-	-	-	0.0055%	0.0161%	0.0341%	0.0493%	0.0720%	0.0106%
	Females	0.9991%	-	-	0.0148%	0.1844%	0.9169%	2.0545%	3.5208%	4.4394%	4.4716%	1.0684%
Asian/ Pacific Islander ^c	Both Sexes	0.5562%	-	-	0.0067%	0.0828%	0.4764%	1.1203%	1.7076%	2.1247%	2.2900%	0.5370%
	Males	0.0044%	-	-	-	-	0.0025%	0.0058%	0.0129%	0.0275%	0.0329%	0.0049%
	Females	1.0616%	-	-	0.0132%	0.1563%	0.8958%	2.0786%	3.1083%	3.7465%	3.6687%	0.9632%
Hispanic ^d	Both Sexes	0.2989%	-	-	0.0045%	0.0623%	0.3287%	0.8821%	1.5670%	2.0875%	2.3259%	0.4763%
	Males	0.0022%	-	-	-	-	0.0017%	0.0043%	0.0153%	0.0247%	0.0327%	0.0045%
	Females	0.6026%	-	-	0.0093%	0.1286%	0.6700%	1.7232%	2.8993%	3.6157%	3.6983%	0.8667%

^a US 2012 cancer prevalence counts are based on 2012 cancer prevalence proportions from the SEER registries and 1/1/2012 US population estimates based on the average of 2011 and 2012 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 37 years for 1975-2012 SEER 9 data; 20 years for 1992-2012 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 37 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <37 and >=37. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 5.15
Cancer of the Cervix Uteri (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2012
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <20 ^e	0 to <37 ^e	>=37 ^g	Complete ^h
<u>Race</u>										
All Races ^b	39,323	32,593	34,878	32,772	28,429	21,631	151,928	214,008	35,504	249,512
White ^b	31,133	25,865	27,920	27,182	23,762	17,910	122,201	173,836	30,059	203,895
Black ^b	5,191	4,548	4,549	3,363	2,982	2,737	19,129	26,586	+	+
Asian/Pacific Islander ^c	2,104	1,727	1,678	+	+	+	7,030	+	+	+
Hispanic ^d	7,897	7,205	6,782	+	+	+	27,657	+	+	+

Estimated prevalence percent^a on January 1, 2012, of the SEER^c population diagnosed in the previous 20 years
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f All Ages	
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+		
<u>Race</u>												
All Races ^c	0.0953%	-	0.0002%	0.0087%	0.0723%	0.1825%	0.2039%	0.1846%	0.1608%	0.1031%	0.0896%	
White ^c	0.0996%	-	-	0.0101%	0.0830%	0.2026%	0.2119%	0.1762%	0.1457%	0.0853%	0.0934%	
Black ^c	0.0801%	-	-	0.0054%	0.0599%	0.1449%	0.1807%	0.1993%	0.1863%	0.1394%	0.0825%	
Asian/Pacific Islander ^c	0.0834%	-	-	0.0042%	0.0302%	0.1070%	0.1718%	0.2126%	0.2242%	0.1991%	0.0760%	
Hispanic ^d	0.1105%	-	-	0.0088%	0.0783%	0.2340%	0.3404%	0.3503%	0.3416%	0.2635%	0.1419%	

^a US 2012 cancer prevalence counts are based on 2012 cancer prevalence proportions from the SEER registries and 1/1/2012 US population estimates based on the average of 2011 and 2012 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 37 years for 1975-2012 SEER 9 data; 20 years for 1992-2012 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 37 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <37 and >=37. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 6.28
Cancer of the Colon and Rectum (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2012
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <20 ^e	0 to <37 ^e	>=37 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	398,737	281,172	199,881	118,333	77,219	45,038	1,033,951	1,149,814	19,115	1,168,929
	Males	204,170	140,415	100,896	58,365	37,777	20,275	521,903	573,302	6,599	579,901
	Females	194,567	140,757	98,985	59,968	39,442	24,763	512,048	576,512	12,516	589,028
White ^b	Both Sexes	324,290	236,784	171,603	101,740	67,233	39,748	865,509	967,504	16,749	984,253
	Males	167,409	120,619	87,557	50,622	33,369	17,992	442,016	487,849	5,865	493,714
	Females	156,881	116,165	84,046	51,118	33,864	21,756	423,493	479,655	10,884	490,539
Black ^b	Both Sexes	47,802	28,970	18,851	10,912	6,993	3,772	109,787	119,414	1,456	120,870
	Males	22,482	11,967	8,317	4,755	2,949	1,443	49,065	52,531	452	52,983
	Females	25,320	17,003	10,534	6,157	4,044	2,329	60,722	66,883	1,004	67,887
Asian/ Pacific Islander ^c	Both Sexes	18,283	11,130	6,965	+	+	+	40,533	+	+	+
	Males	9,532	5,669	3,500	+	+	+	20,817	+	+	+
	Females	8,751	5,461	3,465	+	+	+	19,716	+	+	+
Hispanic ^d	Both Sexes	31,428	17,484	10,149	+	+	+	64,639	+	+	+
	Males	16,954	9,073	5,339	+	+	+	34,188	+	+	+
	Females	14,474	8,411	4,810	+	+	+	30,451	+	+	+

Estimated prevalence percent^a on January 1, 2012, of the SEER^c population diagnosed in the previous 20 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	Age-Adjusted ^f All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.2958%	-	0.0004%	0.0051%	0.0263%	0.1096%	0.3496%	0.7612%	1.5234%	2.3653%	0.2845%
	Males	0.3032%	-	0.0004%	0.0048%	0.0265%	0.1172%	0.3837%	0.8917%	1.8042%	2.6637%	0.3268%
	Females	0.2886%	-	0.0004%	0.0055%	0.0260%	0.1021%	0.3169%	0.6432%	1.2958%	2.1888%	0.2507%
White ^c	Both Sexes	0.3079%	-	0.0004%	0.0057%	0.0271%	0.1077%	0.3325%	0.7303%	1.5176%	2.3849%	0.2804%
	Males	0.3158%	-	0.0003%	0.0052%	0.0268%	0.1140%	0.3684%	0.8607%	1.8043%	2.6787%	0.3227%
	Females	0.3000%	-	0.0005%	0.0063%	0.0275%	0.1011%	0.2969%	0.6085%	1.2793%	2.2093%	0.2456%
Black ^c	Both Sexes	0.2516%	-	-	0.0035%	0.0226%	0.1193%	0.4202%	0.9605%	1.6861%	2.2680%	0.3153%
	Males	0.2384%	-	-	0.0043%	0.0225%	0.1213%	0.4308%	1.0773%	1.8373%	2.4198%	0.3410%
	Females	0.2636%	-	-	0.0028%	0.0226%	0.1177%	0.4109%	0.8688%	1.5802%	2.1943%	0.2975%
Asian/ Pacific Islander ^c	Both Sexes	0.2724%	-	-	0.0026%	0.0231%	0.1051%	0.3620%	0.7564%	1.4121%	2.2434%	0.2738%
	Males	0.2889%	-	-	0.0021%	0.0249%	0.1239%	0.4029%	0.8950%	1.7160%	2.6145%	0.3231%
	Females	0.2574%	-	-	0.0031%	0.0214%	0.0884%	0.3269%	0.6418%	1.1771%	2.0167%	0.2366%
Hispanic ^d	Both Sexes	0.1270%	-	-	0.0032%	0.0204%	0.0805%	0.2775%	0.6417%	1.2106%	1.7552%	0.2253%
	Males	0.1318%	-	-	0.0031%	0.0214%	0.0848%	0.2982%	0.7645%	1.5577%	2.1268%	0.2714%
	Females	0.1222%	-	-	0.0032%	0.0192%	0.0760%	0.2576%	0.5362%	0.9535%	1.5327%	0.1911%

^a US 2012 cancer prevalence counts are based on 2012 cancer prevalence proportions from the SEER registries and 1/1/2012 US population estimates based on the average of 2011 and 2012 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 37 years for 1975-2012 SEER 9 data; 20 years for 1992-2012 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 37 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <37 and >=37. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 7.15
Cancer of the Corpus and Uterus, NOS (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2012
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <20 ^e	0 to <37 ^e	>=37 ^g	Complete ^h
<u>Race</u>										
All Races ^b	183,434	127,357	101,371	70,925	47,636	32,609	504,608	598,766	22,846	621,612
White ^b	156,395	112,816	92,247	65,091	44,433	30,759	446,663	535,109	19,170	554,279
Black ^b	16,207	7,915	4,477	2,990	1,678	957	32,295	35,198	730	35,928
Asian/Pacific Islander ^c	7,132	4,407	2,976	+	+	+	16,394	+	+	+
Hispanic ^d	13,533	8,381	5,340	+	+	+	30,687	+	+	+

Estimated prevalence percent^a on January 1, 2012, of the SEER^c population diagnosed in the previous 20 years
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f All Ages
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	
<u>Race</u>											
All Races ^c	0.2740%	-	-	0.0033%	0.0305%	0.1177%	0.3935%	0.9253%	1.2039%	1.0375%	0.2397%
White ^c	0.3120%	-	-	0.0029%	0.0300%	0.1186%	0.4121%	1.0052%	1.3507%	1.1596%	0.2602%
Black ^c	0.1417%	-	-	0.0040%	0.0240%	0.0808%	0.2209%	0.6058%	0.8518%	0.5672%	0.1552%
Asian/Pacific Islander ^c	0.1984%	-	-	0.0040%	0.0326%	0.1348%	0.4207%	0.7006%	0.6230%	0.4936%	0.1766%
Hispanic ^d	0.1228%	-	-	0.0041%	0.0387%	0.1270%	0.3167%	0.6654%	0.7679%	0.5984%	0.1754%

^a US 2012 cancer prevalence counts are based on 2012 cancer prevalence proportions from the SEER registries and 1/1/2012 US population estimates based on the average of 2011 and 2012 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 37 years for 1975-2012 SEER 9 data; 20 years for 1992-2012 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 37 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <37 and >=37. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 8.21
Cancer of the Esophagus (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2012
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <20 ^e	0 to <37 ^e	>=37 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	21,344	7,819	4,025	1,560	584	237	35,130	35,667	114	35,781
	Males	16,607	6,020	3,180	1,254	420	185	27,341	27,750	58	27,808
	Females	4,737	1,799	845	306	164	52	7,789	7,917	56	7,973
White ^b	Both Sexes	19,188	7,012	3,600	1,367	519	206	31,500	31,990	89	32,079
	Males	15,216	5,539	2,926	1,105	377	157	25,036	25,405	52	25,457
	Females	3,972	1,473	674	262	142	49	6,464	6,585	37	6,622
Black ^b	Both Sexes	1,544	591	298	147	51	28	2,619	2,659	27	2,686
	Males	930	320	165	111	38	28	1,551	1,591	7	1,598
	Females	614	271	133	36	13	0	1,068	1,068	20	1,088
Asian/ Pacific Islander ^c	Both Sexes	431	123	89	+	+	+	676	+	+	+
	Males	316	84	66	+	+	+	489	+	+	+
	Females	115	39	23	+	+	+	187	+	+	+
Hispanic ^d	Both Sexes	974	247	123	+	+	+	1,430	+	+	+
	Males	794	207	106	+	+	+	1,175	+	+	+
	Females	180	40	17	+	+	+	255	+	+	+

Estimated prevalence percent^a on January 1, 2012, of the SEER^c population diagnosed in the previous 20 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	Age-Adjusted ^f All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0092%	-	-	0.0001%	0.0004%	0.0025%	0.0118%	0.0330%	0.0517%	0.0458%	0.0088%
	Males	0.0144%	-	-	-	0.0005%	0.0041%	0.0189%	0.0571%	0.0873%	0.0840%	0.0150%
	Females	0.0042%	-	-	-	0.0002%	0.0009%	0.0050%	0.0111%	0.0229%	0.0232%	0.0037%
White ^c	Both Sexes	0.0108%	-	-	-	0.0005%	0.0029%	0.0130%	0.0379%	0.0563%	0.0488%	0.0097%
	Males	0.0171%	-	-	-	0.0007%	0.0050%	0.0208%	0.0658%	0.0968%	0.0890%	0.0167%
	Females	0.0045%	-	-	-	-	0.0007%	0.0052%	0.0118%	0.0227%	0.0247%	0.0038%
Black ^c	Both Sexes	0.0057%	-	-	-	-	0.0009%	0.0123%	0.0238%	0.0423%	0.0322%	0.0069%
	Males	0.0072%	-	-	-	-	-	0.0168%	0.0316%	0.0574%	0.0646%	0.0100%
	Females	0.0043%	-	-	-	-	-	0.0083%	0.0177%	0.0317%	0.0164%	0.0048%
Asian/ Pacific Islander ^c	Both Sexes	0.0045%	-	-	-	-	0.0018%	0.0055%	0.0115%	0.0315%	0.0314%	0.0047%
	Males	0.0068%	-	-	-	-	0.0019%	0.0095%	0.0211%	0.0496%	0.0553%	0.0077%
	Females	0.0024%	-	-	-	-	0.0018%	0.0021%	0.0035%	0.0174%	0.0169%	0.0024%
Hispanic ^d	Both Sexes	0.0028%	-	-	-	0.0004%	0.0019%	0.0063%	0.0149%	0.0275%	0.0329%	0.0049%
	Males	0.0045%	-	-	-	0.0007%	0.0032%	0.0110%	0.0261%	0.0548%	0.0624%	0.0091%
	Females	0.0010%	-	-	-	-	-	0.0018%	0.0052%	0.0073%	0.0152%	0.0016%

^a US 2012 cancer prevalence counts are based on 2012 cancer prevalence proportions from the SEER registries and 1/1/2012 US population estimates based on the average of 2011 and 2012 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 37 years for 1975-2012 SEER 9 data; 20 years for 1992-2012 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 37 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <37 and >=37. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 9.21
Hodgkin LymphomaEstimated United States Cancer Prevalence Counts^a on January 1, 2012
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <20 ^e	0 to <37 ^e	>=37 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	38,167	33,462	28,363	24,965	21,696	16,536	134,051	176,990	12,636	189,626
	Males	20,781	17,382	14,349	12,735	10,761	8,636	70,014	91,797	6,285	98,082
	Females	17,386	16,080	14,014	12,230	10,935	7,900	64,037	85,193	6,351	91,544
White ^b	Both Sexes	31,580	28,310	24,730	22,151	19,990	15,371	115,070	155,246	13,975	169,221
	Males	17,279	14,894	12,469	11,314	9,961	7,826	60,338	80,396	7,224	87,620
	Females	14,301	13,416	12,261	10,837	10,029	7,545	54,732	74,850	6,751	81,601
Black ^b	Both Sexes	4,902	3,930	2,839	2,320	1,338	954	14,649	16,783	1,005	17,788
	Males	2,648	1,806	1,440	1,197	612	702	7,405	8,790	708	9,498
	Females	2,254	2,124	1,399	1,123	726	252	7,244	7,993	297	8,290
Asian/ Pacific Islander ^c	Both Sexes	1,010	798	490	+	+	+	2,590	+	+	+
	Males	515	404	266	+	+	+	1,300	+	+	+
	Females	495	394	224	+	+	+	1,290	+	+	+
Hispanic ^d	Both Sexes	4,206	3,162	2,559	+	+	+	11,603	+	+	+
	Males	2,220	1,767	1,415	+	+	+	6,319	+	+	+
	Females	1,986	1,395	1,144	+	+	+	5,284	+	+	+

Estimated prevalence percent^a on January 1, 2012, of the SEER^c population diagnosed in the previous 20 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted ^f	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0365%	0.0007%	0.0103%	0.0396%	0.0643%	0.0624%	0.0440%	0.0346%	0.0326%	0.0255%	0.0364%
	Males	0.0387%	0.0011%	0.0115%	0.0390%	0.0624%	0.0661%	0.0506%	0.0421%	0.0391%	0.0294%	0.0386%
	Females	0.0343%	0.0003%	0.0091%	0.0403%	0.0663%	0.0587%	0.0376%	0.0279%	0.0273%	0.0232%	0.0344%
White ^c	Both Sexes	0.0413%	0.0007%	0.0112%	0.0446%	0.0747%	0.0708%	0.0495%	0.0388%	0.0369%	0.0288%	0.0414%
	Males	0.0437%	0.0010%	0.0125%	0.0431%	0.0718%	0.0744%	0.0560%	0.0473%	0.0436%	0.0332%	0.0434%
	Females	0.0390%	0.0004%	0.0098%	0.0462%	0.0779%	0.0671%	0.0431%	0.0308%	0.0314%	0.0262%	0.0395%
Black ^c	Both Sexes	0.0330%	0.0007%	0.0100%	0.0309%	0.0598%	0.0630%	0.0437%	0.0337%	0.0297%	0.0176%	0.0341%
	Males	0.0347%	0.0013%	0.0100%	0.0312%	0.0581%	0.0664%	0.0559%	0.0349%	0.0415%	0.0215%	0.0366%
	Females	0.0315%	-	0.0101%	0.0306%	0.0613%	0.0600%	0.0332%	0.0327%	0.0214%	0.0157%	0.0319%
Asian/ Pacific Islander ^c	Both Sexes	0.0143%	0.0010%	0.0050%	0.0231%	0.0233%	0.0205%	0.0132%	0.0108%	0.0105%	0.0075%	0.0137%
	Males	0.0150%	0.0014%	0.0067%	0.0254%	0.0226%	0.0203%	0.0126%	0.0134%	0.0128%	0.0103%	0.0145%
	Females	0.0137%	-	0.0032%	0.0209%	0.0239%	0.0206%	0.0138%	0.0087%	0.0087%	0.0058%	0.0129%
Hispanic ^d	Both Sexes	0.0225%	0.0010%	0.0095%	0.0270%	0.0364%	0.0343%	0.0304%	0.0337%	0.0389%	0.0228%	0.0247%
	Males	0.0242%	0.0013%	0.0117%	0.0278%	0.0345%	0.0362%	0.0399%	0.0360%	0.0423%	0.0412%	0.0272%
	Females	0.0208%	0.0007%	0.0072%	0.0261%	0.0385%	0.0323%	0.0212%	0.0317%	0.0363%	0.0118%	0.0226%

^a US 2012 cancer prevalence counts are based on 2012 cancer prevalence proportions from the SEER registries and 1/1/2012 US population estimates based on the average of 2011 and 2012 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 37 years for 1975-2012 SEER 9 data; 20 years for 1992-2012 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 37 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <37 and >=37. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 10.15
Kaposi SarcomaEstimated United States Cancer Prevalence Counts^a on January 1, 2012
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <20 ^e	0 to <37 ^e	>=37 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	7,384	5,524	4,521	6,509	1,589	579	24,738	26,300	+	+
	Males	6,836	5,165	4,277	6,367	1,489	545	23,345	24,829	+	+
	Females	548	359	244	142	100	34	1,393	1,471	+	+
White ^b	Both Sexes	4,557	4,106	3,235	5,325	1,272	495	17,877	19,162	+	+
	Males	4,177	3,841	3,037	5,229	1,193	472	16,858	18,077	+	+
	Females	380	265	198	96	79	23	1,019	1,085	+	+
Black ^b	Both Sexes	2,359	1,214	990	804	144	38	5,438	5,560	+	+
	Males	2,238	1,138	978	769	134	38	5,185	5,307	+	+
	Females	121	76	12	35	10	0	253	253	+	+
Asian/ Pacific Islander ^c	Both Sexes	134	89	62	+	+	+	359	+	+	+
	Males	134	84	62	+	+	+	354	+	+	+
	Females	0	5	0	+	+	+	5	+	+	+
Hispanic ^d	Both Sexes	1,259	1,080	926	+	+	+	4,360	+	+	+
	Males	1,115	1,019	875	+	+	+	4,083	+	+	+
	Females	144	61	51	+	+	+	277	+	+	+

Estimated prevalence percent^a on January 1, 2012, of the SEER^c population diagnosed in the previous 20 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	Age-Adjusted ^f All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0085%	-	-	0.0012%	0.0054%	0.0192%	0.0207%	0.0121%	0.0095%	0.0152%	0.0082%
	Males	0.0165%	-	-	0.0022%	0.0106%	0.0377%	0.0416%	0.0245%	0.0179%	0.0288%	0.0161%
	Females	0.0008%	-	-	-	0.0002%	0.0008%	0.0006%	0.0009%	0.0027%	0.0072%	0.0007%
White ^c	Both Sexes	0.0086%	-	-	0.0006%	0.0046%	0.0182%	0.0212%	0.0127%	0.0104%	0.0161%	0.0080%
	Males	0.0164%	-	-	0.0012%	0.0089%	0.0350%	0.0422%	0.0253%	0.0191%	0.0301%	0.0155%
	Females	0.0009%	-	-	-	-	0.0008%	0.0005%	0.0009%	0.0031%	0.0078%	0.0007%
Black ^c	Both Sexes	0.0128%	-	-	0.0046%	0.0147%	0.0358%	0.0284%	0.0133%	0.0066%	0.0124%	0.0133%
	Males	0.0259%	-	-	0.0091%	0.0309%	0.0751%	0.0589%	0.0284%	0.0137%	0.0215%	0.0277%
	Females	0.0009%	-	-	-	-	-	0.0018%	-	-	0.0080%	0.0009%
Asian/ Pacific Islander ^c	Both Sexes	0.0021%	-	-	-	0.0011%	0.0048%	0.0049%	0.0032%	0.0032%	0.0045%	0.0020%
	Males	0.0044%	-	-	-	0.0023%	0.0099%	0.0105%	0.0070%	0.0074%	0.0103%	0.0044%
	Females	-	-	-	-	-	-	-	-	-	-	-
Hispanic ^d	Both Sexes	0.0085%	-	-	0.0009%	0.0065%	0.0233%	0.0256%	0.0177%	0.0168%	0.0428%	0.0113%
	Males	0.0158%	-	-	0.0016%	0.0126%	0.0441%	0.0514%	0.0361%	0.0291%	0.0693%	0.0214%
	Females	0.0011%	-	-	-	-	0.0016%	-	0.0019%	0.0078%	0.0270%	0.0019%

^a US 2012 cancer prevalence counts are based on 2012 cancer prevalence proportions from the SEER registries and 1/1/2012 US population estimates based on the average of 2011 and 2012 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 37 years for 1975-2012 SEER 9 data; 20 years for 1992-2012 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 37 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <37 and >=37. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 11.21
Cancer of the Kidney And Renal Pelvis (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2012
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <20 ^e	0 to <37 ^e	>=37 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	154,353	93,811	51,100	31,248	18,187	9,960	339,047	365,660	10,265	375,925
	Males	94,867	55,938	29,534	18,295	10,017	5,667	203,620	217,927	5,756	223,683
	Females	59,486	37,873	21,566	12,953	8,170	4,293	135,427	147,733	4,509	152,242
White ^b	Both Sexes	127,833	80,333	43,767	27,102	16,237	8,794	286,582	310,261	11,564	321,825
	Males	79,042	48,043	25,733	16,331	9,050	5,094	173,577	186,613	5,093	191,706
	Females	48,791	32,290	18,034	10,771	7,187	3,700	113,005	123,648	6,471	130,119
Black ^b	Both Sexes	18,611	9,480	5,438	3,156	1,524	938	37,425	39,819	2,023	41,842
	Males	10,872	5,572	2,698	1,432	750	432	21,002	22,005	689	22,694
	Females	7,739	3,908	2,740	1,724	774	506	16,423	17,814	1,334	19,148
Asian/ Pacific Islander ^c	Both Sexes	4,049	2,099	1,081	+	+	+	7,755	+	+	+
	Males	2,508	1,272	643	+	+	+	4,743	+	+	+
	Females	1,541	827	438	+	+	+	3,012	+	+	+
Hispanic ^d	Both Sexes	14,872	7,921	4,151	+	+	+	29,087	+	+	+
	Males	8,530	4,456	2,294	+	+	+	16,449	+	+	+
	Females	6,342	3,465	1,857	+	+	+	12,638	+	+	+

Estimated prevalence percent^a on January 1, 2012, of the SEER^c population diagnosed in the previous 20 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted ^f	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0942%	0.0073%	0.0107%	0.0059%	0.0150%	0.0572%	0.1417%	0.2885%	0.4291%	0.3900%	0.0897%
	Males	0.1156%	0.0068%	0.0096%	0.0054%	0.0162%	0.0712%	0.1839%	0.3902%	0.5855%	0.5452%	0.1181%
	Females	0.0734%	0.0078%	0.0119%	0.0064%	0.0138%	0.0434%	0.1014%	0.1967%	0.3023%	0.2982%	0.0660%
White ^c	Both Sexes	0.1038%	0.0076%	0.0115%	0.0063%	0.0156%	0.0612%	0.1486%	0.3034%	0.4595%	0.4058%	0.0947%
	Males	0.1270%	0.0069%	0.0109%	0.0057%	0.0161%	0.0759%	0.1900%	0.4051%	0.6256%	0.5734%	0.1241%
	Females	0.0808%	0.0082%	0.0122%	0.0069%	0.0150%	0.0460%	0.1076%	0.2084%	0.3214%	0.3056%	0.0697%
Black ^c	Both Sexes	0.0830%	0.0089%	0.0124%	0.0085%	0.0164%	0.0606%	0.1604%	0.3225%	0.4353%	0.3802%	0.0959%
	Males	0.0988%	0.0101%	0.0085%	0.0072%	0.0187%	0.0744%	0.2159%	0.4464%	0.5737%	0.4852%	0.1243%
	Females	0.0686%	0.0077%	0.0164%	0.0097%	0.0144%	0.0485%	0.1121%	0.2252%	0.3385%	0.3292%	0.0746%
Asian/ Pacific Islander ^c	Both Sexes	0.0501%	0.0040%	0.0040%	0.0017%	0.0078%	0.0300%	0.0756%	0.1593%	0.2211%	0.2720%	0.0495%
	Males	0.0637%	0.0025%	0.0033%	0.0023%	0.0093%	0.0367%	0.1063%	0.2310%	0.3114%	0.3728%	0.0679%
	Females	0.0377%	0.0057%	0.0046%	0.0012%	0.0065%	0.0240%	0.0493%	0.0999%	0.1513%	0.2104%	0.0352%
Hispanic ^d	Both Sexes	0.0570%	0.0060%	0.0087%	0.0037%	0.0150%	0.0549%	0.1353%	0.2792%	0.4475%	0.4576%	0.0910%
	Males	0.0633%	0.0051%	0.0072%	0.0026%	0.0141%	0.0630%	0.1622%	0.3609%	0.6133%	0.5830%	0.1147%
	Females	0.0505%	0.0069%	0.0102%	0.0050%	0.0159%	0.0465%	0.1095%	0.2091%	0.3248%	0.3826%	0.0724%

^a US 2012 cancer prevalence counts are based on 2012 cancer prevalence proportions from the SEER registries and 1/1/2012 US population estimates based on the average of 2011 and 2012 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 37 years for 1975-2012 SEER 9 data; 20 years for 1992-2012 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 37 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <37 and >=37. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 12.21
Cancer of the Larynx (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2012
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <20 ^e	0 to <37 ^e	>=37 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	31,457	19,544	14,348	9,341	6,347	3,986	77,548	87,479	1,373	88,852
	Males	25,542	15,526	11,604	7,487	5,002	3,235	62,341	70,303	1,111	71,414
	Females	5,915	4,018	2,744	1,854	1,345	751	15,207	17,176	262	17,438
White ^b	Both Sexes	26,218	16,460	12,312	8,091	5,536	3,494	65,593	74,257	1,065	75,322
	Males	21,285	13,075	10,033	6,488	4,424	2,887	52,786	59,838	859	60,697
	Females	4,933	3,385	2,279	1,603	1,112	607	12,807	14,419	206	14,625
Black ^b	Both Sexes	4,102	2,598	1,625	1,043	633	409	9,648	10,641	100	10,741
	Males	3,272	2,045	1,199	842	477	292	7,583	8,322	85	8,407
	Females	830	553	426	201	156	117	2,065	2,319	15	2,334
Asian/ Pacific Islander ^c	Both Sexes	591	353	257	+	+	+	1,338	+	+	+
	Males	510	308	222	+	+	+	1,160	+	+	+
	Females	81	45	35	+	+	+	178	+	+	+
Hispanic ^d	Both Sexes	1,883	1,039	812	+	+	+	4,186	+	+	+
	Males	1,623	875	655	+	+	+	3,515	+	+	+
	Females	260	164	157	+	+	+	671	+	+	+

Estimated prevalence percent^a on January 1, 2012, of the SEER^c population diagnosed in the previous 20 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	Age-Adjusted ^f All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0203%	-	-	0.0003%	0.0008%	0.0048%	0.0265%	0.0706%	0.1197%	0.0979%	0.0194%
	Males	0.0336%	-	-	0.0004%	0.0010%	0.0067%	0.0416%	0.1236%	0.2244%	0.2174%	0.0355%
	Females	0.0074%	-	-	0.0002%	0.0007%	0.0030%	0.0120%	0.0227%	0.0349%	0.0273%	0.0066%
White ^c	Both Sexes	0.0224%	-	-	0.0003%	0.0009%	0.0056%	0.0281%	0.0735%	0.1252%	0.1003%	0.0203%
	Males	0.0366%	-	-	0.0004%	0.0010%	0.0075%	0.0438%	0.1262%	0.2319%	0.2201%	0.0365%
	Females	0.0083%	-	-	-	0.0009%	0.0036%	0.0126%	0.0242%	0.0366%	0.0287%	0.0070%
Black ^c	Both Sexes	0.0215%	-	-	-	0.0009%	0.0041%	0.0380%	0.1027%	0.1673%	0.1120%	0.0263%
	Males	0.0355%	-	-	-	-	0.0059%	0.0571%	0.1910%	0.3330%	0.2750%	0.0511%
	Females	0.0088%	-	-	-	-	0.0025%	0.0213%	0.0335%	0.0514%	0.0329%	0.0093%
Asian/ Pacific Islander ^c	Both Sexes	0.0090%	-	-	-	-	0.0016%	0.0081%	0.0301%	0.0559%	0.0730%	0.0093%
	Males	0.0164%	-	-	-	-	0.0030%	0.0152%	0.0588%	0.1086%	0.1708%	0.0188%
	Females	0.0023%	-	-	-	-	-	0.0020%	0.0064%	0.0152%	0.0132%	0.0022%
Hispanic ^d	Both Sexes	0.0082%	-	-	-	0.0006%	0.0036%	0.0158%	0.0448%	0.0991%	0.0973%	0.0149%
	Males	0.0135%	-	-	-	-	0.0050%	0.0261%	0.0838%	0.2049%	0.2166%	0.0295%
	Females	0.0027%	-	-	-	0.0008%	0.0022%	0.0059%	0.0113%	0.0207%	0.0258%	0.0041%

^a US 2012 cancer prevalence counts are based on 2012 cancer prevalence proportions from the SEER registries and 1/1/2012 US population estimates based on the average of 2011 and 2012 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 37 years for 1975-2012 SEER 9 data; 20 years for 1992-2012 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 37 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <37 and >=37. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 13.29
LeukemiaEstimated United States Cancer Prevalence Counts^a on January 1, 2012
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <20 ^e	0 to <37 ^e	>=37 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	120,252	77,467	46,472	30,150	18,163	11,594	282,475	312,923	5,466	318,389
	Males	70,027	43,970	26,212	17,070	9,951	6,383	161,521	177,810	2,421	180,231
	Females	50,225	33,497	20,260	13,080	8,212	5,211	120,954	135,113	3,045	138,158
White ^b	Both Sexes	103,680	68,433	41,514	26,850	16,228	10,578	247,816	275,376	4,574	279,950
	Males	60,589	39,019	23,441	15,284	9,046	5,895	142,242	157,183	1,698	158,881
	Females	43,091	29,414	18,073	11,566	7,182	4,683	105,574	118,193	2,876	121,069
Black ^b	Both Sexes	9,177	5,387	2,761	1,771	1,028	500	19,539	21,066	775	21,841
	Males	5,147	3,002	1,366	915	455	257	10,612	11,290	358	11,648
	Females	4,030	2,385	1,395	856	573	243	8,927	9,776	417	10,193
Asian/ Pacific Islander ^c	Both Sexes	2,924	1,989	1,213	+	+	+	7,090	+	+	+
	Males	1,720	1,063	712	+	+	+	4,006	+	+	+
	Females	1,204	926	501	+	+	+	3,084	+	+	+
Hispanic ^d	Both Sexes	11,633	7,482	4,749	+	+	+	27,364	+	+	+
	Males	6,637	4,246	2,603	+	+	+	15,540	+	+	+
	Females	4,996	3,236	2,146	+	+	+	11,824	+	+	+

Estimated prevalence percent^a on January 1, 2012, of the SEER^c population diagnosed in the previous 20 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted ^f	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0800%	0.0323%	0.0605%	0.0424%	0.0275%	0.0430%	0.0818%	0.1649%	0.2784%	0.3306%	0.0785%
	Males	0.0932%	0.0339%	0.0666%	0.0458%	0.0317%	0.0511%	0.1010%	0.2132%	0.3724%	0.4367%	0.0967%
	Females	0.0672%	0.0307%	0.0541%	0.0388%	0.0234%	0.0349%	0.0634%	0.1213%	0.2021%	0.2678%	0.0631%
White ^c	Both Sexes	0.0915%	0.0362%	0.0694%	0.0476%	0.0305%	0.0456%	0.0890%	0.1831%	0.3133%	0.3652%	0.0872%
	Males	0.1060%	0.0372%	0.0763%	0.0509%	0.0351%	0.0535%	0.1099%	0.2345%	0.4170%	0.4812%	0.1068%
	Females	0.0770%	0.0351%	0.0622%	0.0441%	0.0256%	0.0373%	0.0683%	0.1352%	0.2271%	0.2959%	0.0703%
Black ^c	Both Sexes	0.0440%	0.0172%	0.0258%	0.0274%	0.0191%	0.0343%	0.0556%	0.1097%	0.1719%	0.2086%	0.0497%
	Males	0.0500%	0.0201%	0.0295%	0.0293%	0.0209%	0.0386%	0.0704%	0.1412%	0.2272%	0.2529%	0.0607%
	Females	0.0385%	0.0142%	0.0219%	0.0255%	0.0174%	0.0305%	0.0428%	0.0849%	0.1333%	0.1871%	0.0414%
Asian/ Pacific Islander ^c	Both Sexes	0.0414%	0.0266%	0.0492%	0.0294%	0.0210%	0.0331%	0.0435%	0.0604%	0.0991%	0.1168%	0.0421%
	Males	0.0489%	0.0301%	0.0538%	0.0339%	0.0237%	0.0427%	0.0504%	0.0751%	0.1321%	0.1701%	0.0513%
	Females	0.0344%	0.0230%	0.0443%	0.0250%	0.0185%	0.0245%	0.0375%	0.0482%	0.0737%	0.0842%	0.0346%
Hispanic ^d	Both Sexes	0.0525%	0.0357%	0.0692%	0.0460%	0.0279%	0.0395%	0.0560%	0.0957%	0.1584%	0.1802%	0.0602%
	Males	0.0587%	0.0389%	0.0745%	0.0518%	0.0354%	0.0459%	0.0623%	0.1160%	0.1834%	0.2547%	0.0705%
	Females	0.0462%	0.0324%	0.0636%	0.0396%	0.0199%	0.0328%	0.0500%	0.0783%	0.1398%	0.1356%	0.0512%

^a US 2012 cancer prevalence counts are based on 2012 cancer prevalence proportions from the SEER registries and 1/1/2012 US population estimates based on the average of 2011 and 2012 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 37 years for 1975-2012 SEER 9 data; 20 years for 1992-2012 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 37 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <37 and >=37. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 13.30
Acute Lymphocytic Leukemia

Estimated United States Cancer Prevalence Counts^a on January 1, 2012
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <20 ^e	0 to <37 ^e	>=37 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	17,081	13,446	11,446	9,730	8,142	6,255	55,252	71,898	3,278	75,176
	Males	9,480	7,618	6,175	5,431	4,443	3,384	30,632	39,375	1,680	41,055
	Females	7,601	5,828	5,271	4,299	3,699	2,871	24,620	32,523	1,598	34,121
White ^b	Both Sexes	14,348	11,402	10,036	8,376	7,130	5,665	47,334	62,330	+	+
	Males	7,881	6,453	5,384	4,726	3,945	3,080	26,198	34,133	+	+
	Females	6,467	4,949	4,652	3,650	3,185	2,585	21,136	28,197	+	+
Black ^b	Both Sexes	1,471	1,013	738	657	520	280	4,081	4,884	+	+
	Males	926	608	376	320	235	162	2,321	2,711	+	+
	Females	545	405	362	337	285	118	1,760	2,173	+	+
Asian/ Pacific Islander ^c	Both Sexes	770	712	462	+	+	+	2,422	+	+	+
	Males	432	396	282	+	+	+	1,375	+	+	+
	Females	338	316	180	+	+	+	1,047	+	+	+
Hispanic ^d	Both Sexes	5,171	3,540	2,644	+	+	+	13,595	+	+	+
	Males	2,976	2,054	1,392	+	+	+	7,775	+	+	+
	Females	2,195	1,486	1,252	+	+	+	5,820	+	+	+

Estimated prevalence percent^a on January 1, 2012, of the SEER^c population diagnosed in the previous 20 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted ^f	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0178%	0.0273%	0.0516%	0.0297%	0.0080%	0.0051%	0.0040%	0.0043%	0.0038%	0.0022%	0.0184%
	Males	0.0203%	0.0287%	0.0570%	0.0331%	0.0103%	0.0056%	0.0045%	0.0045%	0.0039%	0.0031%	0.0203%
	Females	0.0154%	0.0259%	0.0460%	0.0260%	0.0058%	0.0046%	0.0036%	0.0041%	0.0036%	0.0017%	0.0164%
White ^c	Both Sexes	0.0201%	0.0309%	0.0600%	0.0345%	0.0098%	0.0055%	0.0044%	0.0046%	0.0039%	0.0020%	0.0211%
	Males	0.0226%	0.0318%	0.0657%	0.0382%	0.0125%	0.0061%	0.0050%	0.0049%	0.0042%	0.0025%	0.0232%
	Females	0.0177%	0.0300%	0.0540%	0.0306%	0.0069%	0.0049%	0.0039%	0.0044%	0.0036%	0.0018%	0.0190%
Black ^c	Both Sexes	0.0090%	0.0124%	0.0204%	0.0158%	0.0032%	0.0028%	0.0021%	0.0024%	-	-	0.0083%
	Males	0.0107%	0.0156%	0.0237%	0.0179%	0.0035%	0.0028%	-	-	-	-	0.0094%
	Females	0.0075%	0.0090%	0.0170%	0.0138%	0.0030%	0.0027%	0.0031%	0.0024%	-	-	0.0072%
Asian/ Pacific Islander ^c	Both Sexes	0.0128%	0.0227%	0.0396%	0.0189%	0.0045%	0.0045%	0.0030%	0.0030%	0.0043%	0.0028%	0.0139%
	Males	0.0152%	0.0252%	0.0443%	0.0227%	0.0053%	0.0048%	0.0040%	0.0031%	-	0.0074%	0.0157%
	Females	0.0105%	0.0201%	0.0347%	0.0151%	0.0038%	0.0042%	0.0020%	0.0029%	0.0057%	-	0.0120%
Hispanic ^d	Both Sexes	0.0257%	0.0307%	0.0607%	0.0339%	0.0101%	0.0071%	0.0068%	0.0069%	0.0076%	-	0.0220%
	Males	0.0290%	0.0334%	0.0651%	0.0396%	0.0130%	0.0083%	0.0079%	0.0057%	-	-	0.0242%
	Females	0.0224%	0.0278%	0.0561%	0.0275%	0.0069%	0.0059%	0.0058%	0.0078%	0.0105%	-	0.0195%

^a US 2012 cancer prevalence counts are based on 2012 cancer prevalence proportions from the SEER registries and 1/1/2012 US population estimates based on the average of 2011 and 2012 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 37 years for 1975-2012 SEER 9 data; 20 years for 1992-2012 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 37 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <37 and >=37. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 14.21
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2012
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <20 ^e	0 to <37 ^e	>=37 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	36,620	8,385	2,786	1,115	697	475	49,235	50,271	463	50,734
	Males	27,039	5,883	1,792	662	311	219	35,560	35,990	73	36,063
	Females	9,581	2,502	994	453	386	256	13,675	14,281	390	14,671
White ^b	Both Sexes	26,566	6,036	1,977	814	547	390	35,626	36,468	356	36,824
	Males	19,706	4,407	1,224	473	230	162	25,937	26,268	227 ⁱ	26,495
	Females	6,860	1,629	753	341	317	228	9,689	10,200	129 ⁱ	10,329
Black ^b	Both Sexes	5,543	884	180	60	72	36	6,713	6,806	+	+
	Males	4,120	477	107	25	24	26	4,751	4,788	+	+
	Females	1,423	407	73	35	48	10	1,962	2,018	+	+
Asian/ Pacific Islander ^c	Both Sexes	3,815	1,293	505	+	+	+	5,755	+	+	+
	Males	2,716	901	360	+	+	+	4,081	+	+	+
	Females	1,099	392	145	+	+	+	1,674	+	+	+
Hispanic ^d	Both Sexes	5,720	1,147	362	+	+	+	7,459	+	+	+
	Males	4,141	824	195	+	+	+	5,320	+	+	+
	Females	1,579	323	167	+	+	+	2,139	+	+	+

Estimated prevalence percent^a on January 1, 2012, of the SEER^c population diagnosed in the previous 20 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted ^f	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0178%	0.0028%	0.0025%	0.0011%	0.0020%	0.0067%	0.0356%	0.0658%	0.0670%	0.0429%	0.0163%
	Males	0.0258%	0.0033%	0.0024%	0.0014%	0.0024%	0.0094%	0.0570%	0.1058%	0.0957%	0.0671%	0.0245%
	Females	0.0100%	0.0024%	0.0026%	0.0008%	0.0016%	0.0040%	0.0151%	0.0296%	0.0437%	0.0286%	0.0091%
White ^c	Both Sexes	0.0148%	0.0032%	0.0026%	0.0011%	0.0016%	0.0052%	0.0307%	0.0539%	0.0440%	0.0278%	0.0128%
	Males	0.0214%	0.0035%	0.0025%	0.0013%	0.0017%	0.0071%	0.0490%	0.0866%	0.0625%	0.0436%	0.0191%
	Females	0.0081%	0.0029%	0.0027%	0.0010%	0.0016%	0.0032%	0.0127%	0.0233%	0.0286%	0.0184%	0.0071%
Black ^c	Both Sexes	0.0147%	0.0009%	0.0008%	0.0013%	0.0021%	0.0046%	0.0334%	0.0868%	0.0547%	0.0294%	0.0155%
	Males	0.0215%	-	-	0.0019%	0.0023%	0.0061%	0.0521%	0.1482%	0.0812%	0.0431%	0.0243%
	Females	0.0084%	-	-	-	0.0019%	0.0033%	0.0171%	0.0386%	0.0362%	0.0228%	0.0086%
Asian/ Pacific Islander ^c	Both Sexes	0.0370%	0.0026%	0.0035%	0.0008%	0.0040%	0.0157%	0.0631%	0.1183%	0.2036%	0.1457%	0.0366%
	Males	0.0543%	0.0041%	0.0031%	-	0.0063%	0.0243%	0.1078%	0.1934%	0.3004%	0.2271%	0.0571%
	Females	0.0210%	-	0.0040%	-	0.0020%	0.0081%	0.0247%	0.0562%	0.1287%	0.0959%	0.0201%
Hispanic ^d	Both Sexes	0.0146%	0.0038%	0.0024%	0.0010%	0.0014%	0.0073%	0.0487%	0.0916%	0.0821%	0.0656%	0.0216%
	Males	0.0205%	0.0045%	0.0024%	0.0012%	0.0013%	0.0114%	0.0807%	0.1451%	0.1162%	0.0958%	0.0327%
	Females	0.0085%	0.0031%	0.0024%	0.0009%	0.0016%	0.0030%	0.0181%	0.0457%	0.0569%	0.0475%	0.0120%

^a US 2012 cancer prevalence counts are based on 2012 cancer prevalence proportions from the SEER registries and 1/1/2012 US population estimates based on the average of 2011 and 2012 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 37 years for 1975-2012 SEER 9 data; 20 years for 1992-2012 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 37 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <37 and >=37. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 15.27
Cancer of the Lung And Bronchus (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2012
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <20 ^e	0 to <37 ^e	>=37 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	233,827	80,267	42,112	23,166	12,241	6,603	385,066	402,326	6,482	408,808
	Males	107,351	34,954	17,903	10,199	5,280	2,905	172,829	180,530	3,700	184,230
	Females	126,476	45,313	24,209	12,967	6,961	3,698	212,237	221,796	2,782	224,578
White ^b	Both Sexes	197,170	69,783	36,974	20,131	10,983	5,734	329,144	344,368	2,551	346,919
	Males	90,128	29,757	15,541	8,709	4,643	2,518	146,275	152,955	1,180	154,135
	Females	107,042	40,026	21,433	11,422	6,340	3,216	182,869	191,413	1,371	192,784
Black ^b	Both Sexes	26,406	7,618	3,738	2,242	891	693	40,454	41,975	229	42,204
	Males	12,146	3,790	1,628	1,046	458	301	18,809	19,582	114	19,696
	Females	14,260	3,828	2,110	1,196	433	392	21,645	22,393	115	22,508
Asian/ Pacific Islander ^c	Both Sexes	7,914	2,119	1,072	+	+	+	11,640	+	+	+
	Males	3,940	1,016	542	+	+	+	5,806	+	+	+
	Females	3,974	1,103	530	+	+	+	5,834	+	+	+
Hispanic ^d	Both Sexes	9,212	2,741	1,427	+	+	+	14,092	+	+	+
	Males	4,247	1,256	609	+	+	+	6,468	+	+	+
	Females	4,965	1,485	818	+	+	+	7,624	+	+	+

Estimated prevalence percent^a on January 1, 2012, of the SEER^c population diagnosed in the previous 20 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted ^f	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.1031%	0.0002%	0.0004%	0.0018%	0.0058%	0.0243%	0.1087%	0.3206%	0.6462%	0.6341%	0.1004%
	Males	0.0947%	0.0002%	0.0003%	0.0019%	0.0053%	0.0217%	0.1009%	0.3252%	0.6558%	0.7040%	0.1022%
	Females	0.1114%	-	0.0004%	0.0018%	0.0063%	0.0268%	0.1161%	0.3165%	0.6385%	0.5927%	0.0997%
White ^c	Both Sexes	0.1113%	0.0002%	0.0004%	0.0021%	0.0062%	0.0241%	0.1066%	0.3252%	0.6728%	0.6532%	0.1028%
	Males	0.0993%	-	0.0003%	0.0020%	0.0052%	0.0217%	0.0975%	0.3195%	0.6584%	0.7030%	0.1014%
	Females	0.1232%	-	0.0005%	0.0021%	0.0072%	0.0266%	0.1156%	0.3305%	0.6848%	0.6234%	0.1046%
Black ^c	Both Sexes	0.0900%	-	-	0.0021%	0.0059%	0.0298%	0.1561%	0.4044%	0.6765%	0.5118%	0.1106%
	Males	0.0883%	-	-	0.0024%	0.0076%	0.0256%	0.1533%	0.4567%	0.7597%	0.6697%	0.1241%
	Females	0.0915%	-	-	0.0018%	0.0044%	0.0336%	0.1586%	0.3633%	0.6182%	0.4352%	0.1020%
Asian/ Pacific Islander ^c	Both Sexes	0.0781%	-	-	0.0043%	0.0216%	0.0851%	0.2389%	0.4937%	0.5748%	0.0799%	
	Males	0.0812%	-	-	0.0039%	0.0203%	0.0836%	0.2730%	0.5930%	0.7189%	0.0928%	
	Females	0.0754%	-	-	0.0046%	0.0228%	0.0864%	0.2108%	0.4169%	0.4868%	0.0707%	
Hispanic ^d	Both Sexes	0.0276%	-	-	0.0016%	0.0037%	0.0137%	0.0471%	0.1476%	0.3289%	0.3576%	0.0508%
	Males	0.0249%	-	-	0.0014%	0.0033%	0.0129%	0.0463%	0.1509%	0.3366%	0.4114%	0.0533%
	Females	0.0305%	-	-	0.0019%	0.0041%	0.0145%	0.0478%	0.1448%	0.3232%	0.3254%	0.0492%

^a US 2012 cancer prevalence counts are based on 2012 cancer prevalence proportions from the SEER registries and 1/1/2012 US population estimates based on the average of 2011 and 2012 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 37 years for 1975-2012 SEER 9 data; 20 years for 1992-2012 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 37 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <37 and >=37. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 16.21
Melanoma of the Skin (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2012
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <20 ^e	0 to <37 ^e	>=37 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	285,569	216,879	161,838	112,858	82,325	56,741	813,687	962,438	34,149	996,587
	Males	153,364	110,812	80,586	55,278	36,818	23,816	417,183	478,737	10,877	489,614
	Females	132,205	106,067	81,252	57,580	45,507	32,925	396,504	483,701	23,272	506,973
White ^b	Both Sexes	273,641	210,027	156,380	108,243	79,323	54,316	783,649	926,489	32,186	958,675
	Males	147,752	107,549	77,889	53,080	35,829	22,752	402,978	462,226	10,357	472,583
	Females	125,889	102,478	78,491	55,163	43,494	31,564	380,671	464,263	21,829	486,092
Black ^b	Both Sexes	1,174	655	609	390	324	146	2,978	3,420	99	3,519
	Males	489	228	276	242	106	37	1,307	1,417	40	1,457
	Females	685	427	333	148	218	109	1,671	2,003	59	2,062
Asian/ Pacific Islander ^c	Both Sexes	744	543	399	+	+	+	1,902	+	+	+
	Males	363	244	149	+	+	+	853	+	+	+
	Females	381	299	250	+	+	+	1,049	+	+	+
Hispanic ^d	Both Sexes	5,092	3,745	2,916	+	+	+	13,681	+	+	+
	Males	2,032	1,303	1,049	+	+	+	5,073	+	+	+
	Females	3,060	2,442	1,867	+	+	+	8,608	+	+	+

Estimated prevalence percent^a on January 1, 2012, of the SEER^c population diagnosed in the previous 20 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted ^f	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.2036%	0.0003%	0.0031%	0.0279%	0.0980%	0.2044%	0.3527%	0.5300%	0.6791%	0.7522%	0.1937%
	Males	0.2142%	0.0002%	0.0029%	0.0184%	0.0689%	0.1635%	0.3528%	0.6487%	0.9461%	1.1560%	0.2199%
	Females	0.1933%	0.0005%	0.0032%	0.0378%	0.1273%	0.2449%	0.3526%	0.4227%	0.4627%	0.5134%	0.1770%
White ^c	Both Sexes	0.2650%	0.0003%	0.0038%	0.0367%	0.1296%	0.2642%	0.4469%	0.6594%	0.8445%	0.9001%	0.2436%
	Males	0.2766%	-	0.0037%	0.0241%	0.0882%	0.2071%	0.4404%	0.7946%	1.1641%	1.3783%	0.2710%
	Females	0.2534%	0.0003%	0.0039%	0.0503%	0.1731%	0.3231%	0.4533%	0.5331%	0.5790%	0.6142%	0.2273%
Black ^c	Both Sexes	0.0065%	-	-	-	0.0044%	0.0063%	0.0096%	0.0186%	0.0324%	0.0569%	0.0081%
	Males	0.0062%	-	-	-	0.0027%	0.0043%	0.0105%	0.0209%	0.0433%	0.0834%	0.0093%
	Females	0.0068%	-	-	-	0.0058%	0.0081%	0.0089%	0.0169%	0.0247%	0.0441%	0.0074%
Asian/ Pacific Islander ^c	Both Sexes	0.0118%	0.0008%	0.0010%	0.0028%	0.0065%	0.0137%	0.0181%	0.0288%	0.0356%	0.0461%	0.0115%
	Males	0.0111%	-	-	0.0028%	0.0064%	0.0091%	0.0162%	0.0337%	0.0459%	0.0519%	0.0116%
	Females	0.0124%	-	-	0.0028%	0.0067%	0.0177%	0.0197%	0.0248%	0.0276%	0.0425%	0.0115%
Hispanic ^d	Both Sexes	0.0269%	0.0003%	0.0013%	0.0043%	0.0175%	0.0415%	0.0710%	0.1068%	0.1332%	0.1737%	0.0392%
	Males	0.0196%	-	0.0011%	0.0021%	0.0107%	0.0236%	0.0499%	0.0932%	0.1417%	0.2337%	0.0344%
	Females	0.0344%	-	0.0014%	0.0068%	0.0247%	0.0601%	0.0912%	0.1184%	0.1269%	0.1378%	0.0451%

^a US 2012 cancer prevalence counts are based on 2012 cancer prevalence proportions from the SEER registries and 1/1/2012 US population estimates based on the average of 2011 and 2012 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 37 years for 1975-2012 SEER 9 data; 20 years for 1992-2012 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 37 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <37 and >=37. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 17.13
MesotheliomaEstimated United States Cancer Prevalence Counts^a on January 1, 2012
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <20 ^e	0 to <37 ^e	>=37 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	3,247	644	404	232	185	128	4,600	4,957	+	+
	Males	2,341	338	170	47	39	71	2,920	3,042	+	+
	Females	906	306	234	185	146	57	1,680	1,915	+	+
White ^b	Both Sexes	3,056	585	310	208	182	116	4,233	4,549	+	+
	Males	2,231	302	135	47	36	59	2,739	2,832	+	+
	Females	825	283	175	161	146	57	1,494	1,717	+	+
Black ^b	Both Sexes	109	27	71	12	0	12	219	254	+	+
	Males	48	15	35	0	0	12	98	122	+	+
	Females	61	12	36	12	0	0	121	132	+	+
Asian/ Pacific Islander ^c	Both Sexes	56	18	7	+	+	+	86	+	+	+
	Males	36	8	4	+	+	+	53	+	+	+
	Females	20	10	3	+	+	+	33	+	+	+
Hispanic ^d	Both Sexes	277	41	61	+	+	+	379	+	+	+
	Males	193	20	49	+	+	+	262	+	+	+
	Females	84	21	12	+	+	+	117	+	+	+

Estimated prevalence percent^a on January 1, 2012, of the SEER^c population diagnosed in the previous 20 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted ^f	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0013%	-	-	-	0.0004%	0.0004%	0.0014%	0.0042%	0.0058%	0.0074%	0.0012%
	Males	0.0017%	-	-	-	0.0003%	0.0003%	0.0016%	0.0064%	0.0087%	0.0139%	0.0017%
	Females	0.0009%	-	-	-	0.0005%	0.0005%	0.0012%	0.0022%	0.0035%	0.0036%	0.0008%
White ^c	Both Sexes	0.0015%	-	-	-	0.0004%	0.0005%	0.0016%	0.0048%	0.0068%	0.0077%	0.0014%
	Males	0.0020%	-	-	-	0.0004%	0.0005%	0.0019%	0.0071%	0.0105%	0.0138%	0.0020%
	Females	0.0010%	-	-	-	0.0004%	0.0005%	0.0014%	0.0026%	0.0038%	0.0041%	0.0009%
Black ^c	Both Sexes	0.0006%	-	-	-	-	-	0.0010%	0.0023%	0.0045%	0.0050%	0.0008%
	Males	0.0007%	-	-	-	-	-	-	0.0047%	-	-	0.0009%
	Females	0.0006%	-	-	-	-	-	-	-	0.0048%	-	0.0007%
Asian/ Pacific Islander ^c	Both Sexes	0.0006%	-	-	-	0.0006%	-	-	0.0020%	-	0.0045%	0.0005%
	Males	0.0008%	-	-	-	-	-	-	0.0035%	-	0.0118%	0.0009%
	Females	0.0004%	-	-	-	0.0010%	-	-	-	-	-	0.0004%
Hispanic ^d	Both Sexes	0.0007%	-	-	-	0.0004%	-	0.0020%	0.0053%	0.0030%	0.0074%	0.0012%
	Males	0.0010%	-	-	-	-	-	0.0027%	0.0092%	0.0050%	0.0137%	0.0019%
	Females	0.0005%	-	-	-	-	-	0.0014%	0.0019%	-	-	0.0006%

^a US 2012 cancer prevalence counts are based on 2012 cancer prevalence proportions from the SEER registries and 1/1/2012 US population estimates based on the average of 2011 and 2012 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 37 years for 1975-2012 SEER 9 data; 20 years for 1992-2012 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 37 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <37 and >=37. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 18.21
Myeloma (Invasive)Estimated United States Cancer Prevalence Counts^a on January 1, 2012
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <20 ^e	0 to <37 ^e	>=37 ^g	Complete ^h	
Race	Sex											
All Races ^b	Both Sexes	56,508	20,540	7,352	3,082	1,394	365	88,267	89,416	242	89,658	
	Males	30,245	11,069	4,053	1,603	754	192	47,394	47,990	84	48,074	
	Females	26,263	9,471	3,299	1,479	640	173	40,873	41,426	158	41,584	
White ^b	Both Sexes	41,956	15,946	5,693	2,401	1,044	304	66,608	67,438	0	67,438	
	Males	23,218	8,979	3,361	1,246	569	174	37,140	37,583	0 ⁱ	37,583	
	Females	18,738	6,967	2,332	1,155	475	130	29,468	29,855	0 ⁱ	29,855	
Black ^b	Both Sexes	11,675	3,969	1,304	564	294	41	17,660	17,922	+	+	
	Males	5,512	1,791	508	302	173	13	8,189	8,338	+	+	
	Females	6,163	2,178	796	262	121	28	9,471	9,584	+	+	
Asian/ Pacific Islander ^c	Both Sexes	1,548	395	208	+	+	+	2,231	+	+	+	
	Males	824	196	101	+	+	+	1,158	+	+	+	
	Females	724	199	107	+	+	+	1,073	+	+	+	
Hispanic ^d	Both Sexes	4,223	1,366	618	+	+	+	6,386	+	+	+	
	Males	2,296	749	340	+	+	+	3,488	+	+	+	
	Females	1,927	617	278	+	+	+	2,898	+	+	+	

Estimated prevalence percent^a on January 1, 2012, of the SEER^c population diagnosed in the previous 20 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)								Age-Adjusted ^f		
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
Race	Sex											
All Races ^c	Both Sexes	0.0250%	-	-	0.0004%	0.0022%	0.0117%	0.0358%	0.0839%	0.1306%	0.1122%	0.0238%
	Males	0.0274%	-	-	0.0006%	0.0027%	0.0138%	0.0401%	0.0999%	0.1538%	0.1425%	0.0283%
	Females	0.0227%	-	-	-	0.0017%	0.0096%	0.0317%	0.0695%	0.1119%	0.0942%	0.0202%
White ^c	Both Sexes	0.0245%	-	-	0.0005%	0.0022%	0.0104%	0.0326%	0.0782%	0.1232%	0.1054%	0.0222%
	Males	0.0276%	-	-	0.0007%	0.0027%	0.0129%	0.0370%	0.0955%	0.1487%	0.1408%	0.0271%
	Females	0.0214%	-	-	-	0.0018%	0.0078%	0.0282%	0.0619%	0.1021%	0.0843%	0.0181%
Black ^c	Both Sexes	0.0401%	-	-	-	0.0034%	0.0277%	0.0729%	0.1754%	0.2486%	0.2201%	0.0481%
	Males	0.0394%	-	-	-	0.0041%	0.0277%	0.0756%	0.2014%	0.2796%	0.2221%	0.0524%
	Females	0.0407%	-	-	-	0.0028%	0.0277%	0.0707%	0.1549%	0.2269%	0.2191%	0.0449%
Asian/ Pacific Islander ^c	Both Sexes	0.0147%	-	-	-	0.0009%	0.0054%	0.0207%	0.0471%	0.0877%	0.0815%	0.0147%
	Males	0.0157%	-	-	-	0.0014%	0.0072%	0.0246%	0.0521%	0.0996%	0.0988%	0.0172%
	Females	0.0137%	-	-	-	-	0.0038%	0.0174%	0.0429%	0.0786%	0.0710%	0.0127%
Hispanic ^d	Both Sexes	0.0125%	-	-	0.0006%	0.0023%	0.0095%	0.0327%	0.0663%	0.1233%	0.0977%	0.0211%
	Males	0.0134%	-	-	0.0010%	0.0029%	0.0121%	0.0385%	0.0806%	0.1312%	0.1226%	0.0247%
	Females	0.0116%	-	-	-	0.0016%	0.0069%	0.0272%	0.0540%	0.1175%	0.0828%	0.0181%

^a US 2012 cancer prevalence counts are based on 2012 cancer prevalence proportions from the SEER registries and 1/1/2012 US population estimates based on the average of 2011 and 2012 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 37 years for 1975-2012 SEER 9 data; 20 years for 1992-2012 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 37 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <37 and >=37. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 19.25
Non-Hodgkin LymphomaEstimated United States Cancer Prevalence Counts^a on January 1, 2012
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <20 ^e	0 to <37 ^e	>=37 ^g	Complete ^h	
Race	Sex											
All Races ^b	Both Sexes	205,597	143,043	87,628	50,507	28,179	15,884	499,961	541,393	8,232	549,625	
	Males	110,512	73,527	45,359	26,351	14,922	8,603	262,705	284,672	4,332	289,004	
	Females	95,085	69,516	42,269	24,156	13,257	7,281	237,256	256,721	3,900	260,621	
White ^b	Both Sexes	177,024	125,584	77,296	44,460	25,369	14,430	436,199	473,815	8,597	482,412	
	Males	95,569	64,896	40,143	23,215	13,477	7,779	230,047	250,031	4,868	254,899	
	Females	81,455	60,688	37,153	21,245	11,892	6,651	206,152	223,784	3,729	227,513	
Black ^b	Both Sexes	16,520	11,569	6,486	3,710	1,796	916	39,149	41,616	683	42,299	
	Males	8,420	5,675	3,240	1,899	923	534	19,661	20,969	289	21,258	
	Females	8,100	5,894	3,246	1,811	873	382	19,488	20,647	394	21,041	
Asian/ Pacific Islander ^c	Both Sexes	6,342	3,800	2,382	+	+	+	13,826	+	+	+	
	Males	3,369	1,919	1,274	+	+	+	7,211	+	+	+	
	Females	2,973	1,881	1,108	+	+	+	6,615	+	+	+	
Hispanic ^d	Both Sexes	16,134	9,712	5,785	+	+	+	34,651	+	+	+	
	Males	8,441	4,947	3,073	+	+	+	18,053	+	+	+	
	Females	7,693	4,765	2,712	+	+	+	16,598	+	+	+	

Estimated prevalence percent^a on January 1, 2012, of the SEER^c population diagnosed in the previous 20 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted ^f	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
Race	Sex											
All Races ^c	Both Sexes	0.1402%	0.0029%	0.0120%	0.0252%	0.0461%	0.0991%	0.1958%	0.3710%	0.5961%	0.6826%	0.1347%
	Males	0.1498%	0.0037%	0.0159%	0.0306%	0.0540%	0.1166%	0.2266%	0.4203%	0.6685%	0.7803%	0.1546%
	Females	0.1309%	0.0021%	0.0078%	0.0196%	0.0382%	0.0818%	0.1664%	0.3265%	0.5375%	0.6248%	0.1173%
White ^c	Both Sexes	0.1571%	0.0030%	0.0125%	0.0264%	0.0479%	0.1057%	0.2098%	0.3998%	0.6519%	0.7289%	0.1446%
	Males	0.1671%	0.0039%	0.0170%	0.0322%	0.0560%	0.1248%	0.2425%	0.4471%	0.7281%	0.8267%	0.1653%
	Females	0.1472%	0.0021%	0.0077%	0.0202%	0.0393%	0.0860%	0.1774%	0.3557%	0.5885%	0.6705%	0.1260%
Black ^c	Both Sexes	0.0878%	0.0026%	0.0110%	0.0244%	0.0509%	0.0921%	0.1623%	0.2844%	0.3672%	0.3532%	0.0996%
	Males	0.0934%	0.0027%	0.0130%	0.0312%	0.0621%	0.1046%	0.1884%	0.3394%	0.3521%	0.4004%	0.1117%
	Females	0.0828%	0.0025%	0.0090%	0.0178%	0.0411%	0.0811%	0.1395%	0.2413%	0.3778%	0.3302%	0.0896%
Asian/ Pacific Islander ^c	Both Sexes	0.0883%	0.0029%	0.0102%	0.0189%	0.0305%	0.0618%	0.1243%	0.2260%	0.3764%	0.4951%	0.0876%
	Males	0.0955%	0.0035%	0.0134%	0.0209%	0.0343%	0.0693%	0.1429%	0.2605%	0.4486%	0.5988%	0.1027%
	Females	0.0818%	0.0023%	0.0067%	0.0169%	0.0270%	0.0552%	0.1084%	0.1974%	0.3206%	0.4317%	0.0758%
Hispanic ^d	Both Sexes	0.0679%	0.0031%	0.0094%	0.0196%	0.0324%	0.0771%	0.1502%	0.2847%	0.4602%	0.5289%	0.1042%
	Males	0.0695%	0.0036%	0.0126%	0.0231%	0.0384%	0.0875%	0.1692%	0.2984%	0.4720%	0.5701%	0.1130%
	Females	0.0663%	0.0026%	0.0061%	0.0157%	0.0260%	0.0662%	0.1319%	0.2730%	0.4515%	0.5043%	0.0962%

^a US 2012 cancer prevalence counts are based on 2012 cancer prevalence proportions from the SEER registries and 1/1/2012 US population estimates based on the average of 2011 and 2012 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 37 years for 1975-2012 SEER 9 data; 20 years for 1992-2012 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 37 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <37 and >=37. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 20.27
Cancer of the Oral Cavity and Pharynx (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2012
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <20 ^e	0 to <37 ^e	>=37 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	108,762	66,555	42,301	26,989	17,258	11,366	252,895	282,380	8,728	291,108
	Males	76,856	45,595	27,558	16,900	10,165	6,484	171,885	188,848	4,931	193,779
	Females	31,906	20,960	14,743	10,089	7,093	4,882	81,010	93,532	3,797	97,329
White ^b	Both Sexes	93,692	57,853	36,599	23,463	15,216	9,917	218,927	244,661	7,471	252,132
	Males	66,971	40,346	24,141	14,938	9,167	5,807	150,908	166,123	4,232	170,355
	Females	26,721	17,507	12,458	8,525	6,049	4,110	68,019	78,538	3,239	81,777
Black ^b	Both Sexes	8,294	4,536	3,183	1,860	1,080	891	18,408	20,530	422	20,952
	Males	5,530	2,719	1,881	962	478	380	11,314	12,172	167	12,339
	Females	2,764	1,817	1,302	898	602	511	7,094	8,358	255	8,613
Asian/ Pacific Islander ^c	Both Sexes	4,029	2,673	1,722	+	+	+	9,517	+	+	+
	Males	2,620	1,631	1,010	+	+	+	5,892	+	+	+
	Females	1,409	1,042	712	+	+	+	3,625	+	+	+
Hispanic ^d	Both Sexes	6,104	3,112	2,047	+	+	+	12,560	+	+	+
	Males	3,997	1,887	1,250	+	+	+	7,795	+	+	+
	Females	2,107	1,225	797	+	+	+	4,765	+	+	+

Estimated prevalence percent^a on January 1, 2012, of the SEER^c population diagnosed in the previous 20 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)									Age-Adjusted ^f	
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0708%	0.0005%	0.0024%	0.0062%	0.0163%	0.0474%	0.1310%	0.2300%	0.2663%	0.2386%	0.0658%
	Males	0.0970%	0.0005%	0.0020%	0.0060%	0.0157%	0.0596%	0.1946%	0.3547%	0.4014%	0.3507%	0.0953%
	Females	0.0453%	0.0004%	0.0029%	0.0065%	0.0169%	0.0353%	0.0700%	0.1173%	0.1568%	0.1723%	0.0406%
White ^c	Both Sexes	0.0772%	0.0005%	0.0024%	0.0062%	0.0152%	0.0461%	0.1358%	0.2467%	0.2863%	0.2489%	0.0687%
	Males	0.1067%	0.0004%	0.0019%	0.0056%	0.0143%	0.0595%	0.2043%	0.3803%	0.4296%	0.3658%	0.1000%
	Females	0.0478%	0.0005%	0.0030%	0.0068%	0.0161%	0.0323%	0.0679%	0.1220%	0.1673%	0.1791%	0.0410%
Black ^c	Both Sexes	0.0426%	-	0.0023%	0.0063%	0.0137%	0.0354%	0.1040%	0.1634%	0.1812%	0.1263%	0.0472%
	Males	0.0557%	-	0.0017%	0.0081%	0.0123%	0.0376%	0.1473%	0.2578%	0.2760%	0.2030%	0.0672%
	Females	0.0306%	-	0.0029%	0.0046%	0.0149%	0.0335%	0.0662%	0.0894%	0.1149%	0.0891%	0.0321%
Asian/ Pacific Islander ^c	Both Sexes	0.0594%	-	0.0024%	0.0063%	0.0232%	0.0588%	0.1173%	0.1643%	0.1967%	0.2040%	0.0570%
	Males	0.0767%	-	0.0030%	0.0061%	0.0259%	0.0725%	0.1653%	0.2398%	0.2906%	0.2810%	0.0784%
	Females	0.0435%	-	0.0017%	0.0066%	0.0208%	0.0467%	0.0760%	0.1018%	0.1241%	0.1570%	0.0398%
Hispanic ^d	Both Sexes	0.0246%	0.0004%	0.0015%	0.0046%	0.0098%	0.0240%	0.0640%	0.1261%	0.1635%	0.1739%	0.0380%
	Males	0.0300%	-	0.0011%	0.0044%	0.0094%	0.0281%	0.0891%	0.1783%	0.2480%	0.2288%	0.0520%
	Females	0.0190%	-	0.0019%	0.0048%	0.0101%	0.0198%	0.0399%	0.0813%	0.1008%	0.1410%	0.0266%

^a US 2012 cancer prevalence counts are based on 2012 cancer prevalence proportions from the SEER registries and 1/1/2012 US population estimates based on the average of 2011 and 2012 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 37 years for 1975-2012 SEER 9 data; 20 years for 1992-2012 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 37 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <37 and >=37. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 21.15
Cancer of the Ovary (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2012
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <20 ^e	0 to <37 ^e	>=37 ^g	Complete ^h
<u>Race</u>										
All Races ^b	59,878	31,489	24,695	19,686	16,447	13,256	142,838	177,706	14,740	192,446
White ^b	51,577	27,236	21,555	17,247	14,932	11,885	124,164	155,277	12,490	167,767
Black ^b	4,582	2,223	1,612	1,413	938	836	10,113	12,583	1,505	14,088
Asian/Pacific Islander ^c	2,711	1,646	1,110	+	+	+	6,210	+	+	+
Hispanic ^d	5,785	2,815	1,865	+	+	+	11,901	+	+	+

Estimated prevalence percent^a on January 1, 2012, of the SEER^c population diagnosed in the previous 20 years
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f All Ages	
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+		
<u>Race</u>												
All Races ^c	0.0819%	0.0004%	0.0063%	0.0183%	0.0331%	0.0705%	0.1510%	0.2258%	0.2461%	0.1971%	0.0734%	
White ^c	0.0904%	0.0004%	0.0064%	0.0194%	0.0344%	0.0720%	0.1597%	0.2451%	0.2708%	0.2115%	0.0783%	
Black ^c	0.0447%	-	0.0040%	0.0130%	0.0256%	0.0514%	0.0891%	0.1242%	0.1525%	0.1413%	0.0468%	
Asian/Pacific Islander ^c	0.0726%	-	0.0081%	0.0181%	0.0333%	0.0774%	0.1531%	0.1916%	0.1698%	0.1324%	0.0658%	
Hispanic ^d	0.0475%	-	0.0085%	0.0217%	0.0336%	0.0554%	0.1181%	0.1827%	0.1944%	0.1747%	0.0615%	

- ^a Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.
^a US 2012 cancer prevalence counts are based on 2012 cancer prevalence proportions from the SEER registries and 1/1/2012 US population estimates based on the average of 2011 and 2012 population estimates from the US Bureau of the Census.
^{b c d} Prevalence was calculated using the First Malignant Primary Only for a person. Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.
^e Maximum limited-duration prevalence: 37 years for 1975-2012 SEER 9 data; 20 years for 1992-2012 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).
^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.
^{g h i} (g) Cases diagnosed more than 37 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <37 and >=37. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.
 - Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.
 + Not available.

Table 22.21
Cancer of the Pancreas (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2012
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <20 ^e	0 to <37 ^e	>=37 ^g	Complete ^h	
Race	Sex											
All Races ^b	Both Sexes	34,042	5,941	2,492	1,380	789	430	44,240	45,432	270	45,702	
	Males	17,628	2,644	1,241	479	395	107	22,190	22,666	117	22,783	
	Females	16,414	3,297	1,251	901	394	323	22,050	22,766	153	22,919	
White ^b	Both Sexes	27,956	5,052	1,946	1,019	597	332	36,320	37,195	220	37,415	
	Males	14,674	2,268	1,083	345	336	74	18,561	18,920	104	19,024	
	Females	13,282	2,784	863	674	261	258	17,759	18,275	116	18,391	
Black ^b	Both Sexes	4,327	581	345	199	153	60	5,476	5,722	127	5,849	
	Males	2,084	261	93	42	52	26	2,481	2,581	53	2,634	
	Females	2,243	320	252	157	101	34	2,995	3,141	74	3,215	
Asian/ Pacific Islander ^c	Both Sexes	1,335	239	121	+	+	+	1,755	+	+	+	
	Males	614	109	41	+	+	+	789	+	+	+	
	Females	721	130	80	+	+	+	966	+	+	+	
Hispanic ^d	Both Sexes	2,386	473	203	+	+	+	3,216	+	+	+	
	Males	1,247	155	83	+	+	+	1,518	+	+	+	
	Females	1,139	318	120	+	+	+	1,698	+	+	+	

Estimated prevalence percent^a on January 1, 2012, of the SEER^c population diagnosed in the previous 20 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)									Age-Adjusted ^f	
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
Race	Sex											
All Races ^c	Both Sexes	0.0129%	-	0.0001%	0.0008%	0.0020%	0.0070%	0.0187%	0.0404%	0.0624%	0.0617%	0.0123%
	Males	0.0130%	-	-	0.0005%	0.0016%	0.0073%	0.0209%	0.0449%	0.0690%	0.0661%	0.0133%
	Females	0.0129%	-	-	0.0011%	0.0025%	0.0068%	0.0167%	0.0363%	0.0570%	0.0591%	0.0115%
White ^c	Both Sexes	0.0134%	-	-	0.0007%	0.0018%	0.0070%	0.0180%	0.0407%	0.0623%	0.0601%	0.0122%
	Males	0.0137%	-	-	0.0005%	0.0015%	0.0073%	0.0204%	0.0457%	0.0702%	0.0658%	0.0134%
	Females	0.0130%	-	-	0.0009%	0.0022%	0.0066%	0.0156%	0.0361%	0.0557%	0.0567%	0.0111%
Black ^c	Both Sexes	0.0124%	-	-	0.0010%	0.0030%	0.0089%	0.0261%	0.0466%	0.0661%	0.0699%	0.0146%
	Males	0.0114%	-	-	-	0.0019%	0.0092%	0.0282%	0.0514%	0.0642%	0.0554%	0.0143%
	Females	0.0133%	-	-	0.0013%	0.0040%	0.0087%	0.0243%	0.0428%	0.0675%	0.0770%	0.0147%
Asian/ Pacific Islander ^c	Both Sexes	0.0115%	-	-	0.0009%	0.0020%	0.0061%	0.0169%	0.0331%	0.0595%	0.0659%	0.0114%
	Males	0.0107%	-	-	-	0.0019%	0.0054%	0.0173%	0.0345%	0.0607%	0.0726%	0.0117%
	Females	0.0121%	-	-	0.0017%	0.0021%	0.0067%	0.0166%	0.0320%	0.0586%	0.0618%	0.0113%
Hispanic ^d	Both Sexes	0.0063%	-	-	0.0008%	0.0013%	0.0056%	0.0143%	0.0356%	0.0451%	0.0675%	0.0104%
	Males	0.0058%	-	-	0.0007%	0.0007%	0.0059%	0.0150%	0.0400%	0.0525%	0.0493%	0.0105%
	Females	0.0068%	-	-	0.0010%	0.0020%	0.0053%	0.0137%	0.0318%	0.0396%	0.0784%	0.0101%

^a US 2012 cancer prevalence counts are based on 2012 cancer prevalence proportions from the SEER registries and 1/1/2012 US population estimates based on the average of 2011 and 2012 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 37 years for 1975-2012 SEER 9 data; 20 years for 1992-2012 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 37 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <37 and >=37. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 23.15
Cancer of the Prostate (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2012
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <20 ^e	0 to <37 ^e	>=37 ^g	Complete ^h
<u>Race</u>										
All Races ^b	1,044,262	809,960	554,045	290,151	79,382	12,743	2,749,706	2,794,570	1,022	2,795,592
White ^b	836,144	669,947	468,474	250,656	72,322	11,765	2,272,023	2,312,969	1,192	2,314,161
Black ^b	151,342	107,605	68,225	32,343	5,501	857	363,061	366,166	102	366,268
Asian/Pacific Islander ^c	21,838	15,684	9,110	+	+	+	50,334	+	+	+
Hispanic ^d	60,598	44,337	26,562	+	+	+	143,146	+	+	+

Estimated prevalence percent^a on January 1, 2012, of the SEER^c population diagnosed in the previous 20 years
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f All Ages
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	
<u>Race</u>											
All Races ^c	1.5229%	-	-	0.0003%	0.0011%	0.0846%	1.0365%	5.0125%	11.9057%	14.8461%	1.7054%
White ^c	1.6077%	-	-	0.0003%	0.0009%	0.0717%	0.9901%	4.9572%	11.9866%	14.6943%	1.6925%
Black ^c	1.7122%	-	-	-	0.0032%	0.2305%	2.0081%	8.5683%	17.8128%	20.6319%	2.6571%
Asian/Pacific Islander ^c	0.7345%	-	-	-	-	0.0227%	0.3551%	2.0303%	6.0832%	9.9317%	0.8893%
Hispanic ^d	0.5504%	-	-	-	-	0.0410%	0.6150%	3.4399%	9.5140%	12.8163%	1.3359%

^a US 2012 cancer prevalence counts are based on 2012 cancer prevalence proportions from the SEER registries and 1/1/2012 US population estimates based on the average of 2011 and 2012 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 37 years for 1975-2012 SEER 9 data; 20 years for 1992-2012 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 37 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <37 and >=37. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 24.21
Cancer of the Stomach (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2012
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <20 ^e	0 to <37 ^e	>=37 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	37,369	16,901	9,494	5,313	3,425	2,037	70,550	75,640	1,189	76,829
	Males	22,014	9,683	5,468	2,948	1,539	1,053	40,773	43,300	675	43,975
	Females	15,355	7,218	4,026	2,365	1,886	984	29,777	32,340	514	32,854
White ^b	Both Sexes	27,715	12,472	6,831	3,917	2,451	1,490	51,925	55,769	947	56,716
	Males	17,131	7,459	4,006	2,091	1,057	801	31,097	33,063	524	33,587
	Females	10,584	5,013	2,825	1,826	1,394	689	20,828	22,706	423	23,129
Black ^b	Both Sexes	5,729	2,640	1,562	701	494	327	10,870	11,530	164	11,694
	Males	2,682	1,243	804	443	265	158	5,316	5,635	91	5,726
	Females	3,047	1,397	758	258	229	169	5,554	5,895	73	5,968
Asian/ Pacific Islander ^c	Both Sexes	3,360	1,783	1,029	+	+	+	6,816	+	+	+
	Males	1,884	984	581	+	+	+	3,825	+	+	+
	Females	1,476	799	448	+	+	+	2,991	+	+	+
Hispanic ^d	Both Sexes	5,659	2,385	1,221	+	+	+	9,978	+	+	+
	Males	3,168	1,294	658	+	+	+	5,404	+	+	+
	Females	2,491	1,091	563	+	+	+	4,574	+	+	+

Estimated prevalence percent^a on January 1, 2012, of the SEER^c population diagnosed in the previous 20 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	Age-Adjusted ^f All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0251%	-	0.0001%	0.0007%	0.0035%	0.0125%	0.0296%	0.0671%	0.1265%	0.1773%	0.0242%
	Males	0.0291%	-	0.0002%	0.0005%	0.0033%	0.0130%	0.0359%	0.0891%	0.1711%	0.2358%	0.0312%
	Females	0.0212%	-	-	0.0009%	0.0037%	0.0121%	0.0236%	0.0473%	0.0903%	0.1428%	0.0187%
White ^c	Both Sexes	0.0213%	-	0.0001%	0.0008%	0.0032%	0.0111%	0.0246%	0.0565%	0.1013%	0.1303%	0.0195%
	Males	0.0254%	-	-	0.0006%	0.0032%	0.0117%	0.0304%	0.0775%	0.1398%	0.1740%	0.0256%
	Females	0.0173%	-	-	0.0009%	0.0033%	0.0104%	0.0189%	0.0370%	0.0693%	0.1041%	0.0146%
Black ^c	Both Sexes	0.0239%	-	-	-	0.0034%	0.0154%	0.0437%	0.0828%	0.1527%	0.1990%	0.0295%
	Males	0.0244%	-	-	-	0.0028%	0.0151%	0.0485%	0.0999%	0.1962%	0.2113%	0.0341%
	Females	0.0234%	-	-	-	0.0039%	0.0156%	0.0396%	0.0694%	0.1222%	0.1931%	0.0262%
Asian/ Pacific Islander ^c	Both Sexes	0.0466%	-	-	0.0007%	0.0047%	0.0179%	0.0458%	0.1152%	0.2504%	0.4726%	0.0478%
	Males	0.0544%	-	-	-	0.0042%	0.0183%	0.0584%	0.1505%	0.3362%	0.6467%	0.0629%
	Females	0.0396%	-	-	-	0.0052%	0.0174%	0.0351%	0.0859%	0.1840%	0.3662%	0.0369%
Hispanic ^d	Both Sexes	0.0196%	-	-	0.0009%	0.0055%	0.0164%	0.0394%	0.0923%	0.1857%	0.2508%	0.0342%
	Males	0.0208%	-	-	0.0009%	0.0055%	0.0166%	0.0441%	0.1168%	0.2478%	0.2943%	0.0419%
	Females	0.0184%	-	-	0.0008%	0.0054%	0.0162%	0.0350%	0.0713%	0.1396%	0.2248%	0.0284%

^a US 2012 cancer prevalence counts are based on 2012 cancer prevalence proportions from the SEER registries and 1/1/2012 US population estimates based on the average of 2011 and 2012 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 37 years for 1975-2012 SEER 9 data; 20 years for 1992-2012 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 37 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <37 and >=37. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 25.15
Cancer of the Testis (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2012
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <20 ^e	0 to <37 ^e	>=37 ^g	Complete ^h
<u>Race</u>										
All Races ^b	42,435	39,377	36,757	31,998	28,620	21,406	162,426	218,903	14,699	233,602
White ^b	38,830	37,097	34,463	30,469	27,485	20,712	152,188	206,644	14,041	220,685
Black ^b	1,301	938	1,012	619	391	306	4,042	4,941	160	5,101
Asian/Pacific Islander ^c	909	647	626	+	+	+	2,666	+	+	+
Hispanic ^d	6,548	4,837	3,673	+	+	+	17,653	+	+	+

Estimated prevalence percent^a on January 1, 2012, of the SEER^c population diagnosed in the previous 20 years
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f All Ages	
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+		
<u>Race</u>												
All Races ^c	0.0925%	0.0025%	0.0075%	0.0708%	0.1685%	0.2102%	0.1476%	0.0668%	0.0253%	0.0132%	0.0921%	
White ^c	0.1139%	0.0024%	0.0088%	0.0879%	0.2107%	0.2591%	0.1781%	0.0797%	0.0297%	0.0154%	0.1133%	
Black ^c	0.0200%	0.0016%	0.0025%	0.0143%	0.0316%	0.0490%	0.0397%	0.0147%	0.0069%	-	0.0208%	
Asian/Pacific Islander ^c	0.0311%	0.0044%	0.0045%	0.0268%	0.0519%	0.0647%	0.0469%	0.0222%	0.0083%	-	0.0300%	
Hispanic ^d	0.0675%	0.0027%	0.0106%	0.0773%	0.1477%	0.1359%	0.0843%	0.0344%	0.0143%	-	0.0683%	

^a US 2012 cancer prevalence counts are based on 2012 cancer prevalence proportions from the SEER registries and 1/1/2012 US population estimates based on the average of 2011 and 2012 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 37 years for 1975-2012 SEER 9 data; 20 years for 1992-2012 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 37 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <37 and >=37. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 26.21
Cancer of the Thyroid (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2012
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <20 ^e	0 to <37 ^e	>=37 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	185,714	126,992	81,900	58,468	42,019	32,763	471,409	563,884	37,905	601,789
	Males	40,645	27,654	17,782	12,720	8,569	6,868	102,483	121,802	8,576	130,378
	Females	145,069	99,338	64,118	45,748	33,450	25,895	368,926	442,082	29,329	471,411
White ^b	Both Sexes	159,121	111,235	72,095	51,715	37,904	29,499	410,753	494,191	34,338	528,529
	Males	35,956	24,773	16,097	11,563	7,919	6,193	91,800	109,477	7,199	116,676
	Females	123,165	86,462	55,998	40,152	29,985	23,306	318,953	384,714	27,139	411,853
Black ^b	Both Sexes	12,464	7,829	4,487	3,094	1,808	1,527	28,673	32,855	1,608	34,463
	Males	2,000	1,411	756	571	355	286	4,882	5,582	179	5,761
	Females	10,464	6,418	3,731	2,523	1,453	1,241	23,791	27,273	1,429	28,702
Asian/ Pacific Islander ^c	Both Sexes	9,269	5,797	3,805	+	+	+	21,530	+	+	+
	Males	1,753	994	686	+	+	+	3,862	+	+	+
	Females	7,516	4,803	3,119	+	+	+	17,668	+	+	+
Hispanic ^d	Both Sexes	19,138	12,254	7,605	+	+	+	44,177	+	+	+
	Males	3,249	2,094	1,247	+	+	+	7,414	+	+	+
	Females	15,889	10,160	6,358	+	+	+	36,763	+	+	+

Estimated prevalence percent^a on January 1, 2012, of the SEER^c population diagnosed in the previous 20 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted ^f	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.1371%	0.0001%	0.0051%	0.0404%	0.1320%	0.2244%	0.2707%	0.2767%	0.2548%	0.1680%	0.1318%
	Males	0.0592%	-	0.0020%	0.0119%	0.0425%	0.0853%	0.1173%	0.1506%	0.1561%	0.1109%	0.0584%
	Females	0.2130%	-	0.0084%	0.0703%	0.2220%	0.3624%	0.4175%	0.3907%	0.3347%	0.2018%	0.2018%
White ^c	Both Sexes	0.1500%	0.0002%	0.0058%	0.0464%	0.1473%	0.2462%	0.2922%	0.2881%	0.2626%	0.1640%	0.1420%
	Males	0.0667%	-	0.0023%	0.0140%	0.0464%	0.0947%	0.1300%	0.1624%	0.1678%	0.1113%	0.0638%
	Females	0.2330%	-	0.0095%	0.0811%	0.2536%	0.4028%	0.4526%	0.4054%	0.3413%	0.1955%	0.2197%
Black ^c	Both Sexes	0.0663%	-	0.0013%	0.0132%	0.0565%	0.1153%	0.1432%	0.1794%	0.1694%	0.1305%	0.0718%
	Males	0.0233%	-	-	0.0035%	0.0154%	0.0379%	0.0499%	0.0784%	0.0786%	0.0934%	0.0279%
	Females	0.1054%	-	0.0016%	0.0225%	0.0924%	0.1836%	0.2245%	0.2587%	0.2329%	0.1486%	0.1083%
Asian/ Pacific Islander ^c	Both Sexes	0.1282%	-	0.0049%	0.0326%	0.1155%	0.1990%	0.2473%	0.2679%	0.2591%	0.2070%	0.1224%
	Males	0.0485%	-	0.0017%	0.0071%	0.0410%	0.0710%	0.0930%	0.1211%	0.1337%	0.1075%	0.0490%
	Females	0.2011%	-	0.0082%	0.0581%	0.1823%	0.3122%	0.3800%	0.3892%	0.3562%	0.2678%	0.1851%
Hispanic ^d	Both Sexes	0.0868%	-	0.0054%	0.0334%	0.1010%	0.1639%	0.2287%	0.2540%	0.2483%	0.1739%	0.1106%
	Males	0.0286%	-	0.0021%	0.0073%	0.0267%	0.0480%	0.0799%	0.1079%	0.1450%	0.1038%	0.0416%
	Females	0.1463%	-	0.0089%	0.0623%	0.1805%	0.2848%	0.3713%	0.3794%	0.3248%	0.2159%	0.1767%

^a US 2012 cancer prevalence counts are based on 2012 cancer prevalence proportions from the SEER registries and 1/1/2012 US population estimates based on the average of 2011 and 2012 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 37 years for 1975-2012 SEER 9 data; 20 years for 1992-2012 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 37 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <37 and >=37. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 27.21
Cancer of the Urinary Bladder (Invasive and *In Situ*)

Estimated United States Cancer Prevalence Counts^a on January 1, 2012
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <20 ^e	0 to <37 ^e	>=37 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	201,191	137,251	93,700	60,683	36,747	22,208	509,716	567,531	9,872	577,403
	Males	153,198	102,629	69,335	44,678	27,365	16,037	382,334	424,112	6,373	430,485
	Females	47,993	34,622	24,365	16,005	9,382	6,171	127,382	143,419	3,499	146,918
White ^b	Both Sexes	183,732	127,045	87,671	57,113	35,092	21,111	471,627	526,751	7,746	534,497
	Males	140,619	95,528	65,102	41,940	26,132	15,166	355,088	394,869	4,905	399,774
	Females	43,113	31,517	22,569	15,173	8,960	5,945	116,539	131,882	2,841	134,723
Black ^b	Both Sexes	9,462	5,372	3,390	2,240	1,025	654	20,936	22,532	171	22,703
	Males	6,575	3,529	2,191	1,740	758	540	14,335	15,551	139	15,690
	Females	2,887	1,843	1,199	500	267	114	6,601	6,981	32	7,013
Asian/ Pacific Islander ^c	Both Sexes	3,723	2,419	1,330	+	+	+	8,273	+	+	+
	Males	2,817	1,874	961	+	+	+	6,271	+	+	+
	Females	906	545	369	+	+	+	2,002	+	+	+
Hispanic ^d	Both Sexes	7,902	4,846	2,952	+	+	+	17,563	+	+	+
	Males	5,846	3,688	2,221	+	+	+	13,089	+	+	+
	Females	2,056	1,158	731	+	+	+	4,474	+	+	+

Estimated prevalence percent^a on January 1, 2012, of the SEER^c population diagnosed in the previous 20 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Race	Sex	Age Specific (Crude)								Age-Adjusted ^f		
			All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
	All Races ^c	Both Sexes	0.1341%	0.0002%	0.0004%	0.0015%	0.0070%	0.0275%	0.1093%	0.3531%	0.8054%	1.1910%	0.1312%
		Males	0.2049%	0.0002%	0.0005%	0.0020%	0.0098%	0.0414%	0.1721%	0.5765%	1.3785%	2.3179%	0.2296%
		Females	0.0650%	-	0.0002%	0.0009%	0.0041%	0.0138%	0.0492%	0.1511%	0.3409%	0.5247%	0.0570%
	White ^c	Both Sexes	0.1618%	0.0002%	0.0004%	0.0017%	0.0085%	0.0320%	0.1243%	0.4052%	0.9282%	1.3163%	0.1490%
		Males	0.2454%	-	0.0006%	0.0024%	0.0117%	0.0469%	0.1927%	0.6519%	1.5715%	2.5648%	0.2581%
		Females	0.0784%	-	-	0.0010%	0.0050%	0.0166%	0.0566%	0.1748%	0.3936%	0.5697%	0.0647%
	Black ^c	Both Sexes	0.0496%	-	-	-	0.0030%	0.0164%	0.0663%	0.1838%	0.3749%	0.5505%	0.0647%
		Males	0.0721%	-	-	-	0.0047%	0.0263%	0.1086%	0.3101%	0.6198%	0.9906%	0.1104%
		Females	0.0291%	-	-	-	0.0014%	0.0076%	0.0295%	0.0846%	0.2034%	0.3369%	0.0343%
	Asian/ Pacific Islander ^c	Both Sexes	0.0576%	-	-	-	0.0026%	0.0101%	0.0472%	0.1437%	0.3333%	0.6561%	0.0595%
		Males	0.0906%	-	-	-	0.0033%	0.0170%	0.0781%	0.2450%	0.6012%	1.2223%	0.1068%
		Females	0.0274%	-	-	-	0.0021%	0.0040%	0.0205%	0.0600%	0.1262%	0.3102%	0.0254%
	Hispanic ^d	Both Sexes	0.0344%	-	-	0.0013%	0.0036%	0.0147%	0.0503%	0.1615%	0.3976%	0.6415%	0.0660%
		Males	0.0505%	-	-	0.0016%	0.0050%	0.0195%	0.0774%	0.2613%	0.7170%	1.2369%	0.1171%
		Females	0.0180%	-	-	0.0009%	0.0023%	0.0096%	0.0243%	0.0758%	0.1610%	0.2852%	0.0295%

^a US 2012 cancer prevalence counts are based on 2012 cancer prevalence proportions from the SEER registries and 1/1/2012 US population estimates based on the average of 2011 and 2012 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 37 years for 1975-2012 SEER 9 data; 20 years for 1992-2012 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 37 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <37 and >=37. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 28.12
Childhood Cancer (Invasive, <20 Years old at diagnosis)

Estimated United States Cancer Prevalence Counts^a on January 1, 2012
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <20 ^e	0 to <37 ^e	>=37 ^f	Complete ^g
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	65,893	56,052	50,529	45,384	38,463	33,067	234,055	326,769	+	+
	Males	34,508	29,895	26,046	23,607	19,837	16,567	122,721	167,928	+	+
	Females	31,385	26,157	24,483	21,777	18,626	16,500	111,334	158,841	+	+
White ^b	Both Sexes	53,100	46,757	42,637	37,801	32,859	29,117	194,105	275,292	+	+
	Males	27,958	25,069	21,916	19,706	17,019	14,676	102,124	141,990	+	+
	Females	25,142	21,688	20,721	18,095	15,840	14,441	91,981	133,302	+	+
Black ^b	Both Sexes	7,885	5,822	4,937	4,699	3,602	2,582	24,832	32,639	+	+
	Males	4,009	3,034	2,544	2,375	1,763	1,223	12,692	16,202	+	+
	Females	3,876	2,788	2,393	2,324	1,839	1,359	12,140	16,437	+	+
Asian/ Pacific Islander ^c	Both Sexes	2,887	2,146	1,848	+	+	+	8,581	+	+	+
	Males	1,517	1,118	1,005	+	+	+	4,497	+	+	+
	Females	1,370	1,028	843	+	+	+	4,084	+	+	+
Hispanic ^d	Both Sexes	13,941	10,460	8,359	+	+	+	39,491	+	+	+
	Males	7,467	5,741	4,453	+	+	+	21,305	+	+	+
	Females	6,474	4,719	3,906	+	+	+	18,186	+	+	+

^a US 2012 cancer prevalence counts are based on 2012 cancer prevalence proportions from the SEER registries and 1/1/2012 US population estimates based on the average of 2011 and 2012 population estimates from the US Bureau of the Census.

^{b c d} Prevalence was calculated using the First Malignant Primary Only for a person. Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 37 years for 1975-2012 SEER 9 data; 20 years for 1992-2012 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^{f g} (f) Cases diagnosed more than 37 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <37 and >=37.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 29.7
U.S. Childhood Cancer Survivors at January 1, 2012
Number of People Previously Diagnosed with Cancer as Children (Ages 0-19 Years) in the United States and Alive January 1, 2012
By International Classification of Childhood Cancer (ICCC)^a Selected Group and Subgroup, Age Group, and Sex

Site	Sex	Complete Prevalence Counts by Age at Prevalence							All Ages		
		0-19	20-29	30-39	40-49	50-59	60+	All Ages	Diagnosed between 1975-2011	Diagnosed before 1975	% ^b
									Count	Count	% ^b
All Sites	Both	115,689	98,695	65,873	55,981	41,852	20,877	398,967	326,512	72,455	18
	Male	62,037	51,574	32,716	25,507	17,574	10,684	200,092	168,045	32,047	16
	Female ^c	53,652	47,121	33,157	30,474	24,278	10,193	198,875	158,467	40,408	20
I Leukemia	Both	36,451	21,624	12,675	5,684	1,981	266	78,681	74,399	4,282	5
	Male	20,275	11,238	6,458	2,955	936	116	41,978	40,061	1,917	5
	Female	16,176	10,386	6,217	2,729	1,045	150	36,703	34,338	2,365	6
Acute lymphocytic leukemia ^d	Both	30,816	17,704	10,733	5,036	840	8	65,137	61,916	3,221	5
	Male	17,122	9,418	5,618	2,799	465	4	35,426	33,759	1,667	5
	Female	13,694	8,286	5,115	2,237	375	4	29,711	28,157	1,554	5
I(b) Acute myeloid leukemia	Both	3,991	2,609	1,236	762	283	7	8,888	8,389	499	6
	Male	2,135	1,183	508	356	154	5	4,341	4,077	264	6
	Female	1,856	1,426	728	406	129	2	4,547	4,312	235	5
II(a) Hodgkin lymphoma	Both	4,686	9,188	8,009	7,778	5,470	1,603	36,734	31,454	5,280	14
	Male	2,667	4,625	4,018	3,845	2,678	820	18,653	15,828	2,825	15
	Female	2,019	4,563	3,991	3,933	2,792	783	18,081	15,626	2,455	14
II(b,c,e) Non-Hodgkin lymphoma	Both	6,848	7,511	4,846	3,366	1,946	1,295	25,812	22,302	3,510	14
	Male	4,523	4,905	3,321	2,150	1,022	748	16,669	14,771	1,898	11
	Female	2,325	2,606	1,525	1,216	924	547	9,143	7,531	1,612	18
III CNS and misc intracranial & intraspinal neoplasms	Both	21,183	15,559	9,245	8,586	5,494	1,997	62,064	51,221	10,843	17
	Male	11,260	8,627	5,046	4,342	2,464	911	32,650	27,727	4,923	15
	Female	9,923	6,932	4,199	4,244	3,030	1,086	29,414	23,494	5,920	20
IV Neuroblastoma & other peripheral nervous cell tumor	Both	9,642	4,053	2,315	1,874	1,602	907	20,393	16,178	4,215	21
	Male	4,976	2,076	1,064	588	557	216	9,477	8,172	1,305	14
	Female	4,666	1,977	1,251	1,286	1,045	691	10,916	8,006	2,910	27
VI Renal tumors	Both	7,849	4,793	3,712	4,483	2,757	991	24,585	16,656	7,929	32
	Male	3,819	2,264	1,640	2,534	1,496	577	12,330	7,810	4,520	37
	Female	4,030	2,529	2,072	1,949	1,261	414	12,255	8,846	3,409	28
VIII Malignant bone tumors	Both	3,642	4,433	3,275	2,535	2,524	1,048	17,457	13,938	3,519	20
	Male	2,124	2,628	1,725	1,318	1,106	548	9,449	7,830	1,619	17
	Female	1,518	1,805	1,550	1,217	1,418	500	8,008	6,108	1,900	24
VIII(a) Osteosarcoma	Both	1,853	2,331	1,782	1,354	1,252	874	9,446	7,376	2,070	22
	Male	1,044	1,307	969	689	543	411	4,963	4,022	941	19
	Female	809	1,024	813	665	709	463	4,483	3,354	1,129	25
VIII(c) Ewing tumor & related sarcomas of bone	Both	1,410	1,424	1,028	757	383	168	5,170	4,581	589	11
	Male	850	877	478	377	194	60	2,836	2,593	243	9
	Female	560	547	550	380	189	108	2,334	1,988	346	15
IX Soft tissue & other extraosseous sarcomas	Both	6,987	7,139	4,429	4,833	4,423	4,952	32,763	22,369	10,394	32
	Male	3,889	3,847	2,234	2,424	2,094	2,777	17,265	11,894	5,371	31
	Female	3,098	3,292	2,195	2,409	2,329	2,175	15,498	10,475	5,023	32
X Germ cell & trophoblastic tumors & neoplasms of gonads	Both	5,233	7,977	5,821	6,018	6,842	7,142	39,033	25,272	13,761	35
	Male	2,702	5,252	3,597	3,211	3,381	3,296	21,439	14,942	6,497	30
	Female	2,531	2,725	2,224	2,807	3,461	3,846	17,594	10,330	7,264	41

^a International Classification of Childhood Cancer is based on ICD-O-3. Steliarova-Foucher E, Stiller C, Lacour B, Kaatsch P. International Classification of Childhood Cancer, Third Edition. Cancer. April 1, 2005; Vol 103, No. 7, pg 1457-1467.

^b Percent diagnosed before 1975 of all ages complete prevalence counts.

^c All sites females excludes breast cancer.

^d SEER site recode Acute Lymphocytic Leukemia was used instead of the ICCC classification lymphocytic leukemia (Ia) because of statistical modeling requirements.

For more information regarding the modeling of the prevalence estimates, see Mariotto et al. Long-Term Survivors of Childhood Cancers in the United States. Cancer Epidemiol Biomarkers Prev 2009, page 1034.