

Table 1.23  
U.S. Complete Prevalence Counts, Invasive Cancers Only, January 1, 2012<sup>a</sup>  
By Age at Prevalence

Site/Sex	Age at Prevalence								
	All Ages <sup>c</sup>	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
All Sites									
Males	6,464,529	19,057	42,522	87,325	156,133	356,474	907,855	1,732,881	3,162,283
Females	7,311,722	17,006	36,354	89,937	229,430	636,759	1,353,262	1,832,064	3,116,909
Oral Cavity & Pharynx									
Males	193,779	73	416	1,506	3,443	14,067	46,864	62,089	65,320
Females	97,329	96	569	1,562	3,831	8,690	18,524	24,531	39,526
Esophagus									
Males	27,808	0	0	34	126	993	4,575	10,168	11,912
Females	7,973	0	0	11	11	207	1,315	2,059	4,369
Stomach									
Males	43,975	0	46	105	596	2,524	7,100	12,229	21,374
Females	32,854	3	23	172	691	2,440	4,745	7,157	17,623
Colon & Rectum									
Males	579,901	11	93	1,215	5,831	26,335	84,353	142,161	319,901
Females	589,028	0	105	1,463	5,773	23,768	72,770	115,560	369,590
Liver & Intrahep									
Males	36,063	560	460	615	605	1,716	11,082	13,605	7,421
Females	14,671	438	522	404	506	925	3,227	4,037	4,612
Pancreas									
Males	22,783	34	18	152	297	1,722	4,835	7,158	8,566
Females	22,919	0	33	237	479	1,550	4,120	6,272	10,228
Larynx									
Males	71,414	0	0	91	244	1,657	10,356	21,554	37,512
Females	17,438	0	0	47	128	806	3,474	4,783	8,200
Lung & Bronchus									
Males	184,230	34	75	504	1,243	5,258	24,519	55,147	97,451
Females	224,578	34	92	414	1,489	6,994	30,650	60,573	124,332
Melanoma of the Skin									
Males	489,614	45	685	5,156	17,606	45,367	99,463	137,601	183,691
Females	506,973	102	812	10,505	34,549	73,225	119,346	119,937	148,497

<sup>a</sup> U.S. 2012 cancer prevalence counts are based on 2012 cancer prevalence proportions from the SEER 9 registries (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta) and 1/1/2012 U.S. population estimates based on the average of 2011 and 2012 population estimates from the U.S. Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b</sup> Cases diagnosed more than 37 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

<sup>c</sup> Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.

Table 1.23 - continued  
 U.S. Complete Prevalence Counts, Invasive Cancers Only, January 1, 2012<sup>a</sup>  
 By Age at Prevalence

Site/Sex	Age at Prevalence								
	All Ages <sup>c</sup>	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
Breast									
Males	15,499	0	0	11	134	699	1,855	4,533	8,267
Females	2,975,314	11	35	3,037	35,604	214,136	559,716	824,053	1,338,721
Cervix									
Females	249,512	0	68	2,062	15,420	41,029	59,636	58,895	72,402
Corpus & Uterus, NOS									
Females	621,612	11	23	614	5,890	24,821	91,757	174,495	324,000
Ovary <sup>d</sup>									
Females	192,446	77	1,080	3,491	6,874	18,099	42,498	51,208	69,119
Prostate									
Males	2,795,592	34	47	102	293	20,286	240,483	807,991	1,726,357
Urinary Bladder									
Males	430,485	56	117	541	2,355	10,502	43,269	104,139	269,506
Females	146,918	45	34	208	1,046	4,028	13,688	31,217	96,652
Kidney & Renal Pelvis									
Males	223,683	1,445	2,452	2,765	4,922	19,349	43,876	64,800	84,074
Females	152,242	1,519	2,612	2,882	4,794	12,658	26,698	37,929	63,150
Hodgkin Lymphoma									
Males	98,082	237	2,413	9,323	16,360	23,948	22,943	15,053	7,806
Females	91,544	39	1,969	9,432	16,470	22,866	20,748	12,271	7,749
Non-Hodgkin Lymphoma									
Males	289,004	794	3,674	8,038	13,729	29,671	55,734	74,542	102,823
Females	260,621	475	1,812	4,755	8,981	21,174	43,182	63,503	116,739
Myeloma									
Males	48,074	0	6	121	521	3,285	8,685	16,099	19,356
Females	41,584	0	0	28	392	2,271	7,446	12,033	19,413
Leukemia									
Males	180,231	6,723	13,446	13,043	11,701	14,648	25,247	36,927	58,495
Females	138,158	5,979	10,128	12,020	10,156	10,667	16,951	24,384	47,873
Acute Lymphocytic Leuk									
Males	41,055	5,618	11,450	9,948	6,685	4,256	1,672	861	564
Females	34,121	5,033	8,635	8,663	5,668	3,311	1,318	916	579

<sup>a</sup> U.S. 2012 cancer prevalence counts are based on 2012 cancer prevalence proportions from the SEER 9 registries (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta) and 1/1/2012 U.S. population estimates based on the average of 2011 and 2012 population estimates from the U.S. Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b</sup> Cases diagnosed more than 37 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

<sup>c</sup> Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.