

Table 1.25
Age-Adjusted SEER Incidence Rates and Trends for the Top 15 Cancer Sites^a by Race/Ethnicity

			Males					
All Races			White			Black		
	Rate ^b	APC ^c		Rate ^b	APC ^c		Rate ^b	APC ^c
	2008-2012	2003-2012		2008-2012	2003-2012		2008-2012	2003-2012
All Sites	516.6	-1.3*	All Sites	519.8	-1.3*	All Sites	590.1	-2.1*
Prostate	137.9	-2.9*	Prostate	130.4	-3.2*	Prostate	214.5	-2.9*
Lung and Bronchus	70.1	-2.6*	Lung and Bronchus	70.3	-2.5*	Lung and Bronchus	90.9	-2.9*
Colon and Rectum	48.9	-3.1*	Colon and Rectum	47.8	-3.3*	Colon and Rectum	61.2	-3.0*
Urinary Bladder	35.8	-1.0*	Urinary Bladder	39.0	-1.0*	Kidney and Renal Pelvis	25.1	1.7*
Melanoma of the Skin	28.2	1.5*	Melanoma of the Skin	33.0	1.4*	Urinary Bladder	21.4	-0.3
Non-Hodgkin Lymphoma	23.9	-0.2	Non-Hodgkin Lymphoma	24.9	-0.2	Non-Hodgkin Lymphoma	17.8	-0.8
Kidney and Renal Pelvis	21.3	1.5*	Kidney and Renal Pelvis	21.9	1.5*	Pancreas	17.2	-0.3
Leukemia	17.0	0.0	Leukemia	17.9	0.0	Liver & IBD ^f	16.2	3.4*
Oral Cavity and Pharynx	16.5	0.5*	Oral Cavity and Pharynx	17.1	0.9*	Myeloma	15.1	0.1
Pancreas	14.0	0.8*	Pancreas	14.0	0.9*	Oral Cavity and Pharynx	14.6	-2.4*
Liver & IBD ^f	12.7	3.2*	Liver & IBD ^f	11.2	3.8*	Stomach	14.6	-3.5*
Stomach	10.1	-1.4*	Stomach	9.2	-0.9*	Leukemia	13.5	-0.6
Myeloma	7.9	1.3*	Brain and ONS ^f	8.4	-0.4	Larynx	8.8	-3.3*
Brain and ONS ^f	7.7	-0.4	Esophagus	8.0	-0.3	Esophagus	7.6	-5.0*
Esophagus	7.6	-1.0*	Myeloma	7.5	1.4*	Brain and ONS ^f	4.9	0.2
Asian/Pacific Islander			American Indian/Alaska Native^d			Hispanic^e		
	Rate ^b	APC ^c		Rate ^b	APC ^c		Rate ^b	APC ^c
	2008-2012	2003-2012		2008-2012	2003-2012		2008-2012	2003-2012
All Sites	325.4	-2.1*	All Sites	340.8	-1.8*	All Sites	395.0	-1.9*
Prostate	74.0	-4.8*	Prostate	67.1	-5.0*	Prostate	114.7	-4.0*
Lung and Bronchus	49.0	-2.3*	Lung and Bronchus	47.6	-3.7*	Colon and Rectum	43.3	-2.4*
Colon and Rectum	42.2	-2.2*	Colon and Rectum	46.3	0.0	Lung and Bronchus	37.9	-3.3*
Liver & IBD ^f	20.9	-1.2	Kidney and Renal Pelvis	24.8	1.0	Non-Hodgkin Lymphoma	20.6	0.3
Non-Hodgkin Lymphoma	16.6	0.1	Liver & IBD ^f	19.9	1.3	Kidney and Renal Pelvis	20.2	0.6
Urinary Bladder	15.8	-1.6*	Urinary Bladder	15.6	-1.0	Urinary Bladder	19.6	-1.5*
Stomach	14.5	-3.9*	Non-Hodgkin Lymphoma	14.6	1.1	Liver & IBD ^f	19.1	2.8*
Kidney and Renal Pelvis	11.6	1.6	Oral Cavity and Pharynx	13.2	4.5	Stomach	14.2	-2.4*
Oral Cavity and Pharynx	11.0	-0.1	Stomach	12.3	-4.1	Leukemia	12.6	-0.3
Pancreas	10.8	0.4	Pancreas	11.1	-	Pancreas	11.9	0.1
Leukemia	9.8	0.3	Leukemia	10.1	2.8	Oral Cavity and Pharynx	10.1	1.1
Thyroid	6.2	6.4*	Esophagus	4.9	-	Myeloma	7.3	0.7
Myeloma	4.6	0.9	Testis	4.8	-	Brain and ONS ^f	5.8	-2.0*
Brain and ONS ^f	4.3	0.3	Myeloma	4.6	-	Esophagus	5.2	-1.2
Esophagus	3.6	-2.8	Larynx	4.3	-	Testis	4.9	2.8*

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Top 15 cancer sites selected based on 2008-2012 age-adjusted rates for the race/ethnic group.

^b Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^c The APC is the Annual Percent Change over the time interval.

Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^d Rates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^e Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

^f IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.

* The APC is significantly different from zero (p<.05).

- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.