

Table 5.7  
Cancer of the Cervix Uteri (Invasive)

SEER Incidence<sup>a</sup> and U.S. Death<sup>b</sup> Rates, Age-Adjusted and Age-Specific Rates, by Race

	SEER Incidence <sup>a</sup>			U.S. Mortality <sup>b</sup>		
	All Races Females	White Females	Black Females	All Races Females	White Females	Black Females
<u>Age-Adjusted Rates, 2008-2012</u>						
Age at Diagnosis/Death:						
All ages	7.7	7.7	9.2	2.3	2.1	4.0
Under 65	7.2	7.3	7.6	1.8	1.7	2.7
65 and over	11.3	10.0	20.2	5.8	5.1	12.4
All ages (IARC world std) <sup>c</sup>	6.1	6.1	6.9	1.7	1.6	2.8
<u>Age-Specific Rates, 2008-2012</u>						
Age at Diagnosis/Death:						
<1	-	-	-	-	-	-
1-4	-	-	-	-	-	-
5-9	-	-	-	-	-	-
10-14	-	-	-	-	-	-
15-19	0.1	0.1	-	-	-	-
20-24	1.3	1.3	1.3	0.1	0.1	0.2
25-29	5.0	5.6	4.3	0.6	0.6	0.8
30-34	10.1	11.1	8.8	1.3	1.2	1.7
35-39	13.6	14.8	11.0	2.3	2.2	3.2
40-44	14.6	15.2	14.5	3.1	3.0	4.3
45-49	13.6	13.4	14.7	4.0	3.8	6.2
50-54	12.7	12.2	16.4	4.5	4.1	7.1
55-59	11.8	11.3	16.1	4.6	4.3	7.3
60-64	12.3	11.6	17.5	5.2	4.8	8.8
65-69	12.5	11.8	18.0	5.4	4.8	10.5
70-74	12.2	11.0	17.4	5.6	5.0	10.9
75-79	10.3	9.2	20.2	5.7	5.0	12.7
80-84	9.9	8.4	22.7	6.3	5.4	14.9
85+	9.8	7.8	28.0	6.8	5.9	16.2

<sup>a</sup> SEER 18 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

<sup>b</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

<sup>c</sup> Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

- Rates are per 100,000 and are age-adjusted to the IARC world standard population.

- Statistic not shown. Rate based on less than 16 cases for the time interval.