


Table 5.14
Cancer of the Cervix Uteri (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012


Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	2.31	0.02		
High Five States				
Arkansas	3.39 ^b	0.21	01(01,10)	47.14 ^c
Mississippi	3.37 ^b	0.21	02(01,11)	46.10 ^c
West Virginia	3.29 ^b	0.26	03(01,17)	42.78 ^c
Oklahoma	3.16 ^b	0.18	04(01,14)	37.07 ^c
Alabama	3.03 ^b	0.15	05(01,17)	31.16 ^c
Low Five States				
Massachusetts	1.40 ^b	0.09	47(40,51)	-39.29 ^c
Rhode Island	1.34 ^b	0.21	48(30,51)	-41.89 ^c
Utah	1.29 ^b	0.15	49(38,51)	-44.02 ^c
Vermont	1.20 ^b	0.26	50(29,51)	-47.85 ^c
North Dakota	1.16 ^b	0.25	51(28,51)	-49.52 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	3.03 ^b	0.15	05(01,17)	31.16 ^c	Montana	1.51	0.24	45(21,51)	-34.56 ^c
Alaska	2.03	0.36	30(03,51)	-11.88	Nebraska	1.95	0.21	33(14,48)	-15.66 ^c
Arizona	1.98	0.11	32(21,42)	-14.11	Nevada	2.20	0.18	23(10,40)	-4.51
Arkansas	3.39 ^b	0.21	01(01,10)	47.14 ^c	New Hampshire	1.64	0.21	41(22,51)	-28.72 ^c
California	2.23	0.05	22(18,30)	-3.37	New Jersey	2.38	0.10	20(12,29)	3.15
Colorado	1.53 ^b	0.11	44(35,50)	-33.77 ^c	New Mexico	2.14	0.20	25(10,44)	-7.02
Connecticut	1.66 ^b	0.13	40(28,49)	-28.11 ^c	New York	2.29	0.06	21(16,30)	-0.75
Delaware	2.56	0.33	17(01,41)	10.80	North Carolina	2.05	0.09	29(21,39)	-11.32
D.C.	2.84	0.43	09(01,40)	23.01 ^c	North Dakota	1.16 ^b	0.25	51(28,51)	-49.52 ^c
Florida	2.71 ^b	0.07	13(06,18)	17.48 ^c	Ohio	2.57	0.09	15(08,22)	11.30
Georgia	2.71 ^b	0.10	12(05,20)	17.60 ^c	Oklahoma	3.16 ^b	0.18	04(01,14)	37.07 ^c
Hawaii	1.83	0.22	38(16,50)	-20.68 ^c	Oregon	1.88	0.13	37(21,45)	-18.42 ^c
Idaho	1.90	0.22	35(14,49)	-17.41 ^c	Pennsylvania	2.13	0.08	26(19,35)	-7.69
Illinois	2.53	0.08	18(09,23)	9.90	Rhode Island	1.34 ^b	0.21	48(30,51)	-41.89 ^c
Indiana	2.41	0.12	19(10,29)	4.38	South Carolina	2.73	0.15	11(03,22)	18.20 ^c
Iowa	1.94	0.16	34(19,45)	-15.91 ^c	South Dakota	2.06	0.31	28(05,50)	-10.77
Kansas	1.99	0.17	31(17,45)	-13.53	Tennessee	2.70	0.12	14(05,21)	17.01 ^c
Kentucky	2.95 ^b	0.16	07(02,18)	27.72 ^c	Texas	2.77 ^b	0.07	10(05,16)	20.06 ^c
Louisiana	3.00 ^b	0.16	06(01,17)	30.21 ^c	Utah	1.29 ^b	0.15	49(38,51)	-44.02 ^c
Maine	1.59 ^b	0.20	42(24,51)	-31.21 ^c	Vermont	1.20 ^b	0.26	50(29,51)	-47.85 ^c
Maryland	2.16	0.12	24(16,38)	-6.28	Virginia	1.90 ^b	0.09	36(23,42)	-17.54 ^c
Massachusetts	1.40 ^b	0.09	47(40,51)	-39.29 ^c	Washington	1.81 ^b	0.10	39(27,45)	-21.70 ^c
Michigan	2.08	0.09	27(19,38)	-9.62	West Virginia	3.29 ^b	0.26	03(01,17)	42.78 ^c
Minnesota	1.46 ^b	0.10	46(37,51)	-36.82 ^c	Wisconsin	1.53 ^b	0.10	43(35,50)	-33.61 ^c
Mississippi	3.37 ^b	0.21	02(01,11)	46.10 ^c	Wyoming	2.86	0.45	08(01,41)	23.92 ^c
Missouri	2.56	0.13	16(07,25)	10.84					



Age-Adjusted Death
Rates per 100,000
Quantile Interval



- ^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- ^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
- ^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
- ^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.
- SE Standard error of the rate.
- PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.