

Table 5.15
Cancer of the Cervix Uteri (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2012
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <20 ^e	0 to <37 ^e	>=37 ^g	Complete ^h
<u>Race</u>										
All Races ^b	39,323	32,593	34,878	32,772	28,429	21,631	151,928	214,008	35,504	249,512
White ^b	31,133	25,865	27,920	27,182	23,762	17,910	122,201	173,836	30,059	203,895
Black ^b	5,191	4,548	4,549	3,363	2,982	2,737	19,129	26,586	+	+
Asian/Pacific Islander ^c	2,104	1,727	1,678	+	+	+	7,030	+	+	+
Hispanic ^d	7,897	7,205	6,782	+	+	+	27,657	+	+	+

Estimated prevalence percent^a on January 1, 2012, of the SEER^c population diagnosed in the previous 20 years
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>											
All Races ^c	0.0953%	-	0.0002%	0.0087%	0.0723%	0.1825%	0.2039%	0.1846%	0.1608%	0.1031%	0.0896%
White ^c	0.0996%	-	-	0.0101%	0.0830%	0.2026%	0.2119%	0.1762%	0.1457%	0.0853%	0.0934%
Black ^c	0.0801%	-	-	0.0054%	0.0599%	0.1449%	0.1807%	0.1993%	0.1863%	0.1394%	0.0825%
Asian/Pacific Islander ^c	0.0834%	-	-	0.0042%	0.0302%	0.1070%	0.1718%	0.2126%	0.2242%	0.1991%	0.0760%
Hispanic ^d	0.1105%	-	-	0.0088%	0.0783%	0.2340%	0.3404%	0.3503%	0.3416%	0.2635%	0.1419%

^a US 2012 cancer prevalence counts are based on 2012 cancer prevalence proportions from the SEER registries and 1/1/2012 US population estimates based on the average of 2011 and 2012 population estimates from the US Bureau of the Census.
^{b c d} Prevalence was calculated using the First Malignant Primary Only for a person. Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.
^e Maximum limited-duration prevalence: 37 years for 1975-2012 SEER 9 data; 20 years for 1992-2012 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).
^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.
^{g h i} (g) Cases diagnosed more than 37 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <37 and >=37. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.
- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.
+ Not available.