


Table 6.25
Cancer of the Colon and Rectum (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	15.50	0.03		
High Five States				
Mississippi	19.69 ^b	0.36	01(01,03)	27.01 ^c
Arkansas	18.54 ^b	0.34	02(01,08)	19.65 ^c
Louisiana	18.46 ^b	0.28	03(01,07)	19.12 ^c
West Virginia	18.43 ^b	0.40	04(01,10)	18.89 ^c
Kentucky	18.10 ^b	0.28	05(02,10)	16.76 ^c
Low Five States				
Arizona	13.44 ^b	0.20	47(43,50)	-13.31 ^c
Hawaii	13.40 ^b	0.41	48(36,50)	-13.54 ^c
Colorado	13.37 ^b	0.24	49(42,50)	-13.76 ^c
Connecticut	12.68 ^b	0.25	50(47,51)	-18.16 ^c
Utah	11.62 ^b	0.34	51(50,51)	-25.01 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	17.19 ^b	0.26	09(05,18)	10.94 ^c	Montana	14.22	0.49	42(23,50)	-8.22
Alaska	15.84	0.86	21(03,48)	2.19	Nebraska	16.96 ^b	0.41	12(04,23)	9.41
Arizona	13.44 ^b	0.20	47(43,50)	-13.31 ^c	Nevada	17.15 ^b	0.37	10(04,21)	10.67 ^c
Arkansas	18.54 ^b	0.34	02(01,08)	19.65 ^c	New Hampshire	14.43	0.44	39(22,49)	-6.90
California	14.22 ^b	0.09	43(36,46)	-8.27	New Jersey	16.26 ^b	0.18	18(13,24)	4.92
Colorado	13.37 ^b	0.24	49(42,50)	-13.76 ^c	New Mexico	14.82	0.37	34(21,46)	-4.37
Connecticut	12.68 ^b	0.25	50(47,51)	-18.16 ^c	New York	15.08 ^b	0.12	31(24,37)	-2.71
Delaware	14.60	0.54	37(19,49)	-5.81	North Carolina	14.75 ^b	0.17	35(27,42)	-4.85
D.C.	17.47	0.78	08(01,29)	12.72 ^c	North Dakota	15.97	0.63	20(05,43)	3.07
Florida	14.40 ^b	0.11	40(33,45)	-7.07	Ohio	17.00 ^b	0.16	11(07,18)	9.70
Georgia	15.76	0.19	22(18,29)	1.66	Oklahoma	17.60 ^b	0.30	06(03,16)	13.58 ^c
Hawaii	13.40 ^b	0.41	48(36,50)	-13.54 ^c	Oregon	14.86	0.26	33(23,44)	-4.13
Idaho	13.75 ^b	0.42	46(30,50)	-11.31 ^c	Pennsylvania	16.73 ^b	0.15	14(09,20)	7.94
Illinois	16.68 ^b	0.16	17(09,21)	7.64	Rhode Island	15.08	0.49	30(16,47)	-2.70
Indiana	16.73 ^b	0.22	15(07,21)	7.92	South Carolina	16.18	0.26	19(11,27)	4.42
Iowa	16.71 ^b	0.30	16(06,23)	7.82	South Dakota	15.69	0.57	23(08,43)	1.26
Kansas	15.52	0.32	25(17,37)	0.16	Tennessee	17.56 ^b	0.23	07(04,13)	13.27 ^c
Kentucky	18.10 ^b	0.28	05(02,10)	16.76 ^c	Texas	15.38	0.12	26(21,32)	-0.77
Louisiana	18.46 ^b	0.28	03(01,07)	19.12 ^c	Utah	11.62 ^b	0.34	51(50,51)	-25.01 ^c
Maine	15.19	0.43	28(17,45)	-2.02	Vermont	14.65	0.63	36(16,49)	-5.46
Maryland	15.36	0.23	27(20,36)	-0.90	Virginia	15.12	0.20	29(22,38)	-2.44
Massachusetts	14.27 ^b	0.20	41(32,47)	-7.90	Washington	14.09 ^b	0.20	44(34,48)	-9.10
Michigan	15.57	0.17	24(19,32)	0.46	West Virginia	18.43 ^b	0.40	04(01,10)	18.89 ^c
Minnesota	13.94 ^b	0.22	45(36,49)	-10.03 ^c	Wisconsin	14.52 ^b	0.21	38(28,45)	-6.30
Mississippi	19.69 ^b	0.36	01(01,03)	27.01 ^c	Wyoming	14.98	0.72	32(09,49)	-3.33
Missouri	16.77 ^b	0.22	13(07,21)	8.20					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 10% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.