

Table 8.9
Cancer of the Esophagus (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males and Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Survival Time																					
1-year	26.0	29.8	34.7	38.3	40.7	41.9	40.1	40.5	44.4	45.6	46.2	41.8	46.8	48.2	45.2	47.6	50.2	47.7	48.5	49.5	49.2
2-year	11.4	14.3	19.0	21.8	22.9	24.2	23.7	24.2	26.9	28.3	29.1	25.9	30.4	30.2	31.2	30.5	33.1	32.2	30.6	34.0	
3-year	7.7	10.0	13.5	16.3	16.8	18.6	17.7	18.4	20.8	23.6	23.1	20.9	23.9	25.8	24.2	25.2	26.9	24.3	24.4		
4-year	5.9	8.2	10.8	13.7	13.3	15.9	14.4	14.2	18.3	20.6	19.3	17.9	20.0	22.3	21.4	21.8	23.3	21.6			
5-year	4.9	7.0	9.5	12.2	11.6	15.0	12.5	12.6	17.4	18.9	17.9	16.8	17.1	20.3	18.9	20.0	21.6				
6-year	4.3	6.1	8.5	11.1	10.8	13.8	11.4	11.7	15.9	17.9	15.8	15.3	14.3	19.8	17.4	18.6					
7-year	3.8	5.6	7.7	10.2	10.2	12.9	10.8	11.3	14.8	16.7	14.8	14.9	13.1	18.3	15.6						
8-year	3.4	4.9	7.1	9.5	9.7	12.3	9.9	9.8	14.3	15.8	13.8	13.6	12.8	17.9							
9-year	3.2	4.5	6.4	9.0	8.9	11.5	9.5	9.6	13.4	15.5	13.5	13.1	12.6								
10-year	2.9	4.3	6.0	8.3	8.5	10.7	9.3	9.4	12.9	15.1	13.3	13.0									
11-year	2.7	3.9	5.4	7.8	7.7	9.7	9.1	9.1	12.7	14.2	12.2										
12-year	2.6	3.8	4.9	7.1	7.2	9.4	7.7	8.5	11.8	14.0											
13-year	2.4	3.5	4.6	7.0	6.9	9.4	7.1	8.5	10.9												
14-year	2.4	3.4	4.3	6.6	5.7	9.0	6.5	8.4													
15-year	2.4	3.4	4.2	6.2	5.4	8.7	6.2														
16-year	2.3	3.2	3.7	5.9	4.8	7.8															
17-year	2.3	3.0	3.2	5.5	4.8																
18-year	2.2	2.7	3.1	5.3																	
19-year	2.2	2.5	2.9																		
20-year	2.2	2.4	2.6																		

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).