

Table 8.10  
Cancer of the Esophagus (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Survival Time																					
1-year	24.0	29.2	33.9	38.7	40.3	43.0	40.7	40.9	43.8	46.9	47.1	43.1	46.3	47.6	46.3	48.5	51.8	48.7	49.8	48.4	48.7
2-year	10.0	13.8	18.4	21.6	23.0	24.6	24.7	24.6	26.4	28.7	29.4	26.4	29.9	30.2	31.1	30.9	33.4	32.8	30.9	33.2	
3-year	6.8	9.2	13.1	16.4	16.7	18.2	18.5	18.6	20.6	23.5	23.2	21.3	23.2	25.8	24.1	25.9	27.3	24.7	25.2		
4-year	5.1	7.3	10.3	13.4	13.2	15.0	15.3	14.2	18.0	19.6	19.4	18.1	19.5	22.4	20.8	22.8	23.5	21.9			
5-year	4.2	6.0	9.3	12.0	11.8	13.9	13.2	12.1	17.2	17.3	18.4	16.8	16.7	20.0	18.4	21.0	21.9				
6-year	3.7	5.1	8.4	10.8	11.0	12.9	11.9	11.5	16.0	16.4	16.2	15.4	13.8	19.3	17.1	19.4					
7-year	3.3	4.5	7.6	10.2	10.7	11.9	11.6	11.2	14.9	15.4	15.4	15.3	12.6	18.3	15.2						
8-year	3.0	4.0	7.0	9.3	10.1	11.4	10.5	9.4	14.7	14.6	14.6	13.7	12.1	17.9							
9-year	2.8	3.6	6.5	8.8	9.6	10.6	10.2	9.3	14.0	14.3	14.2	13.3	12.1								
10-year	2.7	3.4	6.2	7.9	9.2	9.9	9.9	9.3	13.2	14.0	13.8	13.3									
11-year	2.5	3.3	5.7	7.5	8.4	8.9	9.7	9.0	13.1	13.1	13.0										
12-year	2.4	3.2	5.0	6.8	7.7	8.6	8.6	8.8	12.7	12.8											
13-year	2.3	3.0	4.6	6.8	7.2	8.6	8.0	8.8	11.7												
14-year	2.2	3.0	4.2	6.6	6.3	8.3	7.1	8.8													
15-year	2.2	3.0	4.0	6.1	5.9	8.3	6.6														
16-year	2.2	2.8	3.4	5.9	5.3	8.0															
17-year	2.2	2.8	2.9	5.5	5.3																
18-year	2.2	2.5	2.8	5.3																	
19-year	2.1	2.3	2.5																		
20-year	2.1	2.2	2.3																		

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).