


Table 8.19
Cancer of the Esophagus (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012


Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	7.45	0.03		
High Five States				
New Hampshire	10.13 ^b	0.56	01(01,13)	35.94 ^c
Maine	9.85 ^b	0.51	02(01,14)	32.15 ^c
Ohio	9.14 ^b	0.18	03(01,13)	22.62 ^c
West Virginia	9.09 ^b	0.42	04(01,23)	21.93 ^c
Indiana	8.97 ^b	0.24	05(02,16)	20.28 ^c
Low Five States				
Texas	6.10 ^b	0.11	47(45,50)	-18.12 ^c
California	6.05 ^b	0.09	48(45,50)	-18.85 ^c
New Mexico	5.80 ^b	0.34	49(43,51)	-22.26 ^c
Hawaii	5.58 ^b	0.40	50(43,51)	-25.20 ^c
Utah	5.15 ^b	0.33	51(47,51)	-30.95 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	7.30	0.25	37(17,46)	-2.13	Montana	7.40	0.52	33(06,49)	-0.75
Alaska	7.10	0.82	40(02,51)	-4.71	Nebraska	8.35	0.43	13(02,40)	11.95
Arizona	7.10	0.21	41(23,46)	-4.79	Nevada	8.08	0.37	18(04,41)	8.33
Arkansas	6.79	0.30	45(23,49)	-8.87	New Hampshire	10.13 ^b	0.56	01(01,13)	35.94 ^c
California	6.05 ^b	0.09	48(45,50)	-18.85 ^c	New Jersey	7.03	0.18	43(26,46)	-5.72
Colorado	7.08	0.26	42(20,46)	-5.04	New Mexico	5.80 ^b	0.34	49(43,51)	-22.26 ^c
Connecticut	7.37	0.29	34(15,46)	-1.07	New York	7.13	0.12	39(27,45)	-4.37
Delaware	7.43	0.57	32(03,49)	-0.33	North Carolina	7.00	0.18	44(28,46)	-6.09
D.C.	7.46	0.81	30(01,51)	0.06	North Dakota	8.26	0.70	15(01,46)	10.85
Florida	7.15	0.11	38(28,45)	-4.13	Ohio	9.14 ^b	0.18	03(01,13)	22.62 ^c
Georgia	6.64 ^b	0.19	46(35,48)	-10.89	Oklahoma	8.07	0.30	19(06,36)	8.19
Hawaii	5.58 ^b	0.40	50(43,51)	-25.20 ^c	Oregon	7.65	0.28	23(12,44)	2.65
Idaho	7.33	0.45	35(09,48)	-1.61	Pennsylvania	8.77 ^b	0.16	09(03,17)	17.62 ^c
Illinois	8.01	0.17	21(11,30)	7.41	Rhode Island	8.72	0.59	10(01,41)	16.95 ^c
Indiana	8.97 ^b	0.24	05(02,16)	20.28 ^c	South Carolina	7.65	0.26	24(12,44)	2.62
Iowa	8.78 ^b	0.33	08(02,24)	17.72 ^c	South Dakota	8.96	0.65	06(01,40)	20.22 ^c
Kansas	7.79	0.34	22(08,44)	4.51	Tennessee	7.52	0.22	29(16,44)	0.87
Kentucky	8.02	0.28	20(07,36)	7.56	Texas	6.10 ^b	0.11	47(45,50)	-18.12 ^c
Louisiana	7.62	0.27	27(12,44)	2.26	Utah	5.15 ^b	0.33	51(47,51)	-30.95 ^c
Maine	9.85 ^b	0.51	02(01,14)	32.15 ^c	Vermont	8.22	0.71	16(01,47)	10.30
Maryland	7.32	0.24	36(18,46)	-1.82	Virginia	7.62	0.21	26(14,41)	2.26
Massachusetts	8.94 ^b	0.24	07(02,17)	19.91 ^c	Washington	8.16	0.23	17(07,31)	9.48
Michigan	8.65 ^b	0.19	11(03,19)	16.04 ^c	West Virginia	9.09 ^b	0.42	04(01,23)	21.93 ^c
Minnesota	7.63	0.25	25(12,43)	2.42	Wisconsin	8.38 ^b	0.24	12(04,26)	12.44
Mississippi	7.44	0.33	31(12,46)	-0.18	Wyoming	8.32	0.79	14(01,47)	11.62
Missouri	7.53	0.22	28(15,44)	1.04					



Age-Adjusted Death
Rates per 100,000
Quantile Interval



- ^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- ^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
- ^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
- ^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.
- SE Standard error of the rate.
- PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.