

Table 9.8
Hodgkin Lymphoma

5-Year Relative and Period Survival (Percent) by Race, Sex, Diagnosis Year and Age

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
5-Year Relative Survival (Percent)									
Year of Diagnosis:									
1960-1963 ^a	-	-	-	40	34	48	-	-	-
1970-1973 ^a	-	-	-	67	66	69	-	-	-
1975-1977 ^b	71.8	69.9	74.3	72.1	70.2	74.5	70.1	68.5 ^g	73.3 ^g
1978-1980 ^b	71.7	70.3	73.5	71.8	69.8	74.3	70.6	74.7 ^g	63.1 ^g
1981-1983 ^b	74.2	72.9	75.8	74.6	73.7	75.6	71.8	69.6 ^g	75.9 ^g
1984-1986 ^b	78.2	76.4	80.6	78.7	76.8	81.1	74.7	75.7 ^g	72.1 ^g
1987-1989 ^b	79.2	76.6	82.5	79.7	76.8	83.3	72.3	70.9 ^g	74.0 ^g
1990-1992 ^b	81.7	79.2	84.7	82.8	81.0	85.0	74.2	64.9	84.3
1993-1995 ^b	81.7	79.2	84.7	82.1	80.0	84.7	78.3	74.0	83.0
1996-1998 ^b	85.1	84.4	85.9	85.7	84.3	87.3	81.1	84.3	77.7
1999-2001 ^b	85.0	82.7	87.9	85.7	83.9	87.8	81.3	75.1	89.6
2002-2004 ^b	86.2	85.2	87.2	87.7	87.0	88.4	79.0	74.2	82.6
2005-2011 ^b	88.3 ^f	87.3 ^f	89.5 ^f	88.9 ^f	88.5 ^f	89.1 ^f	85.5 ^f	80.1 ^f	91.6 ^f
5-Year Period Survival (Percent) ^{cd}									
2011	85.8	84.6	87.3	86.3	85.5	87.2	81.9	77.4	87.0
Stage Distribution (%) 2005-2011 ^{ce}									
All Stages									
Number of cases	14,979	8,183	6,796	12,174	6,676	5,498	1,851	989	862
Percent	100%	100%	100%	100%	100%	100%	100%	100%	100%
Localized	16	17	15	16	18	14	19	20	19
Regional	41	36	47	42	36	48	33	27	39
Distant	39	43	34	38	42	33	45	51	39
Unstaged	4	4	5	4	4	5	3	3	3
5-Year Relative Survival (Percent), 2005-2011 ^c									
Age at Diagnosis:									
Ages <45	94.1	93.3	95.1	94.9	94.4	95.4	88.6	84.9	92.7
Ages 45-54	86.0	84.2	88.7	87.4	86.1	89.5	76.7	71.0	85.4
Ages 55-64	77.3	77.7	76.7	79.1	80.0	77.5	63.8	56.4 ^g	70.9 ^g
Ages 65-74	64.3	62.5	66.5	63.8	61.2	67.2	70.1 ^g	77.6 ^g	61.6 ^g
Ages 75+	41.5	39.1	43.1	41.8	38.6	43.8	49.0 ^h	-	39.7 ^h
Ages <65	91.0	89.9	92.4	91.9	91.2	92.8	84.3	80.0	89.3
Ages 65+	53.6	52.9	54.3	53.2	51.5	54.8	65.0 ^g	76.1 ^g	54.0 ^g
Stage ^e :									
All Stages	85.9	85.1	86.9	86.1	85.6	86.8	83.1	79.8	86.6
Localized	91.0	92.1	89.4	91.3	92.3	89.8	88.5	88.4	87.7
Regional	92.8	92.3	93.3	93.1	92.6	93.5	90.0	89.1	90.7
Distant	76.9	76.6	77.4	76.7	76.9	76.2	76.0	71.4	82.6
Unstaged	80.8	78.8	82.6	80.9	79.1	82.5	71.9 ^g	71.5 ^h	-

^a Based on End Results data from a series of hospital registries and one population-based registry.

^b SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta).

Based on follow-up of patients into 2012.

^c SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

Based on follow-up of patients into 2012.

^d Period survival provides a 2011 estimate of survival by piecing together the most recent conditional survival estimates from several cohorts. It is computed here using three year calendar blocks (2009-2011: 0-1 year survival), (2008-2010: 1-2 year survival), (2007-2009: 2-3 year survival), (2006-2008: 3-4 year survival), (2005-2007: 4-5 years survival).

^e Stage at diagnosis is classified using SEER Summary Stage 2000. Stage distribution percentages may not sum to 100 due to rounding.

^f The difference between 1975-1977 and 2005-2011 is statistically significant ($p < .05$).

^g The standard error is between 5 and 10 percentage points.

^h The standard error is greater than 10 percentage points.

- Statistic could not be calculated due to fewer than 25 cases during the time period.