

Table 9.15  
Hodgkin Lymphoma

SEER Incidence and U.S. Mortality  
Age-Adjusted Rates and Trends<sup>a</sup>  
By Race/Ethnicity and Sex

SEER Incidence	SEER 18 Areas <sup>b</sup>			SEER 13 Areas <sup>b</sup>			SEER 9 Areas <sup>b</sup>		
	Rate 2008-2012			Trend 2003-2012 <sup>c</sup>			Trend 2003-2012 <sup>d</sup>		
	Rate per 100,000 persons			AAPC (%)			AAPC (%)		
RACE/ETHNICITY	Total	Males	Females	Total	Males	Females	Total	Males	Females
All Races	2.7	3.0	2.4	-0.6	-0.9	0.0	-0.7	-0.3	0.3*
White	2.9	3.2	2.6	-0.9	0.0	-0.1	-0.8	0.1	0.2
White Hispanic <sup>e</sup>	2.4	2.7	2.1	0.1	-0.5	0.7	-	-	-
White Non-Hispanic <sup>e</sup>	3.1	3.5	2.8	-1.0	0.3	-0.1	-	-	-
Black	2.7	3.0	2.4	0.9*	1.0*	0.6	1.1*	0.5	1.9*
Asian/Pacific Islander	1.3	1.5	1.2	-0.1	1.4	2.6*	-	-	-
Amer Ind/Alaska Nat <sup>f</sup>	1.1	1.1	1.0	-	-	-	-	-	-
Hispanic <sup>e</sup>	2.3	2.6	2.0	0.0	-0.6	0.6	-	-	-
U.S. Mortality <sup>g</sup>	Rate 2008-2012			Trend 2003-2012 <sup>h</sup>			AAPC (%)		
RACE/ETHNICITY	Rate per 100,000 persons			Total	Males	Females	Total	Males	Females
All Races	0.4	0.5	0.3	-2.7*	-2.3*	-2.8*			
White	0.4	0.5	0.3	-2.6*	-2.2*	-2.7*			
White Hispanic <sup>e</sup>	0.4	0.6	0.3	-2.3*	-2.4*	-2.2*			
White Non-Hispanic <sup>e</sup>	0.4	0.5	0.3	-2.6*	-2.6*	-2.7*			
Black	0.3	0.4	0.3	-2.8*	-3.2*	-2.5*			
Asian/Pacific Islander	0.1	0.2	0.1	-1.9	-2.7	-			
Amer Ind/Alaska Nat									
Total U.S.	0.1	-	-	-2.9	-	-			
CHSDA Counties	-	-	-	-1.1	-	-			
Non-CHSDA Counties	-	-	-	-	-	-			
Hispanic <sup>e</sup>	0.4	0.5	0.3	-2.5*	-2.5*	-2.4*			

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 4.2.0, April 2015, National Cancer Institute.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

a Trend based on less than 10 cases for at least one year within the time interval.

b Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

c The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta.

d The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry, and Rural Georgia.

e The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG.

f The 2003-2012 AAPC estimates are based on a Joinpoint analysis with up to 3 Joinpoints over diagnosis years 1992-2012.

g The 2003-2012 AAPC estimates are based on a Joinpoint analysis with up to 5 Joinpoints over diagnosis years 1975-2012.

h Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. The 2003-2012 Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

i Incidence data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

j US Mortality Files, National Center for Health Statistics, CDC.

k The 2003-2012 mortality AAPCs are based on a Joinpoint analysis using years of death 1992-2012.

\* The APC is significantly different from zero (p<.05).