

Table 9.19
Hodgkin Lymphoma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012


Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	0.46	0.01		
High Five States				
Rhode Island	0.72	0.17	01(01,41)	56.20 ^c
New Hampshire	0.68	0.16	02(01,41)	48.19 ^c
Nebraska	0.67	0.12	03(01,39)	45.54 ^c
Louisiana	0.62	0.08	04(01,32)	35.19 ^c
Oregon	0.57	0.08	05(01,37)	24.20 ^c
Low Five States				
Maryland	0.38	0.05	38(07,42)	-17.55 ^c
Colorado	0.34	0.06	39(11,42)	-27.02 ^c
Kentucky	0.34	0.06	40(13,42)	-27.31 ^c
Arizona	0.31 ^b	0.04	41(26,42)	-33.70 ^c
Utah	0.30	0.08	42(08,42)	-34.27 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	0.45	0.06	26(03,42)	-1.60	Montana	-	-	-(-,-)	-
Alaska	-	-	-(-,-)	-	Nebraska	0.67	0.12	03(01,39)	45.54 ^c
Arizona	0.31 ^b	0.04	41(26,42)	-33.70 ^c	Nevada	0.52	0.10	07(01,42)	12.30
Arkansas	0.42	0.08	32(03,42)	-9.21	New Hampshire	0.68	0.16	02(01,41)	48.19 ^c
California	0.49	0.02	16(05,31)	6.55	New Jersey	0.42	0.05	31(06,41)	-9.07
Colorado	0.34	0.06	39(11,42)	-27.02 ^c	New Mexico	0.52	0.10	08(01,42)	11.84
Connecticut	0.38	0.07	37(05,42)	-16.51 ^c	New York	0.50	0.03	14(04,33)	8.67
Delaware	-	-	-(-,-)	-	North Carolina	0.45	0.05	27(05,40)	-2.67
D.C.	-	-	-(-,-)	-	North Dakota	-	-	-(-,-)	-
Florida	0.46	0.03	24(07,37)	0.00	Ohio	0.48	0.04	20(04,38)	3.90
Georgia	0.48	0.05	21(03,39)	3.08	Oklahoma	0.49	0.08	17(01,41)	6.54
Hawaii	-	-	-(-,-)	-	Oregon	0.57	0.08	05(01,37)	24.20 ^c
Idaho	0.49	0.11	18(01,42)	6.31	Pennsylvania	0.45	0.04	28(06,39)	-3.08
Illinois	0.42	0.04	33(09,41)	-9.39	Rhode Island	0.72	0.17	01(01,41)	56.20 ^c
Indiana	0.47	0.06	22(03,40)	2.56	South Carolina	0.46	0.07	25(02,41)	-0.08
Iowa	0.41	0.07	34(03,42)	-10.24	South Dakota	-	-	-(-,-)	-
Kansas	0.44	0.08	29(02,42)	-3.63	Tennessee	0.51	0.06	10(02,38)	11.21
Kentucky	0.34	0.06	40(13,42)	-27.31 ^c	Texas	0.51	0.03	12(03,31)	10.02
Louisiana	0.62	0.08	04(01,32)	35.19 ^c	Utah	0.30	0.08	42(08,42)	-34.27 ^c
Maine	0.52	0.13	06(01,42)	12.57	Vermont	-	-	-(-,-)	-
Maryland	0.38	0.05	38(07,42)	-17.55 ^c	Virginia	0.52	0.05	09(02,37)	11.81
Massachusetts	0.46	0.05	23(03,40)	0.53	Washington	0.40	0.05	35(06,42)	-12.31
Michigan	0.51	0.05	13(02,37)	9.72	West Virginia	0.49	0.10	19(01,42)	5.37
Minnesota	0.43	0.06	30(04,42)	-7.40	Wisconsin	0.49	0.06	15(02,39)	7.30
Mississippi	0.51	0.09	11(01,42)	10.80	Wyoming	-	-	-(-,-)	-
Missouri	0.39	0.05	36(07,42)	-16.35 ^c					



Age-Adjusted Death
Rates per 100,000
Quantile Interval



U.S. rate: 0.46

- ^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- ^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
- ^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
- ^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.
- SE Standard error of the rate.
- PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.