


Table 11.18
Cancer of the Kidney and Renal Pelvis (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	3.92	0.02		
High Five States				
Oklahoma	5.30 ^b	0.16	01(01,04)	35.23 ^c
Louisiana	5.14 ^b	0.15	02(01,07)	31.05 ^c
North Dakota	4.96	0.36	03(01,29)	26.46 ^c
Arkansas	4.65 ^b	0.17	04(02,22)	18.54 ^c
Kentucky	4.62 ^b	0.14	05(02,20)	17.77 ^c
Low Five States				
Wyoming	3.30	0.33	47(15,51)	-15.95 ^c
Massachusetts	3.29 ^b	0.09	48(37,50)	-16.05 ^c
Connecticut	3.27 ^b	0.13	49(35,50)	-16.63 ^c
D.C.	2.79 ^b	0.31	50(32,51)	-28.74 ^c
Hawaii	2.68 ^b	0.18	51(46,51)	-31.71 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	3.96	0.12	28(15,37)	0.94	Montana	3.43	0.24	43(20,51)	-12.54
Alaska	4.44	0.45	14(01,49)	13.29	Nebraska	4.43	0.21	15(02,32)	12.86
Arizona	3.75	0.10	33(22,43)	-4.48	Nevada	3.76	0.17	32(16,48)	-4.09
Arkansas	4.65 ^b	0.17	04(02,22)	18.54 ^c	New Hampshire	3.66	0.22	35(14,50)	-6.67
California	3.52 ^b	0.04	39(34,45)	-10.16	New Jersey	3.38 ^b	0.08	45(35,49)	-13.81
Colorado	3.47 ^b	0.12	42(30,49)	-11.65	New Mexico	4.48	0.20	13(02,30)	14.18
Connecticut	3.27 ^b	0.13	49(35,50)	-16.63 ^c	New York	3.30 ^b	0.06	46(39,50)	-15.76 ^c
Delaware	3.50	0.26	41(15,51)	-10.80	North Carolina	3.97	0.09	27(18,36)	1.19
D.C.	2.79 ^b	0.31	50(32,51)	-28.74 ^c	North Dakota	4.96	0.36	03(01,29)	26.46 ^c
Florida	3.42 ^b	0.05	44(36,48)	-12.79	Ohio	4.27 ^b	0.08	19(09,26)	8.83
Georgia	3.51 ^b	0.09	40(32,48)	-10.43	Oklahoma	5.30 ^b	0.16	01(01,04)	35.23 ^c
Hawaii	2.68 ^b	0.18	51(46,51)	-31.71 ^c	Oregon	3.94	0.13	29(15,40)	0.47
Idaho	4.07	0.23	24(05,44)	3.66	Pennsylvania	3.98	0.07	26(19,34)	1.51
Illinois	4.13	0.08	22(14,30)	5.20	Rhode Island	3.55	0.24	38(17,50)	-9.61
Indiana	4.56 ^b	0.12	09(03,20)	16.22 ^c	South Carolina	4.07	0.13	23(12,35)	3.73
Iowa	4.59 ^b	0.16	07(02,22)	17.06 ^c	South Dakota	4.41	0.31	16(01,40)	12.40
Kansas	4.57 ^b	0.17	08(02,24)	16.53 ^c	Tennessee	4.19	0.11	20(09,31)	6.79
Kentucky	4.62 ^b	0.14	05(02,20)	17.77 ^c	Texas	4.54 ^b	0.06	11(04,17)	15.78 ^c
Louisiana	5.14 ^b	0.15	02(01,07)	31.05 ^c	Utah	3.56	0.19	37(21,50)	-9.20
Maine	4.01	0.22	25(05,45)	2.32	Vermont	3.56	0.31	36(10,51)	-9.13
Maryland	3.70	0.11	34(24,45)	-5.64	Virginia	3.81	0.10	31(21,41)	-2.98
Massachusetts	3.29 ^b	0.09	48(37,50)	-16.05 ^c	Washington	3.92	0.11	30(18,38)	-0.13
Michigan	4.13	0.09	21(13,31)	5.37	West Virginia	4.62	0.20	06(01,26)	17.71 ^c
Minnesota	4.27	0.12	18(06,29)	8.87	Wisconsin	4.34 ^b	0.12	17(04,26)	10.59
Mississippi	4.55 ^b	0.17	10(02,25)	16.05 ^c	Wyoming	3.30	0.33	47(15,51)	-15.95 ^c
Missouri	4.51 ^b	0.12	12(03,21)	14.92					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.