

Table 14.9
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males and Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Survival Time																					
1-year	13.8	14.8	17.6	21.5	23.0	24.6	27.8	26.8	27.8	29.0	31.1	32.6	36.0	38.2	39.2	39.4	42.3	44.9	44.4	48.0	46.5
2-year	6.6	8.2	9.8	12.4	13.8	14.9	17.3	17.1	17.7	19.8	21.5	22.1	24.7	26.1	27.0	28.8	30.3	32.6	30.9	35.6	
3-year	4.8	6.0	7.3	8.8	9.4	10.8	12.5	13.0	14.3	15.4	15.9	17.3	20.5	20.5	21.4	23.4	24.0	25.5	23.9		
4-year	4.1	4.6	6.4	6.9	6.9	9.1	10.1	10.7	12.0	12.3	13.6	15.4	16.9	17.5	18.4	19.9	21.0	21.5			
5-year	3.4	3.8	5.4	5.6	5.7	8.1	8.1	9.4	10.1	11.6	11.8	14.0	15.5	14.5	16.8	17.0	18.5				
6-year	3.0	3.5	5.0	5.0	4.9	6.9	7.4	8.4	9.0	10.7	10.6	11.9	14.8	13.3	15.3	15.4					
7-year	2.8	3.1	4.8	4.6	4.4	6.5	6.6	6.9	8.3	9.9	9.5	11.3	13.0	12.6	14.0						
8-year	2.7	2.9	4.4	4.4	4.2	6.4	5.8	6.6	7.8	9.6	8.6	10.3	12.0	12.0							
9-year	2.6	2.8	4.2	4.3	4.1	5.9	5.6	5.6	7.5	9.3	8.2	10.0	11.4								
10-year	2.6	2.8	4.0	4.2	3.6	5.6	5.4	5.5	7.4	8.9	7.6	9.2									
11-year	2.6	2.6	4.0	3.9	3.2	5.5	5.4	4.8	6.9	8.6	7.5										
12-year	2.5	2.6	3.8	3.9	3.1	5.2	5.4	4.6	6.8	8.5											
13-year	2.3	2.4	3.8	3.7	3.1	5.0	5.4	4.5	6.6												
14-year	2.3	2.4	3.7	3.7	3.1	4.7	5.4	4.5													
15-year	2.2	2.4	3.7	3.7	3.0	3.9	5.0														
16-year	2.2	2.3	3.7	3.6	3.0	3.8															
17-year	2.1	2.3	3.7	3.5	2.9																
18-year	2.0	2.2	3.7	3.4																	
19-year	2.0	2.1	3.7																		
20-year	2.0	2.1	3.7																		

^a

Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).