

Table 14.10
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Survival Time																					
1-year	11.4	12.7	14.6	20.2	21.1	24.6	26.8	25.6	28.9	28.3	30.6	31.7	35.0	37.3	40.0	39.7	41.3	45.4	45.2	48.9	46.4
2-year	4.8	6.4	7.4	11.4	12.3	15.8	16.5	16.3	17.6	18.9	20.2	21.1	24.7	25.5	27.8	29.8	29.7	32.9	31.5	36.5	
3-year	3.2	4.9	5.2	8.1	8.9	11.2	12.0	12.0	13.7	15.4	15.2	17.6	20.1	19.9	21.8	23.8	24.1	25.9	24.1		
4-year	2.7	3.4	4.5	6.3	6.2	9.2	9.6	10.6	11.4	12.1	12.6	15.6	16.7	17.4	18.6	20.0	20.3	21.9			
5-year	2.0	2.8	3.6	5.2	5.0	7.8	7.9	8.9	10.1	11.2	10.8	14.2	15.6	14.3	17.0	16.9	18.0				
6-year	1.7	2.6	3.3	4.7	4.2	6.7	7.3	7.6	8.9	10.4	9.7	11.9	14.8	13.0	15.5	15.4					
7-year	1.4	2.2	3.3	4.3	3.7	6.5	6.7	6.2	8.3	9.8	8.9	11.1	13.2	12.2	13.9						
8-year	1.4	1.9	3.0	4.1	3.5	6.4	5.5	6.1	7.7	9.7	7.7	9.9	11.9	11.7							
9-year	1.3	1.9	3.0	3.9	3.4	5.9	5.4	5.2	7.5	9.7	7.2	9.4	11.2								
10-year	1.3	1.9	2.9	3.8	2.8	5.4	5.4	5.2	7.4	9.3	6.7	8.6									
11-year	1.3	1.8	2.9	3.7	2.3	5.2	5.4	4.4	6.9	9.1	6.7										
12-year	1.3	1.7	2.7	3.6	2.1	4.9	5.4	4.3	6.9	8.9											
13-year	1.1	1.4	2.7	3.4	2.1	4.7	5.4	4.3	6.8												
14-year	1.1	1.4	2.6	3.4	2.1	4.4	5.3	4.2													
15-year	1.0	1.4	2.6	3.4	2.1	3.7	4.9														
16-year	1.0	1.4	2.6	3.4	2.1	3.7															
17-year	1.0	1.4	2.6	3.4	2.1																
18-year	1.0	1.4	2.6	3.3																	
19-year	1.0	1.4	2.6																		
20-year	1.0	1.4	2.6																		

^a

Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).