

Table 14.11
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Survival Time																					
1-year	18.5	18.9	23.4	24.2	26.6	24.6	29.7	29.4	25.6	30.2	32.3	34.5	38.7	40.5	37.3	38.7	45.3	43.6	42.2	45.3	46.6
2-year	10.0	11.8	14.5	14.3	16.5	13.2	19.0	18.7	17.9	21.5	24.4	24.4	24.7	27.8	25.2	26.3	32.2	31.7	29.1	33.1	
3-year	7.8	8.2	11.4	10.2	10.4	9.9	13.6	15.2	15.3	15.3	17.7	16.6	21.4	22.4	20.3	22.3	23.7	24.7	23.3		
4-year	6.6	6.7	9.9	8.1	8.3	8.7	11.0	10.8	13.3	12.6	15.9	14.9	17.5	17.9	17.9	19.6	22.8	20.7			
5-year	6.1	5.6	8.8	6.4	7.0	8.5	8.4	10.2	10.3	12.3	14.0	13.3	15.0	15.1	16.4	17.4	19.8				
6-year	5.4	5.1	8.4	5.6	6.0	7.3	7.3	9.9	9.3	11.3	12.6	11.8	14.6	14.0	15.0	15.6					
7-year	5.4	4.7	7.7	5.1	5.7	6.7	6.2	8.4	8.2	10.0	11.0	11.6	12.7	13.5	14.1						
8-year	5.2	4.6	7.0	5.1	5.4	6.4	6.2	7.5	7.9	9.2	10.7	11.1	12.1	12.7							
9-year	4.9	4.5	6.6	5.0	5.4	5.9	5.8	6.2	7.4	8.4	10.6	11.1	11.5								
10-year	4.9	4.3	6.2	5.0	5.1	5.9	5.2	5.9	7.4	8.1	9.4	10.8									
11-year	4.9	4.3	6.0	4.5	4.8	5.9	5.2	5.6	6.9	7.6	9.2										
12-year	4.7	4.3	6.0	4.4	4.8	5.7	5.2	5.3	6.3	7.6											
13-year	4.4	4.2	5.8	4.4	4.8	5.4	5.2	5.0	5.9												
14-year	4.4	4.2	5.8	4.2	4.8	5.4	5.2	5.0													
15-year	4.4	4.2	5.8	4.0	4.4	4.3	5.0														
16-year	4.3	4.1	5.8	4.0	4.4	3.9															
17-year	4.0	3.9	5.8	3.8	4.0																
18-year	3.8	3.6	5.7	3.6																	
19-year	3.8	3.3	5.7																		
20-year	3.8	3.3	5.6																		

^a

Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).