


Table 14.18
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)
Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012


Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	5.96	0.02		
High Five States				
D.C.	9.17 ^b	0.56	01(01,04)	53.83 ^c
Texas	7.71 ^b	0.08	02(01,04)	29.35 ^c
Louisiana	7.69 ^b	0.18	03(01,06)	29.00 ^c
Hawaii	7.58 ^b	0.31	04(01,08)	27.15 ^c
California	7.24 ^b	0.06	05(03,07)	21.58 ^c
Low Five States				
Nebraska	4.55 ^b	0.21	47(36,51)	-23.58 ^c
Idaho	4.33 ^b	0.23	48(38,51)	-27.32 ^c
Utah	4.19 ^b	0.20	49(42,51)	-29.63 ^c
Montana	4.17 ^b	0.26	50(40,51)	-30.04 ^c
North Dakota	3.78 ^b	0.31	51(44,51)	-36.61 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	6.14	0.15	14(07,22)	3.08	Montana	4.17 ^b	0.26	50(40,51)	-30.04 ^c
Alaska	5.69	0.47	22(06,48)	-4.57	Nebraska	4.55 ^b	0.21	47(36,51)	-23.58 ^c
Arizona	5.72	0.13	21(13,34)	-4.00	Nevada	5.48	0.20	29(14,42)	-8.04
Arkansas	6.34	0.19	09(06,21)	6.45	New Hampshire	4.63 ^b	0.25	45(32,51)	-22.29 ^c
California	7.24 ^b	0.06	05(03,07)	21.58 ^c	New Jersey	5.50 ^b	0.11	27(19,37)	-7.66
Colorado	5.16 ^b	0.15	37(24,44)	-13.37	New Mexico	6.68	0.25	07(04,17)	12.14
Connecticut	5.45	0.16	33(16,41)	-8.53	New York	5.93	0.07	17(12,23)	-0.50
Delaware	6.28	0.35	10(04,34)	5.47	North Carolina	5.47 ^b	0.10	32(19,38)	-8.27
D.C.	9.17 ^b	0.56	01(01,04)	53.83 ^c	North Dakota	3.78 ^b	0.31	51(44,51)	-36.61 ^c
Florida	5.66 ^b	0.07	23(17,31)	-5.04	Ohio	5.27 ^b	0.09	35(26,41)	-11.50
Georgia	5.47 ^b	0.11	31(19,38)	-8.19	Oklahoma	5.78	0.17	20(10,34)	-2.95
Hawaii	7.58 ^b	0.31	04(01,08)	27.15 ^c	Oregon	6.05	0.16	16(08,27)	1.48
Idaho	4.33 ^b	0.23	48(38,51)	-27.32 ^c	Pennsylvania	5.53 ^b	0.08	25(19,35)	-7.26
Illinois	5.65	0.09	24(16,33)	-5.16	Rhode Island	6.18	0.32	13(06,35)	3.67
Indiana	5.10 ^b	0.12	38(29,44)	-14.39	South Carolina	5.81	0.15	19(11,33)	-2.52
Iowa	4.71 ^b	0.16	44(36,49)	-21.03 ^c	South Dakota	4.62 ^b	0.31	46(25,51)	-22.49 ^c
Kansas	4.81 ^b	0.17	43(33,49)	-19.34 ^c	Tennessee	6.19	0.13	12(07,20)	3.97
Kentucky	5.52	0.15	26(16,39)	-7.38	Texas	7.71 ^b	0.08	02(01,04)	29.35 ^c
Louisiana	7.69 ^b	0.18	03(01,06)	29.00 ^c	Utah	4.19 ^b	0.20	49(42,51)	-29.63 ^c
Maine	4.86 ^b	0.24	41(26,50)	-18.49 ^c	Vermont	5.32	0.38	34(08,49)	-10.68
Maryland	6.06	0.14	15(08,25)	1.69	Virginia	5.49 ^b	0.12	28(19,38)	-7.84
Massachusetts	6.24	0.13	11(07,19)	4.65	Washington	6.40	0.14	08(06,16)	7.40
Michigan	5.47 ^b	0.10	30(20,37)	-8.19	West Virginia	4.84 ^b	0.20	42(29,49)	-18.80 ^c
Minnesota	5.02 ^b	0.13	39(31,46)	-15.79 ^c	Wisconsin	5.17 ^b	0.13	36(26,44)	-13.22
Mississippi	7.04 ^b	0.21	06(03,09)	18.22 ^c	Wyoming	4.94	0.41	40(12,51)	-17.05 ^c
Missouri	5.82	0.13	18(11,31)	-2.26					



Age-Adjusted Death
Rates per 100,000
Quantile Interval



- ^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- ^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
- ^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
- ^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.
- SE Standard error of the rate.
- PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.