


Table 14.19
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)
Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012


Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	8.83	0.03		
High Five States				
D.C.	14.71 ^b	1.08	01(01,02)	66.48 ^c
Louisiana	11.71 ^b	0.34	02(01,05)	32.54 ^c
Texas	11.29 ^b	0.15	03(02,05)	27.87 ^c
Hawaii	11.06 ^b	0.55	04(02,12)	25.26 ^c
California	10.55 ^b	0.11	05(04,08)	19.45 ^c
Low Five States				
Idaho	6.37 ^b	0.41	47(36,51)	-27.92 ^c
South Dakota	5.85 ^b	0.52	48(36,51)	-33.75 ^c
Montana	5.62 ^b	0.44	49(41,51)	-36.35 ^c
Utah	5.53 ^b	0.33	50(45,51)	-37.40 ^c
North Dakota	5.40 ^b	0.55	51(40,51)	-38.82 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	9.06	0.28	15(07,27)	2.61	Montana	5.62 ^b	0.44	49(41,51)	-36.35 ^c
Alaska	8.15	0.84	31(04,49)	-7.70	Nebraska	6.62 ^b	0.38	46(34,50)	-25.05 ^c
Arizona	8.12	0.22	32(18,38)	-8.09	Nevada	7.83	0.36	35(16,44)	-11.36
Arkansas	9.48	0.36	11(05,24)	7.35	New Hampshire	6.96 ^b	0.45	44(22,49)	-21.15 ^c
California	10.55 ^b	0.11	05(04,08)	19.45 ^c	New Jersey	8.27	0.20	26(17,36)	-6.33
Colorado	7.61 ^b	0.27	36(23,44)	-13.82	New Mexico	9.67	0.43	10(04,27)	9.45
Connecticut	8.52	0.31	21(11,37)	-3.57	New York	8.90	0.14	17(11,23)	0.80
Delaware	9.82	0.65	08(02,34)	11.13	North Carolina	8.28	0.19	24(17,36)	-6.28
D.C.	14.71 ^b	1.08	01(01,02)	66.48 ^c	North Dakota	5.40 ^b	0.55	51(40,51)	-38.82 ^c
Florida	8.54	0.12	20(16,30)	-3.36	Ohio	7.98 ^b	0.16	34(22,38)	-9.64
Georgia	8.24	0.21	27(17,37)	-6.70	Oklahoma	8.22	0.30	29(14,39)	-6.97
Hawaii	11.06 ^b	0.55	04(02,12)	25.26 ^c	Oregon	9.00	0.30	16(08,30)	1.85
Idaho	6.37 ^b	0.41	47(36,51)	-27.92 ^c	Pennsylvania	8.32	0.15	23(18,35)	-5.78
Illinois	8.28	0.17	25(17,36)	-6.28	Rhode Island	9.72	0.60	09(02,32)	10.08
Indiana	7.32 ^b	0.22	38(32,45)	-17.12 ^c	South Carolina	8.84	0.28	18(09,32)	0.06
Iowa	6.72 ^b	0.29	45(36,49)	-23.94 ^c	South Dakota	5.85 ^b	0.52	48(36,51)	-33.75 ^c
Kansas	6.98 ^b	0.31	42(32,48)	-21.03 ^c	Tennessee	9.46	0.25	12(06,20)	7.14
Kentucky	8.21	0.28	30(15,38)	-7.06	Texas	11.29 ^b	0.15	03(02,05)	27.87 ^c
Louisiana	11.71 ^b	0.34	02(01,05)	32.54 ^c	Utah	5.53 ^b	0.33	50(45,51)	-37.40 ^c
Maine	7.16 ^b	0.44	40(21,49)	-18.99 ^c	Vermont	8.22	0.70	28(06,47)	-6.90
Maryland	9.29	0.27	14(07,23)	5.20	Virginia	8.36	0.22	22(15,36)	-5.37
Massachusetts	9.85 ^b	0.25	07(05,15)	11.53	Washington	9.34	0.24	13(07,21)	5.71
Michigan	7.99 ^b	0.18	33(21,38)	-9.51	West Virginia	6.99 ^b	0.37	41(29,49)	-20.92 ^c
Minnesota	7.19 ^b	0.24	39(32,47)	-18.62 ^c	Wisconsin	7.45 ^b	0.23	37(28,45)	-15.67 ^c
Mississippi	10.43 ^b	0.40	06(02,14)	18.09 ^c	Wyoming	6.97	0.71	43(13,51)	-21.10 ^c
Missouri	8.69	0.24	19(11,32)	-1.61					



Age-Adjusted Death
Rates per 100,000
Quantile Interval



- ^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- ^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
- ^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
- ^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.
- SE Standard error of the rate.
- PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.