



Table 14.20
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)
Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Females									
State	Rate	SE	Rank(CI) ^d	PD					
TOTAL U.S.	3.53	0.02							
High Five States									
D.C.	4.81	0.54	01(01,29)	36.01 ^c					
Texas	4.60 ^b	0.09	02(01,06)	30.31 ^c					
Hawaii	4.41	0.32	03(01,17)	24.80 ^c					
California	4.40 ^b	0.07	04(01,07)	24.68 ^c					
Louisiana	4.40 ^b	0.19	05(01,09)	24.60 ^c					
Low Five States									
Vermont	2.80	0.38	47(07,51)	-20.79 ^c					
Montana	2.78	0.30	48(10,51)	-21.22 ^c					
New Hampshire	2.55 ^b	0.25	49(24,51)	-27.77 ^c					
Idaho	2.52 ^b	0.24	50(27,51)	-28.65 ^c					
North Dakota	2.32 ^b	0.32	51(22,51)	-34.42 ^c					

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	3.77	0.16	09(05,27)	6.75	Montana	2.78	0.30	48(10,51)	-21.22 ^c
Alaska	3.34	0.52	25(01,51)	-5.59	Nebraska	2.84	0.23	46(15,51)	-19.68 ^c
Arizona	3.58	0.14	12(07,33)	1.30	Nevada	3.35	0.22	23(07,48)	-5.18
Arkansas	3.67	0.20	11(05,38)	3.96	New Hampshire	2.55 ^b	0.25	49(24,51)	-27.77 ^c
California	4.40 ^b	0.07	04(01,07)	24.68 ^c	New Jersey	3.30	0.11	27(11,43)	-6.56
Colorado	3.01	0.15	40(16,50)	-14.69	New Mexico	4.02	0.26	07(01,28)	13.90
Connecticut	2.96 ^b	0.16	43(18,50)	-16.18 ^c	New York	3.51	0.08	14(09,28)	-0.62
Delaware	3.35	0.34	22(03,51)	-5.16	North Carolina	3.15 ^b	0.11	35(17,46)	-10.85
D.C.	4.81	0.54	01(01,29)	36.01 ^c	North Dakota	2.32 ^b	0.32	51(22,51)	-34.42 ^c
Florida	3.17 ^b	0.07	33(21,43)	-10.18	Ohio	3.06 ^b	0.09	38(22,47)	-13.31
Georgia	3.25	0.11	29(13,45)	-8.06	Oklahoma	3.68	0.18	10(06,34)	4.24
Hawaii	4.41	0.32	03(01,17)	24.80 ^c	Oregon	3.41	0.17	19(07,44)	-3.53
Idaho	2.52 ^b	0.24	50(27,51)	-28.65 ^c	Pennsylvania	3.20 ^b	0.09	32(18,44)	-9.51
Illinois	3.52	0.10	13(08,31)	-0.34	Rhode Island	3.43	0.32	18(04,49)	-2.87
Indiana	3.27	0.13	28(11,45)	-7.36	South Carolina	3.31	0.15	26(09,45)	-6.45
Iowa	2.95	0.17	44(17,50)	-16.46 ^c	South Dakota	3.46	0.37	16(02,50)	-2.13
Kansas	2.91	0.19	45(16,51)	-17.55 ^c	Tennessee	3.51	0.13	15(08,37)	-0.64
Kentucky	3.24	0.16	30(10,47)	-8.39	Texas	4.60 ^b	0.09	02(01,06)	30.31 ^c
Louisiana	4.40 ^b	0.19	05(01,09)	24.60 ^c	Utah	2.97	0.23	42(11,51)	-15.93 ^c
Maine	3.01	0.26	41(09,51)	-14.78	Vermont	2.80	0.38	47(07,51)	-20.79 ^c
Maryland	3.45	0.14	17(08,40)	-2.30	Virginia	3.14	0.12	36(16,47)	-11.14
Massachusetts	3.35	0.13	24(10,43)	-5.24	Washington	3.79	0.14	08(05,23)	7.33
Michigan	3.36	0.10	21(11,40)	-5.02	West Virginia	3.06	0.22	39(10,50)	-13.43
Minnesota	3.13	0.14	37(14,48)	-11.42	Wisconsin	3.23	0.14	31(12,46)	-8.55
Mississippi	4.30	0.22	06(01,13)	21.66 ^c	Wyoming	3.16	0.46	34(02,51)	-10.65
Missouri	3.38	0.13	20(09,42)	-4.25					



Age-Adjusted Death
Rates per 100,000
Quantile Interval



U.S. rate: 3.53

- ^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- ^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
- ^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
- ^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.
- SE Standard error of the rate.
- PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.