

Table 15.23
Cancer of the Lung And Bronchus (Invasive)

Age-Adjusted SEER Death Rates^a
By Registry, Race and Sex

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>SEER Death Rates^a, 2008-2012</u>									
California	36.15	43.67	30.51	37.39	43.95	32.43	46.82	60.70	37.08
Greater Bay Area	32.83	38.93	28.35	33.72	37.71	30.80	50.69	63.70	41.51
San Francisco-Oakland	34.40	40.74	29.74	34.90	38.36	32.39	52.81	67.86	42.45
San Jose-Monterey	29.77	35.44	25.61	31.67	36.61	28.01	36.13	36.70	35.22
Los Angeles	32.42	41.34	25.84	32.12	39.53	26.51	47.09	63.07	36.87
Greater California	39.00	46.35	33.43	40.42	47.35	35.17	44.27	56.10	34.94
Connecticut	41.58	49.73	35.82	42.28	49.93	36.88	38.55	52.76	29.42
Detroit	52.74	65.98	43.36	51.99	63.19	44.08	58.90	81.69	43.95
Georgia	50.06	68.00	37.23	52.44	68.47	40.55	44.94	70.36	29.36
Atlanta	40.13	52.10	32.06	40.30	49.18	34.09	42.31	63.90	30.14
Rural Georgia	52.35	82.06	30.33	52.19	73.20	35.93	53.30	106.74	20.40
Greater Georgia	54.33	74.22	39.81	56.62	74.89	42.93	46.69	73.47	29.43
Hawaii	33.46	44.37	24.98	38.86	44.31	33.73	44.15	61.29	-
Iowa	47.75	61.40	37.49	47.78	61.47	37.50	59.86	70.37	51.66
Kentucky	70.92	92.19	55.23	71.21	92.30	55.55	74.45	101.13	56.67
Louisiana	57.64	77.40	42.84	56.66	72.97	44.29	61.55	93.44	39.73
New Jersey	42.14	52.05	35.26	43.55	52.85	37.10	44.00	61.82	32.95
New Mexico	33.27	40.90	27.21	34.91	42.46	28.89	42.47	66.34	20.62
Seattle-Puget Sound	44.36	52.31	38.50	45.65	53.03	40.19	47.52	59.65	37.81
Utah	20.41	26.35	15.56	20.46	26.37	15.63	32.99	-	-
9 SEER Areas ^b	40.49	49.86	33.57	41.27	49.51	35.15	49.44	68.44	37.01
11 SEER Areas ^b	38.10	47.17	31.39	38.90	46.80	32.99	48.67	66.61	36.91
13 SEER Areas ^b	38.31	47.45	31.53	39.06	46.98	33.13	48.71	67.12	36.65
18 SEER Areas ^b	42.90	53.62	34.92	44.08	53.80	36.78	50.02	71.04	36.00
Total U.S.	47.22	59.79	37.80	47.94	59.68	39.05	50.65	73.09	35.83

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta. The SEER 11 areas comprise the SEER 9 areas plus San Jose-Monterey and Los Angeles.

The SEER 13 areas comprise the SEER 11 areas plus the Alaska Native Registry and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG.

- Statistic not shown. Rate based on less than 16 cases for the time interval.