


Table 15.24
Cancer of the Lung and Bronchus (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	47.22	0.05		
High Five States				
Kentucky	70.92 ^b	0.55	01(01,01)	50.18 ^c
Arkansas	63.26 ^b	0.62	02(02,04)	33.95 ^c
West Virginia	62.70 ^b	0.73	03(02,05)	32.77 ^c
Tennessee	61.39 ^b	0.42	04(03,05)	30.00 ^c
Mississippi	61.32 ^b	0.63	05(02,05)	29.86 ^c
Low Five States				
California	36.15 ^b	0.14	47(47,47)	-23.44 ^c
Colorado	34.45 ^b	0.39	48(48,50)	-27.05 ^c
Hawaii	33.46 ^b	0.66	49(48,50)	-29.14 ^c
New Mexico	33.27 ^b	0.55	50(48,50)	-29.54 ^c
Utah	20.41 ^b	0.45	51(51,51)	-56.79 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	58.14 ^b	0.47	07(06,10)	23.12 ^c	Montana	42.77 ^b	0.86	37(34,44)	-9.42 ^c
Alaska	52.07	1.57	17(11,26)	10.25 ^c	Nebraska	44.18 ^b	0.67	35(32,39)	-6.45 ^c
Arizona	38.47 ^b	0.33	46(44,46)	-18.53 ^c	Nevada	49.37	0.63	22(17,27)	4.54
Arkansas	63.26 ^b	0.62	02(02,04)	33.95 ^c	New Hampshire	48.65	0.82	24(17,31)	3.02
California	36.15 ^b	0.14	47(47,47)	-23.44 ^c	New Jersey	42.14 ^b	0.30	39(37,43)	-10.77 ^c
Colorado	34.45 ^b	0.39	48(48,50)	-27.05 ^c	New Mexico	33.27 ^b	0.55	50(48,50)	-29.54 ^c
Connecticut	41.58 ^b	0.45	41(37,44)	-11.95 ^c	New York	42.03 ^b	0.20	40(37,43)	-11.00 ^c
Delaware	54.20 ^b	1.03	12(10,17)	14.78 ^c	North Carolina	53.03 ^b	0.33	15(12,17)	12.30 ^c
D.C.	42.51 ^b	1.23	38(32,45)	-9.98 ^c	North Dakota	40.84 ^b	1.03	43(36,46)	-13.52 ^c
Florida	46.58	0.19	30(28,33)	-1.36	Ohio	55.32 ^b	0.29	11(11,13)	17.14 ^c
Georgia	50.06 ^b	0.34	18(17,23)	6.01 ^c	Oklahoma	58.42 ^b	0.54	06(06,10)	23.71 ^c
Hawaii	33.46 ^b	0.66	49(48,50)	-29.14 ^c	Oregon	47.23	0.47	28(24,32)	0.02
Idaho	39.15 ^b	0.71	45(42,46)	-17.10 ^c	Pennsylvania	48.63 ^b	0.25	25(21,27)	2.98
Illinois	49.74 ^b	0.28	20(17,24)	5.32 ^c	Rhode Island	49.99	0.91	19(16,27)	5.86 ^c
Indiana	57.75 ^b	0.41	08(06,10)	22.30 ^c	South Carolina	53.32 ^b	0.46	14(12,17)	12.91 ^c
Iowa	47.75	0.52	27(22,31)	1.11	South Dakota	44.97	0.98	34(27,39)	-4.77
Kansas	48.92	0.57	23(18,28)	3.59	Tennessee	61.39 ^b	0.42	04(03,05)	30.00 ^c
Kentucky	70.92 ^b	0.55	01(01,01)	50.18 ^c	Texas	43.53 ^b	0.20	36(34,38)	-7.83 ^c
Louisiana	57.64 ^b	0.50	09(06,10)	22.06 ^c	Utah	20.41 ^b	0.45	51(51,51)	-56.79 ^c
Maine	53.77 ^b	0.80	13(11,17)	13.87 ^c	Vermont	49.53	1.16	21(15,31)	4.88
Maryland	46.41	0.40	31(27,34)	-1.72	Virginia	48.55 ^b	0.35	26(20,28)	2.81
Massachusetts	46.80	0.36	29(26,33)	-0.90	Washington	45.77 ^b	0.37	33(29,35)	-3.08
Michigan	52.17 ^b	0.31	16(14,18)	10.47 ^c	West Virginia	62.70 ^b	0.73	03(02,05)	32.77 ^c
Minnesota	41.56 ^b	0.39	42(37,44)	-11.98 ^c	Wisconsin	45.83 ^b	0.38	32(29,35)	-2.95
Mississippi	61.32 ^b	0.63	05(02,05)	29.86 ^c	Wyoming	39.62 ^b	1.19	44(37,46)	-16.10 ^c
Missouri	56.95 ^b	0.41	10(07,11)	20.60 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 5% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.