


Table 15.25
Cancer of the Lung and Bronchus (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	59.79	0.09		
High Five States				
Kentucky	92.19 ^b	0.96	01(01,02)	54.19 ^c
Mississippi	88.45 ^b	1.18	02(01,03)	47.94 ^c
Arkansas	88.00 ^b	1.10	03(01,03)	47.18 ^c
Tennessee	83.57 ^b	0.76	04(04,06)	39.77 ^c
Alabama	82.56 ^b	0.86	05(04,06)	38.09 ^c
Low Five States				
Hawaii	44.37 ^b	1.14	47(44,49)	-25.78 ^c
California	43.67 ^b	0.24	48(46,48)	-26.96 ^c
New Mexico	40.90 ^b	0.93	49(48,50)	-31.59 ^c
Colorado	40.75 ^b	0.65	50(48,50)	-31.84 ^c
Utah	26.35 ^b	0.76	51(51,51)	-55.93 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	82.56 ^b	0.86	05(04,06)	38.09 ^c	Montana	49.66 ^b	1.37	42(37,46)	-16.94 ^c
Alaska	61.25	2.59	24(14,37)	2.44	Nebraska	55.46 ^b	1.13	34(25,38)	-7.24 ^c
Arizona	46.63 ^b	0.55	44(43,47)	-22.00 ^c	Nevada	55.94 ^b	0.99	33(25,38)	-6.44 ^c
Arkansas	88.00 ^b	1.10	03(01,03)	47.18 ^c	New Hampshire	57.71	1.35	27(23,37)	-3.47
California	43.67 ^b	0.24	48(46,48)	-26.96 ^c	New Jersey	52.05 ^b	0.51	39(37,41)	-12.95 ^c
Colorado	40.75 ^b	0.65	50(48,50)	-31.84 ^c	New Mexico	40.90 ^b	0.93	49(48,50)	-31.59 ^c
Connecticut	49.73 ^b	0.76	41(39,44)	-16.83 ^c	New York	51.94 ^b	0.34	40(37,41)	-13.13 ^c
Delaware	66.37 ^b	1.73	16(13,23)	11.01 ^c	North Carolina	72.03 ^b	0.59	12(10,13)	20.48 ^c
D.C.	54.60	2.19	36(23,43)	-8.68 ^c	North Dakota	53.31 ^b	1.76	38(25,43)	-10.84 ^c
Florida	58.11 ^b	0.32	26(24,30)	-2.82	Ohio	70.92 ^b	0.50	13(11,14)	18.62 ^c
Georgia	68.00 ^b	0.63	14(14,16)	13.73 ^c	Oklahoma	75.70 ^b	0.93	08(07,10)	26.61 ^c
Hawaii	44.37 ^b	1.14	47(44,49)	-25.78 ^c	Oregon	55.15 ^b	0.76	35(27,38)	-7.75 ^c
Idaho	45.86 ^b	1.14	46(43,48)	-23.30 ^c	Pennsylvania	62.44 ^b	0.43	20(17,24)	4.44
Illinois	62.40 ^b	0.47	21(17,24)	4.37	Rhode Island	62.52	1.56	18(15,26)	4.56
Indiana	75.33 ^b	0.72	09(07,10)	25.99 ^c	South Carolina	72.72 ^b	0.83	11(09,13)	21.62 ^c
Iowa	61.40	0.88	23(18,26)	2.69	South Dakota	58.16	1.67	25(18,37)	-2.72
Kansas	61.68	0.96	22(17,26)	3.16	Tennessee	83.57 ^b	0.76	04(04,06)	39.77 ^c
Kentucky	92.19 ^b	0.96	01(01,02)	54.19 ^c	Texas	56.44 ^b	0.35	30(26,35)	-5.60 ^c
Louisiana	77.40 ^b	0.90	07(06,09)	29.45 ^c	Utah	26.35 ^b	0.76	51(51,51)	-55.93 ^c
Maine	66.50 ^b	1.34	15(14,19)	11.23 ^c	Vermont	57.23	1.89	28(19,39)	-4.27
Maryland	57.07 ^b	0.69	29(24,35)	-4.54	Virginia	62.48 ^b	0.61	19(17,24)	4.51
Massachusetts	56.18 ^b	0.60	32(26,37)	-6.03 ^c	Washington	53.85 ^b	0.61	37(33,39)	-9.93 ^c
Michigan	64.69 ^b	0.52	17(15,19)	8.20 ^c	West Virginia	80.46 ^b	1.26	06(04,07)	34.57 ^c
Minnesota	49.66 ^b	0.64	43(40,44)	-16.94 ^c	Wisconsin	56.30 ^b	0.64	31(26,36)	-5.83 ^c
Mississippi	88.45 ^b	1.18	02(01,03)	47.94 ^c	Wyoming	46.47 ^b	1.91	45(39,49)	-22.28 ^c
Missouri	72.76 ^b	0.71	10(09,13)	21.70 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 5% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.