Table 18.3 Myeloma

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program, 1975-2012 With up to Five Joinpoints, 1992-2012 With up to Three Joinpoints, Females by Race/Ethnicity

| | JP Tre | JP Trend 1 | | JP Trend 2 | | JP Trend 3 | | JP Trend 4 | | JP Trend 5 | | JP Trend 6 | | AAPC ^d | |
|--|---------|------------|---------|------------|---------|------------|---------|------------|-------|------------|-------|------------|---------|-------------------|--|
| | Years | APC | Years | APC | Years | APC | Years | APC | Years | APC | Years | APC | 2003-12 | 2008-12 | |
| SEER 9 Delay-Adjusted Incidence ^a , 1975-2012 | | | | | | | | | | | | | | | |
| All Races | 1975-12 | 0.8* | | | | | | | | | | | 0.8* | 0.8* | |
| White | 1975-12 | 0.6* | | | | | | | | | | | 0.6* | 0.6* | |
| Black | 1975-12 | 1.1* | | | | | | | | | | | 1.1* | 1.1* | |
| SEER 13 Delay-Adjusted Incidence ^b , 199 | | | 92-2012 | | | | | | | | | | | | |
| All Races | 1992-12 | 0.7* | | | | | | | | | | | 0.7* | 0.7* | |
| White | 1992-12 | 0.6* | | | | | | | | | | | 0.6* | 0.6* | |
| Black | 1992-12 | 0.7* | | | | | | | | | | | 0.7* | 0.7* | |
| SEER 9 Observed Incidence ^a , 1975-2012 | | | | | | | | | | | | | | | |
| All Races | 1975-12 | 0.6* | | | | | | | | | | | 0.6* | 0.6* | |
| White | 1975-12 | 0.5* | | | | | | | | | | | 0.5* | 0.5* | |
| Black | 1975-12 | 0.9* | | | | | | | | | | | 0.9* | 0.9* | |
| SEER 13 Observed | | | 2 | | | | | | | | | | | | |
| All Races | 1992-12 | 0.4* | | | | | | | | | | | 0.4* | 0.4* | |
| White | 1992-12 | 0.3 | | | | | | | | | | | 0.3 | 0.3 | |
| White NH ^{ef} | 1992-12 | 0.3 | | | | | | | | | | | 0.3 | 0.3 | |
| Black | 1992-12 | 0.4 | | | | | | | | | | | 0.4 | 0.4 | |
| Black NH ^{ef} | 1992-12 | 0.4 | | | | | | | | | | | 0.4 | 0.4 | |
| API ^e | 1992-12 | 0.6 | | | | | | | | | | | 0.6 | 0.6 | |
| AI/AN ^{eg} | 1992-12 | 0.0 | | | | | | | | | | | 0.0 | 0.0 | |
| Hispanic ^f | 1992-12 | -0.2 | | | | | | | | | | | -0.2 | -0.2 | |
| U.S. Cancer Mortality ^c , 1975-2012 | | | | | | | | | | | | | | | |
| All Races | 1975-93 | 1.5* | 1993-02 | -0.5 | 2002-09 | -2.7* | 2009-12 | 2.0 | | | | | -1.2* | 0.8 | |
| White | 1975-93 | 1.4* | 1993-02 | -0.5 | 2002-09 | -2.6* | 2009-12 | 1.7 | | | | | -1.2* | 0.6 | |
| Black | 1975-95 | 2.1* | 1995-12 | -1.7* | | | | | | | | | -1.7* | -1.7* | |
| U.S. Cancer Mortality ^c , 1992-2012 | | | | | | | | | | | | | | | |
| All Races | 1992-01 | -0.2 | 2001-09 | -2.5* | 2009-12 | 1.9 | | | | | | | -1.1* | 0.7 | |
| White | 1992-02 | -0.4 | 2002-09 | -2.7* | 2009-12 | 1.8 | | | | | | | -1.2* | 0.7 | |
| White NH ^{ef} | 1992-02 | -0.3 | 2002-08 | -3.0* | 2008-12 | 0.7 | | | | | | | -1.4* | 0.7 | |
| Black | 1992-00 | 0.1 | 2000-09 | -2.8* | 2009-12 | 2.4 | | | | | | | -1.1 | 1.1 | |
| Black NH ^{ef} | 1992-99 | 0.5 | 1999-09 | -2.6* | 2009-12 | 2.5 | | | | | | | -0.9 | 1.2 | |
| API ^e | 1992-12 | -0.8 | | | | | | | | | | | -0.8 | -0.8 | |
| AI/AN ^{eg} | 1992-12 | -2.8* | | | | | | | | | | | -2.8* | -2.8* | |
| Hispanic ^f | 1992-12 | -0.7 | | | | | | | | | | | -0.7 | -0.7 | |
| | | | | | | | | | | | | | | | |

Joinpoint Regression Program Version 4.2.0, April 2015, National Cancer Institute. (<u>http://surveillance.cancer.gov/joinpoint</u>/). The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

- Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).
- Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
- The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

- Data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.
- The APC/AAPC is significantly different from zero (p<.05).
- Joinpoint regression line analysis could not be performed on data series.

а

b

С

d

e

f

g

*