

Table 18.9
MyelomaSEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males and Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Survival Time																					
1-year	68.0	69.5	72.3	71.8	74.3	70.6	74.3	74.7	73.7	72.3	72.6	73.5	77.3	74.5	75.8	73.9	76.7	77.2	81.4	81.1	79.2
2-year	52.1	54.5	57.6	57.4	58.9	57.5	60.4	62.2	59.8	59.9	59.5	62.8	65.1	66.1	64.3	66.0	66.2	68.3	71.8	71.7	
3-year	39.6	42.9	45.4	46.3	47.9	44.9	48.7	51.1	47.8	49.9	49.4	53.3	55.8	58.1	57.1	57.7	57.3	60.8	64.0		
4-year	31.5	34.3	35.8	36.9	40.1	38.2	38.7	39.7	38.8	41.2	43.9	46.7	47.6	50.4	50.1	50.7	49.5	54.8			
5-year	25.2	26.9	27.4	29.9	33.5	31.8	31.3	34.0	31.1	34.2	37.2	39.6	41.8	44.8	45.8	45.5	45.1				
6-year	20.1	21.0	22.1	24.5	28.3	27.1	26.3	29.6	27.0	29.2	32.4	35.0	37.4	39.2	40.7	40.4					
7-year	16.2	16.8	17.4	20.4	23.7	23.7	22.5	24.2	23.2	25.3	28.9	31.4	32.8	34.7	35.8						
8-year	13.6	13.4	14.9	17.4	20.4	21.3	19.8	22.0	19.8	22.6	25.4	28.2	29.4	30.9							
9-year	11.5	11.2	11.9	15.4	18.3	19.5	17.1	19.3	18.0	20.5	22.8	25.2	26.6								
10-year	9.9	9.7	10.4	13.7	17.2	17.2	15.5	17.6	16.0	19.3	20.7	22.5									
11-year	8.8	8.7	9.2	12.7	15.6	15.6	14.2	16.4	13.9	18.3	19.4										
12-year	7.8	7.7	8.4	11.1	14.3	14.8	12.9	14.7	13.3	16.9											
13-year	6.7	6.6	7.6	10.3	13.5	13.3	11.7	13.7	12.2												
14-year	6.1	6.1	7.0	9.5	12.3	12.5	11.0	13.4													
15-year	5.3	5.7	6.2	9.2	11.9	11.4	10.8														
16-year	4.9	5.2	5.7	8.6	10.7	9.8															
17-year	4.7	4.7	5.4	8.0	9.9																
18-year	4.1	4.3	5.1	7.5																	
19-year	3.8	3.8	4.9																		
20-year	3.4	3.5	4.7																		

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).