

Table 18.10

Myeloma

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Survival Time																					
1-year	67.7	69.6	73.2	72.4	74.6	71.2	74.0	76.1	75.7	73.6	74.2	74.2	76.9	76.6	75.8	73.6	77.3	76.2	83.2	82.9	79.7
2-year	52.2	54.7	59.3	58.5	59.8	58.3	60.8	63.5	61.4	61.6	61.3	61.4	65.5	68.0	63.7	66.5	67.9	67.2	73.0	73.5	
3-year	38.9	42.8	46.3	46.6	49.2	45.0	50.4	53.5	50.6	52.5	51.1	52.4	56.4	60.0	56.3	58.0	59.6	58.4	65.4		
4-year	30.9	33.7	36.8	37.0	41.2	39.8	40.7	43.5	42.8	43.2	45.9	46.6	47.6	51.5	50.0	51.6	51.4	51.6			
5-year	24.4	25.8	28.4	30.3	36.2	33.7	31.4	37.9	34.5	35.4	39.6	41.2	42.2	45.6	46.4	47.1	47.1				
6-year	19.3	20.8	23.5	25.1	31.6	29.7	26.3	33.7	29.9	29.9	34.6	36.1	37.7	39.8	40.6	42.7					
7-year	15.5	16.9	19.0	21.3	26.0	25.8	22.3	27.8	25.8	26.1	30.9	32.7	34.3	35.4	35.1						
8-year	13.1	13.8	16.2	18.6	22.0	23.9	20.0	25.4	21.4	23.3	26.9	29.4	30.4	32.0							
9-year	11.1	11.7	12.9	16.7	20.0	21.2	18.1	23.1	19.1	19.7	24.3	26.3	28.0								
10-year	9.3	9.9	11.2	14.8	19.0	18.3	16.3	21.4	17.1	18.4	21.7	23.6									
11-year	7.6	8.9	9.9	13.6	17.3	17.0	14.8	20.5	14.0	17.0	20.5										
12-year	7.0	7.7	9.4	12.0	15.4	16.6	14.2	18.4	13.3	15.5											
13-year	5.8	6.7	8.6	11.2	14.8	14.4	13.0	16.9	12.5												
14-year	5.3	6.1	8.0	10.4	13.1	13.9	12.6	16.4													
15-year	4.7	5.7	7.0	10.0	12.9	12.6	12.0														
16-year	4.5	5.2	6.7	9.6	11.1	9.6															
17-year	4.0	4.8	6.3	8.8	10.3																
18-year	3.6	4.4	6.1	8.1																	
19-year	3.4	4.2	6.0																		
20-year	2.7	3.9	5.9																		

^a

Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).