

Table 18.11
MyelomaSEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Survival Time																					
1-year	68.4	69.4	71.4	71.0	74.0	69.9	74.6	73.0	71.4	70.8	70.8	72.7	77.8	72.1	75.8	74.3	75.9	78.4	79.3	78.9	78.6
2-year	52.0	54.3	55.9	56.3	58.0	56.6	60.1	60.8	57.9	57.9	57.5	64.4	64.5	63.9	65.1	65.6	63.8	69.6	70.4	69.5	
3-year	40.4	42.9	44.4	45.8	46.3	44.9	46.9	48.3	44.6	46.9	47.6	54.5	55.0	55.9	58.0	57.5	54.0	63.3	62.4		
4-year	32.2	34.9	34.8	36.8	39.0	36.4	36.7	35.5	34.2	38.9	41.6	46.8	47.5	49.2	50.2	49.7	46.8	58.2			
5-year	26.1	28.0	26.5	29.4	30.4	29.8	31.1	29.7	27.3	32.9	34.5	37.6	41.4	43.7	45.1	43.8	42.3				
6-year	21.0	21.2	20.7	23.8	24.7	24.4	26.3	25.0	23.8	28.3	29.8	33.6	37.1	38.5	40.8	37.9					
7-year	17.0	16.6	15.9	19.5	21.3	21.5	22.6	20.2	20.2	24.4	26.5	29.7	31.2	33.8	36.7						
8-year	14.1	13.0	13.7	16.0	18.5	18.5	19.4	18.3	18.0	21.8	23.7	26.7	28.3	29.5							
9-year	12.0	10.8	10.9	13.9	16.3	17.7	16.1	15.2	16.6	21.3	21.1	23.9	24.9								
10-year	10.5	9.4	9.6	12.5	15.1	15.9	14.5	13.5	14.7	20.3	19.6	21.0									
11-year	10.0	8.4	8.4	11.6	13.7	14.1	13.6	12.0	13.5	19.7	18.1										
12-year	8.7	7.6	7.3	10.1	12.8	13.0	11.4	10.6	13.1	18.4											
13-year	7.5	6.6	6.7	9.3	12.0	11.9	10.2	10.4	11.7												
14-year	6.9	6.0	6.0	8.4	11.4	11.0	9.4	10.1													
15-year	6.0	5.7	5.3	8.3	10.7	10.2	9.4														
16-year	5.4	5.1	4.7	7.5	10.1	9.8															
17-year	5.2	4.6	4.5	7.2	9.3																
18-year	4.5	4.2	4.1	6.8																	
19-year	4.3	3.4	3.8																		
20-year	3.9	3.1	3.5																		

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).