

Table 18.12
Myeloma

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years,
Lifetime Risk of Being Diagnosed with Cancer Given Alive and Cancer-Free at Current Age, and
Lifetime Risk of Dying from Cancer Given Alive at Current Age
Both Sexes, 2010-2012 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer
		+10 yrs	+20 yrs	+30 yrs	Ever				+10 yrs	+20 yrs	+30 yrs	Ever	
All Races	0	0.00	0.00	0.00	0.74	0.43	Asian/ Pacific Islander	0	0.00	0.00	0.00	0.54	0.31
	10	0.00	0.00	0.01	0.75	0.43		10	0.00	0.00	0.01	0.55	0.31
	20	0.00	0.01	0.04	0.75	0.43		20	0.00	0.01	0.02	0.55	0.31
	30	0.01	0.04	0.12	0.76	0.44		30	0.00	0.02	0.07	0.55	0.31
	40	0.03	0.11	0.28	0.76	0.44		40	0.02	0.06	0.16	0.55	0.31
	50	0.08	0.26	0.52	0.75	0.45		50	0.05	0.15	0.33	0.54	0.31
	60	0.19	0.47	0.66	0.70	0.44		60	0.10	0.29	0.43	0.50	0.31
	70	0.32	0.54	-	0.58	0.41		70	0.20	0.36	-	0.43	0.29
80	0.30	-	-	0.36	0.32	80	0.18	-	-	0.27	0.21		
White	0	0.00	0.00	0.00	0.68	0.40	American Indian/ Alaska Native ^a	0	0.00	0.00	0.00	0.54	0.27
	10	0.00	0.00	0.01	0.69	0.40		10	0.00	0.00	0.01	0.55	0.28
	20	0.00	0.01	0.03	0.69	0.41		20	0.00	0.01	0.04	0.55	0.28
	30	0.01	0.03	0.10	0.69	0.41		30	0.00	0.04	0.07	0.55	0.28
	40	0.03	0.10	0.25	0.70	0.41		40	0.03	0.07	0.19	0.56	0.29
	50	0.07	0.23	0.48	0.69	0.42		50	0.04	0.16	0.29	0.55	0.30
	60	0.17	0.43	0.62	0.65	0.42		60	0.13	0.27	0.43	0.55	0.30
	70	0.29	0.50	-	0.54	0.39		70	0.15	0.34	-	0.47	0.28
80	0.28	-	-	0.34	0.30	80	0.26	-	-	0.44	0.19		
Black	0	0.00	0.00	0.00	1.27	0.68	Hispanic ^b	0	0.00	0.00	0.00	0.79	0.43
	10	0.00	0.00	0.01	1.30	0.69		10	0.00	0.00	0.01	0.80	0.44
	20	0.00	0.01	0.08	1.31	0.70		20	0.00	0.01	0.04	0.80	0.44
	30	0.01	0.08	0.25	1.32	0.71		30	0.01	0.03	0.11	0.80	0.44
	40	0.07	0.24	0.58	1.33	0.72		40	0.03	0.10	0.27	0.80	0.44
	50	0.18	0.54	0.98	1.31	0.73		50	0.07	0.24	0.51	0.79	0.44
	60	0.39	0.88	1.18	1.24	0.73		60	0.18	0.46	0.67	0.74	0.44
	70	0.59	0.95	-	1.03	0.69		70	0.31	0.54	-	0.62	0.40
80	0.53	-	-	0.64	0.54	80	0.30	-	-	0.40	0.29		

Devcan 6.8.0, August 2014, National Cancer Institute (<http://surveillance.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic could not be calculated.

A percent of 0.00 represents a value that is below 0.005.