

Table 18.13
Myeloma

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years,
Lifetime Risk of Being Diagnosed with Cancer Given Alive and Cancer-Free at Current Age, and
Lifetime Risk of Dying from Cancer Given Alive at Current Age
Males, 2010-2012 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer
		+10 yrs	+20 yrs	+30 yrs	Ever				+10 yrs	+20 yrs	+30 yrs	Ever	
All Races	0	0.00	0.00	0.00	0.85	0.47	Asian/ Pacific Islander	0	0.00	0.00	0.00	0.62	0.36
	10	0.00	0.00	0.01	0.86	0.48		10	0.00	0.00	0.01	0.62	0.37
	20	0.00	0.01	0.04	0.87	0.48		20	0.00	0.01	0.03	0.62	0.37
	30	0.01	0.04	0.13	0.88	0.49		30	0.01	0.02	0.08	0.63	0.37
	40	0.03	0.13	0.33	0.88	0.49		40	0.02	0.08	0.19	0.63	0.37
	50	0.10	0.30	0.62	0.87	0.50		50	0.06	0.17	0.39	0.62	0.37
	60	0.22	0.56	0.79	0.84	0.51		60	0.12	0.35	0.52	0.58	0.37
	70	0.40	0.66	-	0.71	0.48		70	0.25	0.44	-	0.51	0.35
80	0.38	-	-	0.46	0.38	80	0.24	-	-	0.32	0.27		
White	0	0.00	0.00	0.00	0.81	0.45	American Indian/ Alaska Native ^a	0	0.00	0.00	0.00	0.49	0.29
	10	0.00	0.00	0.01	0.82	0.46		10	0.00	0.00	0.01	0.50	0.30
	20	0.00	0.01	0.04	0.82	0.46		20	0.00	0.00	0.05	0.50	0.30
	30	0.01	0.04	0.12	0.83	0.47		30	0.00	0.05	0.08	0.51	0.31
	40	0.03	0.11	0.30	0.83	0.47		40	0.05	0.08	0.17	0.52	0.32
	50	0.09	0.28	0.58	0.83	0.48		50	0.03	0.13	0.28	0.50	0.33
	60	0.21	0.53	0.75	0.80	0.49		60	0.11	0.26	0.37	0.51	0.34
	70	0.38	0.64	-	0.68	0.46		70	0.18	0.30	-	0.47	0.33
80	0.37	-	-	0.44	0.37	80	0.18	-	-	0.41	0.24		
Black	0	0.00	0.00	0.00	1.33	0.69	Hispanic ^b	0	0.00	0.00	0.00	0.93	0.47
	10	0.00	0.00	0.02	1.36	0.70		10	0.00	0.00	0.01	0.94	0.48
	20	0.00	0.02	0.09	1.37	0.71		20	0.00	0.01	0.04	0.94	0.48
	30	0.02	0.09	0.27	1.39	0.72		30	0.01	0.04	0.12	0.95	0.48
	40	0.08	0.26	0.65	1.41	0.74		40	0.03	0.12	0.31	0.95	0.49
	50	0.19	0.60	1.09	1.40	0.76		50	0.09	0.29	0.62	0.94	0.49
	60	0.45	1.00	1.30	1.35	0.77		60	0.21	0.56	0.82	0.90	0.49
	70	0.70	1.07	-	1.14	0.74		70	0.40	0.68	-	0.78	0.46
80	0.60	-	-	0.71	0.59	80	0.38	-	-	0.51	0.33		

Devcan 6.8.0, August 2014, National Cancer Institute (<http://surveillance.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic could not be calculated.

A percent of 0.00 represents a value that is below 0.005.