

Table 19.4
Non-Hodgkin Lymphoma

Delay-adjusted SEER Incidence^a Rates by Year, Race and Sex

Year of Diagnosis:	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
1975	11.07	12.89	9.59	11.61	13.49	10.11	6.36	7.31	5.31
1976	11.22	12.82	10.06	11.65	13.27	10.51	6.04	7.45	5.03
1977	11.18	12.93	9.82	11.64	13.52	10.20	6.64	7.23	6.10
1978	11.92	14.28	10.12	12.48	14.77	10.74	7.15	9.46	5.17
1979	12.53	14.21	11.21	12.99	14.59	11.77	8.46	11.95	5.91
1980	12.62	15.01	10.81	12.97	15.26	11.22	8.77	10.89	7.20
1981	13.58	15.90	11.81	14.06	16.34	12.35	7.42	10.59	4.98
1982	13.39	15.56	11.61	14.02	16.42	12.05	10.27	10.22	10.01
1983	13.99	16.62	11.95	14.63	17.52	12.39	10.28	10.88	9.62
1984	15.19	18.29	12.79	15.96	19.10	13.54	9.46	12.52	6.92
1985	15.51	18.39	13.17	16.28	19.24	13.86	9.95	11.96	8.43
1986	15.91	19.42	13.03	16.83	20.55	13.78	10.66	13.13	8.52
1987	16.74	20.79	13.42	17.70	22.14	14.00	10.40	10.55	9.95
1988	17.25	21.32	14.08	18.20	22.41	14.90	11.58	15.79	8.49
1989	17.35	21.65	13.84	18.35	23.02	14.50	11.62	14.17	9.72
1990	18.51	22.98	14.81	19.33	23.94	15.50	13.92	17.63	11.02
1991	18.81	23.74	14.60	19.68	24.81	15.29	14.03	18.72	10.37
1992	18.63	23.05	14.94	19.56	23.95	15.84	13.68	18.87	9.65
1993	18.86	23.47	15.01	19.65	24.30	15.71	14.11	19.37	9.78
1994	19.95	24.96	15.80	20.68	25.58	16.56	14.43	21.52	8.77
1995	19.99	25.43	15.43	20.68	26.16	16.02	16.70	23.54	11.38
1996	19.42	24.21	15.49	20.26	25.19	16.20	15.46	19.19	12.30
1997	19.98	24.51	16.44	20.86	25.20	17.44	16.31	24.10	10.84
1998	19.61	22.95	16.84	20.58	24.00	17.74	14.43	16.78	12.55
1999	19.96	24.59	16.34	20.92	25.49	17.31	14.40	18.45	11.32
2000	19.78	24.30	16.12	20.93	25.64	17.07	13.91	18.27	10.68
2001	20.03	24.41	16.48	20.84	25.47	17.08	16.06	18.39	13.97
2002	20.24	24.23	17.10	21.36	25.55	17.98	14.58	17.34	12.50
2003	20.80	25.08	17.56	21.96	26.40	18.52	16.12	19.73	13.60
2004	21.46	25.98	17.99	22.43	26.95	18.91	18.55	24.13	14.51
2005	20.94	25.71	17.13	22.13	26.91	18.27	16.96	20.33	14.23
2006	20.60	24.81	17.33	21.87	26.18	18.47	16.45	20.99	13.08
2007	21.37	26.27	17.47	22.72	27.80	18.55	15.81	19.30	13.18
2008	21.05	25.89	17.10	21.98	27.05	17.79	17.06	19.28	15.07
2009	21.06	25.40	17.54	22.17	26.87	18.24	15.45	18.91	12.90
2010	21.70	26.99	17.40	22.66	27.81	18.43	17.94	23.44	13.58
2011	20.28	24.88	16.63	21.26	26.02	17.45	14.92	17.94	12.68
2012	20.88	25.62	17.05	21.93	26.80	17.92	17.19	19.63	15.00

Additional information on the model used to delay-adjust SEER Incidence rates can be found at [\(http://surveillance.cancer.gov/delay/\)](http://surveillance.cancer.gov/delay/).

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Delay-adjusted rate is not shown for observed rates based on less than 16 cases for the time interval.