

Table 19.14  
Non-Hodgkin Lymphoma

Number of Cases

By SEER Location and Year of Diagnosis, 1975-2012

All Races, Males, Ages 20-54

Year of Diagnosis

| SEER Location                               | 1975-<br>1979 | 1980-<br>1981 | 1982-<br>1983 | 1984-<br>1985 | 1986-<br>1987 | 1988-<br>1989 | 1990-<br>1991 | 1992-<br>1993 | 1994-<br>1995 | 1996-<br>1997 | 1998-<br>1999 | 2000-<br>2001 | 2002-<br>2003 | 2004-<br>2005 | 2006-<br>2007 | 2008-<br>2009 | 2010-<br>2012 |
|---------------------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| SEER 9 <sup>b</sup>                         | 1,464         | 687           | 723           | 958           | 1,210         | 1,410         | 1,765         | 1,831         | 2,090         | 1,764         | 1,607         | 1,689         | 1,621         | 1,716         | 1,650         | 1,689         | 2,454         |
| SEER 9 except<br>SF-Oakland                 | 1,208         | 563           | 588           | 730           | 826           | 921           | 1,194         | 1,295         | 1,484         | 1,321         | 1,269         | 1,320         | 1,307         | 1,403         | 1,378         | 1,397         | 2,054         |
| San Francisco-<br>Oakland SMSA <sup>c</sup> | 256           | 124           | 135           | 228           | 384           | 489           | 571           | 536           | 606           | 443           | 338           | 369           | 314           | 313           | 272           | 292           | 400           |
| San Francisco<br>City/County                | 58            | 29            | 34            | 105           | 215           | 288           | 333           | 303           | 317           | 224           | 105           | 115           | 111           | 112           | 86            | 75            | 113           |

Age-Adjusted Cancer Incidence Rates<sup>a</sup>

By SEER Location and Year of Diagnosis, 1975-2012

All Races, Males, Ages 20-54

Year of Diagnosis

| SEER Location                               | 1975-<br>1979 | 1980-<br>1981 | 1982-<br>1983 | 1984-<br>1985 | 1986-<br>1987 | 1988-<br>1989 | 1990-<br>1991 | 1992-<br>1993 | 1994-<br>1995 | 1996-<br>1997 | 1998-<br>1999 | 2000-<br>2001 | 2002-<br>2003 | 2004-<br>2005 | 2006-<br>2007 | 2008-<br>2009 | 2010-<br>2012 |
|---------------------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| SEER 9 <sup>b</sup>                         | 6.9           | 7.9           | 8.2           | 10.4          | 12.4          | 13.5          | 15.9          | 15.6          | 16.9          | 13.8          | 12.1          | 12.3          | 11.7          | 12.2          | 11.6          | 11.8          | 11.3          |
| SEER 9 except<br>SF-Oakland                 | 6.9           | 7.8           | 8.1           | 9.7           | 10.3          | 10.8          | 13.1          | 13.4          | 14.4          | 12.3          | 11.5          | 11.5          | 11.2          | 11.9          | 11.5          | 11.6          | 11.3          |
| San Francisco-<br>Oakland SMSA <sup>c</sup> | 6.9           | 8.4           | 8.9           | 13.9          | 22.6          | 26.4          | 29.5          | 26.7          | 29.8          | 21.2          | 15.5          | 16.4          | 13.9          | 14.1          | 12.3          | 13.0          | 11.5          |
| San Francisco<br>City/County                | 6.7           | 8.8           | 9.4           | 27.2          | 56.5          | 71.5          | 81.3          | 72.4          | 74.4          | 50.7          | 23.0          | 25.2          | 23.9          | 25.1          | 19.4          | 16.2          | 15.5          |

<sup>a</sup> Rates are per 100,000 males aged 20-54 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> The SEER 9 areas are San Francisco-Oakland SMSA, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta.  
<sup>c</sup> The San Francisco-Oakland SMSA represents 15% of all persons in the SEER areas. For this age group, all SEER areas may not be generalizable to the United States for Non-Hodgkin Lymphoma.

- Statistic not shown. Rate based on less than 4 cases for the time interval.