

Table 19.17  
Non-Hodgkin Lymphoma

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years,  
Lifetime Risk of Being Diagnosed with Cancer Given Alive and Cancer-Free at Current Age, and  
Lifetime Risk of Dying from Cancer Given Alive at Current Age  
Males, 2010-2012 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer
		+10 yrs	+20 yrs	+30 yrs	Ever				+10 yrs	+20 yrs	+30 yrs	Ever	
All Races	0	0.01	0.03	0.06	2.37	0.86	Asian/ Pacific Islander	0	0.01	0.03	0.05	2.09	0.83
	10	0.02	0.05	0.12	2.39	0.88		10	0.02	0.04	0.09	2.10	0.84
	20	0.03	0.10	0.24	2.38	0.88		20	0.03	0.07	0.16	2.09	0.84
	30	0.07	0.21	0.48	2.37	0.88		30	0.04	0.13	0.33	2.08	0.84
	40	0.14	0.42	0.94	2.34	0.89		40	0.09	0.29	0.68	2.05	0.84
	50	0.29	0.82	1.57	2.27	0.90		50	0.20	0.60	1.21	1.99	0.84
	60	0.58	1.38	2.00	2.14	0.90		60	0.41	1.06	1.66	1.86	0.83
	70	0.94	1.66	-	1.82	0.87		70	0.70	1.36	-	1.58	0.78
80	1.03	-	-	1.26	0.75	80	0.83	-	-	1.10	0.65		
White	0	0.01	0.03	0.06	2.47	0.91	American Indian/ Alaska Native <sup>a</sup>	0	0.00	0.02	0.04	1.37	0.58
	10	0.02	0.05	0.12	2.49	0.92		10	0.01	0.03	0.09	1.38	0.59
	20	0.03	0.10	0.24	2.48	0.92		20	0.02	0.08	0.15	1.38	0.59
	30	0.07	0.21	0.49	2.47	0.93		30	0.06	0.13	0.24	1.39	0.60
	40	0.14	0.43	0.97	2.44	0.93		40	0.07	0.19	0.54	1.37	0.62
	50	0.30	0.86	1.65	2.38	0.94		50	0.12	0.49	0.97	1.36	0.64
	60	0.60	1.46	2.10	2.24	0.95		60	0.41	0.93	1.27	1.36	0.66
	70	1.00	1.75	-	1.91	0.91		70	0.62	1.02	-	1.12	0.64
80	1.08	-	-	1.31	0.78	80	0.59	-	-	0.73	0.55		
Black	0	0.01	0.03	0.07	1.43	0.50	Hispanic <sup>b</sup>	0	0.01	0.03	0.06	2.39	0.82
	10	0.02	0.06	0.15	1.45	0.51		10	0.02	0.05	0.10	2.40	0.83
	20	0.04	0.13	0.28	1.44	0.51		20	0.03	0.08	0.20	2.39	0.83
	30	0.08	0.24	0.49	1.42	0.51		30	0.05	0.17	0.42	2.39	0.84
	40	0.16	0.41	0.76	1.37	0.51		40	0.12	0.38	0.86	2.36	0.84
	50	0.27	0.63	1.00	1.27	0.50		50	0.26	0.76	1.49	2.30	0.84
	60	0.40	0.82	1.07	1.12	0.49		60	0.52	1.30	1.96	2.16	0.83
	70	0.53	0.85	-	0.92	0.46		70	0.88	1.63	-	1.84	0.79
80	0.51	-	-	0.62	0.38	80	0.99	-	-	1.28	0.63		

Devcan 6.8.0, August 2014, National Cancer Institute (<http://surveillance.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG). Mortality data are from the NCHS public use data file for the total US.

<sup>a</sup> Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

<sup>b</sup> Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic could not be calculated.

A percent of 0.00 represents a value that is below 0.005.