

Table 19.18
Non-Hodgkin Lymphoma

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years,
Lifetime Risk of Being Diagnosed with Cancer Given Alive and Cancer-Free at Current Age, and
Lifetime Risk of Dying from Cancer Given Alive at Current Age
Females, 2010-2012 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer
		+10 yrs	+20 yrs	+30 yrs	Ever				+10 yrs	+20 yrs	+30 yrs	Ever	
All Races	0	0.01	0.02	0.04	1.89	0.68	Asian/ Pacific Islander	0	0.01	0.02	0.04	1.63	0.67
	10	0.01	0.03	0.08	1.91	0.69		10	0.01	0.03	0.07	1.63	0.68
	20	0.02	0.07	0.17	1.90	0.69		20	0.02	0.06	0.12	1.62	0.68
	30	0.05	0.15	0.35	1.89	0.69		30	0.04	0.10	0.24	1.60	0.68
	40	0.10	0.30	0.70	1.85	0.69		40	0.06	0.20	0.48	1.57	0.68
	50	0.21	0.61	1.19	1.79	0.70		50	0.14	0.42	0.84	1.52	0.67
	60	0.42	1.03	1.53	1.66	0.70		60	0.28	0.71	1.22	1.41	0.66
	70	0.67	1.22	-	1.36	0.67		70	0.45	0.98	-	1.19	0.62
White	80	0.72	-	-	0.90	0.56	American Indian/ Alaska Native ^a	80	0.62	-	-	0.86	0.51
	0	0.00	0.02	0.04	1.99	0.72		0	0.00	0.01	0.02	1.16	0.39
	10	0.01	0.03	0.08	2.00	0.72		10	0.01	0.01	0.05	1.17	0.39
	20	0.02	0.07	0.17	1.99	0.72		20	0.01	0.04	0.10	1.17	0.40
	30	0.05	0.15	0.36	1.98	0.73		30	0.04	0.09	0.27	1.17	0.40
	40	0.10	0.32	0.73	1.95	0.73		40	0.05	0.24	0.52	1.16	0.40
	50	0.22	0.64	1.26	1.89	0.73		50	0.19	0.48	0.83	1.14	0.41
	60	0.44	1.09	1.62	1.74	0.73		60	0.30	0.67	0.98	0.99	0.42
Black	70	0.71	1.29	-	1.43	0.70	Hispanic ^b	70	0.41	0.76	-	0.77	0.43
	80	0.75	-	-	0.93	0.58		80	0.45	-	-	0.46	0.29
	0	0.01	0.02	0.04	1.19	0.42		0	0.01	0.02	0.03	1.99	0.72
	10	0.01	0.03	0.10	1.21	0.43		10	0.01	0.03	0.07	2.00	0.72
	20	0.02	0.09	0.18	1.20	0.43		20	0.02	0.06	0.15	2.00	0.72
	30	0.06	0.16	0.33	1.18	0.42		30	0.04	0.13	0.33	1.98	0.72
	40	0.10	0.27	0.55	1.13	0.42		40	0.09	0.30	0.67	1.95	0.72
	50	0.18	0.46	0.79	1.07	0.42		50	0.21	0.59	1.18	1.89	0.72
	60	0.31	0.66	0.90	0.96	0.41		60	0.39	1.01	1.55	1.73	0.71
	70	0.40	0.68	-	0.76	0.38		70	0.66	1.25	-	1.44	0.67
	80	0.38	-	-	0.48	0.31		80	0.71	-	-	0.95	0.53

Devcan 6.8.0, August 2014, National Cancer Institute (<http://surveillance.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

- Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic could not be calculated.

A percent of 0.00 represents a value that is below 0.005.