


Table 19.22
Non-Hodgkin Lymphoma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012


Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	6.15	0.02		
High Five States				
Minnesota	7.15 ^b	0.16	01(01,11)	16.23 ^c
Michigan	7.02 ^b	0.11	02(01,12)	14.18
Kentucky	7.01 ^b	0.18	03(01,17)	13.89
West Virginia	6.93	0.25	04(01,23)	12.69
Oklahoma	6.89 ^b	0.19	05(01,21)	12.05
Low Five States				
Nevada	5.48	0.21	47(24,51)	-10.85
Wyoming	5.42	0.45	48(07,51)	-11.84
North Dakota	5.36	0.37	49(16,51)	-12.92
New Mexico	5.12 ^b	0.22	50(36,51)	-16.78 ^c
D.C.	5.06	0.43	51(17,51)	-17.68 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	6.27	0.16	22(08,36)	1.94	Montana	6.01	0.32	26(03,50)	-2.31
Alaska	6.37	0.56	18(01,51)	3.52	Nebraska	6.29	0.25	21(03,43)	2.29
Arizona	5.79	0.13	36(23,47)	-5.88	Nevada	5.48	0.21	47(24,51)	-10.85
Arkansas	6.64	0.20	12(01,27)	8.00	New Hampshire	5.56	0.28	44(18,51)	-9.64
California	5.84 ^b	0.06	33(25,42)	-5.14	New Jersey	5.79	0.11	35(23,47)	-5.84
Colorado	5.48 ^b	0.16	46(26,51)	-10.85	New Mexico	5.12 ^b	0.22	50(36,51)	-16.78 ^c
Connecticut	5.58	0.16	42(25,51)	-9.29	New York	5.89 ^b	0.07	29(23,42)	-4.33
Delaware	5.75	0.34	38(08,51)	-6.48	North Carolina	5.86	0.11	30(22,45)	-4.67
D.C.	5.06	0.43	51(17,51)	-17.68 ^c	North Dakota	5.36	0.37	49(16,51)	-12.92
Florida	5.84 ^b	0.07	32(25,43)	-5.13	Ohio	6.86 ^b	0.10	06(01,16)	11.50
Georgia	5.49 ^b	0.12	45(32,51)	-10.77	Oklahoma	6.89 ^b	0.19	05(01,21)	12.05
Hawaii	5.75	0.27	39(13,51)	-6.57	Oregon	6.45	0.17	17(04,29)	4.81
Idaho	6.32	0.29	20(02,46)	2.73	Pennsylvania	6.72 ^b	0.09	08(03,19)	9.16
Illinois	6.34	0.10	19(11,27)	3.12	Rhode Island	5.69	0.31	40(12,51)	-7.55
Indiana	6.69 ^b	0.14	11(02,22)	8.81	South Carolina	5.82	0.16	34(20,48)	-5.39
Iowa	6.70	0.19	10(01,24)	8.94	South Dakota	6.09	0.36	25(01,50)	-0.99
Kansas	6.51	0.21	16(02,31)	5.80	Tennessee	6.59	0.14	13(03,24)	7.06
Kentucky	7.01 ^b	0.18	03(01,17)	13.89	Texas	5.95	0.08	28(22,40)	-3.29
Louisiana	6.58	0.17	14(02,26)	6.95	Utah	5.78	0.24	37(16,50)	-6.08
Maine	6.83	0.29	07(01,28)	11.08	Vermont	6.23	0.41	23(01,50)	1.30
Maryland	5.57 ^b	0.14	43(27,50)	-9.43	Virginia	6.01	0.12	27(17,41)	-2.38
Massachusetts	5.68 ^b	0.12	41(25,49)	-7.69	Washington	6.51	0.14	15(04,26)	5.82
Michigan	7.02 ^b	0.11	02(01,12)	14.18	West Virginia	6.93	0.25	04(01,23)	12.69
Minnesota	7.15 ^b	0.16	01(01,11)	16.23 ^c	Wisconsin	6.70 ^b	0.15	09(02,22)	8.95
Mississippi	5.84	0.20	31(17,50)	-5.12	Wyoming	5.42	0.45	48(07,51)	-11.84
Missouri	6.20	0.14	24(13,36)	0.80					



Age-Adjusted Death
Rates per 100,000
Quantile Interval



- ^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- ^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
- ^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
- ^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.
- SE Standard error of the rate.
- PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.