


Table 19.23
Non-Hodgkin Lymphoma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	7.92	0.03		
High Five States				
Minnesota	9.36 ^b	0.28	01(01,14)	18.13 ^c
Michigan	9.16 ^b	0.20	02(01,13)	15.71 ^c
Ohio	9.00 ^b	0.18	03(01,16)	13.67
Oklahoma	8.89	0.32	04(01,23)	12.24
Maine	8.82	0.50	05(01,36)	11.40
Low Five States				
Delaware	6.87	0.56	47(12,51)	-13.24
Nevada	6.82	0.35	48(23,51)	-13.90
Wyoming	6.32	0.74	49(13,51)	-20.22 ^c
North Dakota	6.31	0.61	50(18,51)	-20.26 ^c
New Mexico	6.05 ^b	0.36	51(42,51)	-23.63 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	7.86	0.28	24(09,44)	-0.80	Montana	8.00	0.55	21(01,49)	0.98
Alaska	7.81	0.92	26(01,51)	-1.34	Nebraska	7.91	0.43	23(02,47)	-0.10
Arizona	7.19	0.22	42(24,49)	-9.17	Nevada	6.82	0.35	48(23,51)	-13.90
Arkansas	8.17	0.35	19(02,42)	3.10	New Hampshire	7.17	0.48	43(09,51)	-9.43
California	7.44 ^b	0.10	35(25,44)	-6.06	New Jersey	7.41	0.19	37(21,47)	-6.46
Colorado	7.02	0.27	46(24,50)	-11.32	New Mexico	6.05 ^b	0.36	51(42,51)	-23.63 ^c
Connecticut	7.13	0.29	44(22,50)	-9.91	New York	7.59	0.13	31(21,43)	-4.12
Delaware	6.87	0.56	47(12,51)	-13.24	North Carolina	7.47	0.20	34(20,47)	-5.67
D.C.	7.05	0.80	45(01,51)	-10.97	North Dakota	6.31	0.61	50(18,51)	-20.26 ^c
Florida	7.49 ^b	0.12	33(23,44)	-5.42	Ohio	9.00 ^b	0.18	03(01,16)	13.67
Georgia	7.32	0.21	39(21,48)	-7.53	Oklahoma	8.89	0.32	04(01,23)	12.24
Hawaii	7.41	0.47	36(07,50)	-6.44	Oregon	8.33	0.30	16(02,36)	5.15
Idaho	8.01	0.49	20(01,48)	1.15	Pennsylvania	8.75 ^b	0.16	08(02,19)	10.46
Illinois	8.20	0.18	18(08,30)	3.50	Rhode Island	7.67	0.55	28(01,50)	-3.17
Indiana	8.64	0.25	12(01,24)	9.11	South Carolina	7.64	0.28	30(12,47)	-3.58
Iowa	8.75	0.34	07(01,27)	10.49	South Dakota	7.51	0.60	32(02,51)	-5.19
Kansas	8.73	0.36	09(01,28)	10.20	Tennessee	8.70	0.25	10(01,24)	9.81
Kentucky	8.81	0.31	06(01,24)	11.27	Texas	7.66	0.13	29(19,40)	-3.23
Louisiana	8.57	0.31	13(01,30)	8.22	Utah	7.19	0.41	41(15,51)	-9.16
Maine	8.82	0.50	05(01,36)	11.40	Vermont	8.25	0.73	17(01,50)	4.15
Maryland	7.29	0.25	40(21,49)	-7.95	Virginia	7.93	0.22	22(11,41)	0.08
Massachusetts	7.38	0.22	38(21,48)	-6.83	Washington	8.35	0.24	15(03,30)	5.41
Michigan	9.16 ^b	0.20	02(01,13)	15.71 ^c	West Virginia	8.44	0.42	14(01,41)	6.58
Minnesota	9.36 ^b	0.28	01(01,14)	18.13 ^c	Wisconsin	8.65	0.25	11(01,25)	9.21
Mississippi	7.85	0.36	25(05,47)	-0.91	Wyoming	6.32	0.74	49(13,51)	-20.22 ^c
Missouri	7.79	0.24	27(12,44)	-1.68					



- ^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- ^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
- ^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
- ^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.
- SE Standard error of the rate.
- PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.