

Table 20.14  
Cancer of the Oral Cavity and Pharynx (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Survival Time																					
1-year	78.7	78.5	79.5	80.4	82.7	81.5	82.2	80.2	84.0	84.0	82.7	83.5	84.4	84.9	83.7	84.9	85.9	86.5	86.1	87.2	87.8
2-year	65.0	65.0	66.1	68.2	70.6	69.0	69.7	70.0	71.3	72.9	73.2	71.6	74.8	75.1	74.1	74.6	76.7	77.9	77.7	77.8	
3-year	59.0	58.2	59.0	61.4	64.7	62.6	64.6	63.7	64.0	67.3	67.5	65.5	69.9	69.8	68.3	69.2	71.0	72.3	72.1		
4-year	54.7	54.0	54.9	57.0	60.3	58.8	60.0	59.1	60.5	63.8	62.2	62.4	66.7	65.5	65.5	65.9	67.8	68.5			
5-year	51.6	50.6	51.4	54.1	57.6	56.3	56.3	55.9	57.2	60.8	59.6	60.4	65.1	63.7	63.6	63.9	65.7				
6-year	49.3	48.0	49.2	51.6	54.9	53.4	53.8	54.3	54.5	58.1	57.1	58.4	64.0	61.1	60.8	61.9					
7-year	47.2	45.9	47.0	49.4	51.9	51.0	52.6	52.6	52.6	56.0	55.3	57.5	62.7	59.9	59.1						
8-year	45.2	44.3	44.7	47.6	49.6	49.1	51.4	51.2	51.1	53.6	52.6	55.3	60.9	58.0							
9-year	43.1	42.5	42.7	45.3	47.5	48.0	49.7	50.2	49.4	51.7	50.5	53.0	58.2								
10-year	41.2	40.6	40.8	43.8	45.7	45.8	48.2	49.2	48.4	49.9	48.7	50.9									
11-year	39.2	39.0	39.5	42.2	43.7	44.1	46.3	47.0	47.3	48.2	47.3										
12-year	37.9	37.8	38.2	40.8	42.0	43.2	44.5	46.1	46.3	46.5											
13-year	36.8	36.3	36.7	39.3	40.2	40.9	42.8	44.5	45.6												
14-year	35.8	35.3	35.5	38.0	36.9	39.8	41.3	42.7													
15-year	34.5	33.9	34.5	36.9	34.9	38.1	40.1														
16-year	33.1	32.7	33.6	35.7	33.9	37.1															
17-year	32.0	31.7	32.6	34.5	32.9																
18-year	31.1	30.5	31.4	33.4																	
19-year	30.4	28.7	29.8																		
20-year	29.3	28.1	29.1																		

<sup>a</sup>

Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).