


Table 20.26
Cancer of the Oral Cavity and Pharynx (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012


Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	1.35	0.01		
High Five States				
Washington	1.91 ^b	0.10	01(01,08)	41.31 ^c
Vermont	1.81	0.28	02(01,45)	34.40 ^c
Mississippi	1.75	0.14	03(01,29)	29.46 ^c
D.C.	1.73	0.32	04(01,50)	28.15 ^c
Arkansas	1.69	0.14	05(01,30)	25.12 ^c
Low Five States				
Rhode Island	1.08	0.18	46(05,50)	-19.84 ^c
Iowa	1.07	0.10	47(21,50)	-21.02 ^c
Connecticut	1.01 ^b	0.09	48(30,50)	-25.43 ^c
Arizona	1.00 ^b	0.07	49(36,50)	-26.19 ^c
North Dakota	0.94	0.23	50(04,50)	-30.06 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	1.39	0.10	19(05,44)	2.64	Montana	1.21	0.20	40(02,50)	-10.07
Alaska	-	-	-(-,-)	-	Nebraska	1.61	0.17	07(01,43)	19.15 ^c
Arizona	1.00 ^b	0.07	49(36,50)	-26.19 ^c	Nevada	1.31	0.14	31(04,49)	-3.14
Arkansas	1.69	0.14	05(01,30)	25.12 ^c	New Hampshire	1.30	0.19	32(02,50)	-3.86
California	1.41	0.04	15(08,32)	4.45	New Jersey	1.25	0.07	36(13,46)	-7.09
Colorado	1.28	0.10	35(07,48)	-5.31	New Mexico	1.32	0.15	28(03,50)	-2.17
Connecticut	1.01 ^b	0.09	48(30,50)	-25.43 ^c	New York	1.22	0.04	39(23,46)	-9.73
Delaware	1.24	0.21	38(02,50)	-8.10	North Carolina	1.40	0.07	16(06,40)	3.47
D.C.	1.73	0.32	04(01,50)	28.15 ^c	North Dakota	0.94	0.23	50(04,50)	-30.06 ^c
Florida	1.42	0.05	13(07,34)	5.01	Ohio	1.36	0.06	24(08,42)	0.37
Georgia	1.36	0.07	22(07,42)	0.59	Oklahoma	1.50	0.12	12(02,41)	11.21
Hawaii	1.32	0.17	30(02,50)	-2.57	Oregon	1.39	0.11	17(04,45)	3.05
Idaho	1.36	0.18	21(02,50)	0.97	Pennsylvania	1.25	0.05	37(16,45)	-7.22
Illinois	1.41	0.06	14(06,38)	4.46	Rhode Island	1.08	0.18	46(05,50)	-19.84 ^c
Indiana	1.28	0.08	34(09,47)	-5.16	South Carolina	1.36	0.10	23(05,45)	0.51
Iowa	1.07	0.10	47(21,50)	-21.02 ^c	South Dakota	1.17	0.21	42(03,50)	-13.42
Kansas	1.09	0.11	45(13,50)	-19.42 ^c	Tennessee	1.55	0.09	09(02,32)	14.62
Kentucky	1.51	0.11	11(02,39)	12.05	Texas	1.33	0.05	26(12,41)	-1.37
Louisiana	1.52	0.11	10(02,39)	12.91	Utah	1.12	0.14	44(09,50)	-16.75 ^c
Maine	1.39	0.17	18(02,50)	2.78	Vermont	1.81	0.28	02(01,45)	34.40 ^c
Maryland	1.33	0.09	25(06,46)	-1.35	Virginia	1.21	0.07	41(15,48)	-10.59
Massachusetts	1.33	0.08	27(07,45)	-1.78	Washington	1.91 ^b	0.10	01(01,08)	41.31 ^c
Michigan	1.32	0.07	29(09,44)	-2.24	West Virginia	1.14	0.13	43(08,50)	-15.79 ^c
Minnesota	1.29	0.09	33(07,47)	-4.29	Wisconsin	1.55	0.10	08(02,33)	15.10 ^c
Mississippi	1.75	0.14	03(01,29)	29.46 ^c	Wyoming	1.61	0.32	06(01,50)	19.60 ^c
Missouri	1.37	0.08	20(06,44)	1.66					



Age-Adjusted Death
Rates per 100,000
Quantile Interval



U.S. rate: 1.35

- ^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- ^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
- ^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
- ^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.
- SE Standard error of the rate.
- PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.