

Table 21.7  
Cancer of the Ovary (Invasive)

SEER Incidence<sup>a</sup> and U.S. Death<sup>b</sup> Rates, Age-Adjusted and Age-Specific Rates, by Race

	SEER Incidence <sup>a</sup>			U.S. Mortality <sup>b</sup>		
	All Races Females	White Females	Black Females	All Races Females	White Females	Black Females
<u>Age-Adjusted Rates, 2008-2012</u>						
Age at Diagnosis/Death:						
All ages	12.1	12.8	9.8	7.7	8.0	6.7
Under 65	7.4	7.8	5.5	3.0	3.1	2.6
65 and over	44.5	47.1	39.7	40.1	41.6	34.6
All ages (IARC world std) <sup>c</sup>	8.7	9.1	6.9	4.8	5.0	4.2
<u>Age-Specific Rates, 2008-2012</u>						
Age at Diagnosis/Death:						
<1	-	-	-	-	-	-
1-4	-	-	-	-	-	-
5-9	0.3	0.2	-	-	-	-
10-14	0.8	0.8	-	-	-	-
15-19	1.4	1.4	1.2	0.1	0.1	-
20-24	1.7	1.7	1.6	0.1	0.1	-
25-29	2.5	2.6	2.3	0.3	0.3	0.4
30-34	3.3	3.3	3.1	0.6	0.6	0.6
35-39	4.9	5.1	3.5	1.1	1.1	1.2
40-44	9.1	9.6	5.6	2.3	2.3	2.2
45-49	14.5	15.1	9.5	4.7	4.9	4.0
50-54	19.9	20.9	13.5	8.6	9.0	7.3
55-59	24.7	26.1	18.3	13.2	13.8	10.9
60-64	30.4	32.3	25.1	19.7	20.6	16.5
65-69	37.7	40.3	33.1	27.6	28.6	24.9
70-74	44.0	46.9	40.6	35.7	37.1	33.1
75-79	47.9	50.9	39.5	44.5	46.3	37.0
80-84	49.3	51.9	46.0	52.4	54.3	43.8
85+	49.0	50.8	45.7	55.1	56.8	44.8

- <sup>a</sup> Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473. SEER 18 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.
- <sup>b</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.
- <sup>c</sup> Rates are per 100,000 and are age-adjusted to the IARC world standard population.
- Statistic not shown. Rate based on less than 16 cases for the time interval.