



Table 21.14  
Cancer of the Ovary (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2008-2012

Females

State	Rate	SE	Rank(CI) <sup>d</sup>	PD
TOTAL U.S.	7.71	0.03		
High Five States				
Oregon	8.98 <sup>b</sup>	0.28	01(01,12)	16.58 <sup>c</sup>
Washington	8.70 <sup>b</sup>	0.22	02(01,15)	12.91
Montana	8.67	0.52	03(01,41)	12.54
Alabama	8.65 <sup>b</sup>	0.24	04(01,19)	12.30
Iowa	8.44	0.29	05(01,31)	9.53
Low Five States				
Louisiana	6.96	0.23	47(28,50)	-9.71
North Dakota	6.78	0.57	48(05,51)	-11.97
Alaska	6.70	0.74	49(01,51)	-13.07
Rhode Island	6.57	0.44	50(19,51)	-14.70
Hawaii	5.31 <sup>b</sup>	0.36	51(49,51)	-31.13 <sup>c</sup>

State	Rate	SE	Rank(CI) <sup>d</sup>	PD	State	Rate	SE	Rank(CI) <sup>d</sup>	PD
Alabama	8.65 <sup>b</sup>	0.24	04(01,19)	12.30	Montana	8.67	0.52	03(01,41)	12.54
Alaska	6.70	0.74	49(01,51)	-13.07	Nebraska	7.00	0.36	45(13,50)	-9.11
Arizona	7.35	0.20	36(17,48)	-4.64	Nevada	6.97	0.32	46(17,50)	-9.57
Arkansas	8.16	0.30	10(01,38)	5.93	New Hampshire	7.66	0.43	28(01,49)	-0.60
California	7.67	0.09	27(16,37)	-0.48	New Jersey	8.26	0.17	07(02,25)	7.25
Colorado	8.26	0.25	08(01,31)	7.22	New Mexico	7.84	0.36	20(01,48)	1.78
Connecticut	7.32	0.25	38(14,49)	-5.03	New York	7.74	0.11	24(13,37)	0.41
Delaware	7.67	0.52	26(01,50)	-0.43	North Carolina	7.51	0.16	33(15,44)	-2.50
D.C.	7.17	0.66	40(01,51)	-6.99	North Dakota	6.78	0.57	48(05,51)	-11.97
Florida	7.04 <sup>b</sup>	0.10	43(34,49)	-8.62	Ohio	7.96	0.15	16(06,33)	3.27
Georgia	7.87	0.18	19(06,38)	2.17	Oklahoma	8.06	0.27	13(02,39)	4.55
Hawaii	5.31 <sup>b</sup>	0.36	51(49,51)	-31.13 <sup>c</sup>	Oregon	8.98 <sup>b</sup>	0.28	01(01,12)	16.58 <sup>c</sup>
Idaho	7.58	0.42	32(02,50)	-1.67	Pennsylvania	8.26 <sup>b</sup>	0.14	09(03,22)	7.12
Illinois	7.77	0.15	23(09,38)	0.87	Rhode Island	6.57	0.44	50(19,51)	-14.70
Indiana	8.15	0.21	11(02,33)	5.81	South Carolina	7.81	0.24	22(05,42)	1.35
Iowa	8.44	0.29	05(01,31)	9.53	South Dakota	7.09	0.53	41(03,51)	-8.01
Kansas	7.88	0.31	18(02,44)	2.25	Tennessee	7.72	0.20	25(08,42)	0.16
Kentucky	7.48	0.24	34(10,48)	-2.93	Texas	7.26 <sup>b</sup>	0.11	39(29,46)	-5.80
Louisiana	6.96	0.23	47(28,50)	-9.71	Utah	7.39	0.36	35(05,50)	-4.05
Maine	7.62	0.40	30(02,49)	-1.12	Vermont	7.32	0.60	37(01,51)	-5.01
Maryland	7.64	0.21	29(08,45)	-0.89	Virginia	8.03	0.19	14(04,35)	4.25
Massachusetts	7.61	0.19	31(10,44)	-1.27	Washington	8.70 <sup>b</sup>	0.22	02(01,15)	12.91
Michigan	7.92	0.16	17(06,36)	2.75	West Virginia	8.01	0.36	15(01,45)	3.88
Minnesota	7.84	0.23	21(04,41)	1.68	Wisconsin	8.12	0.22	12(02,34)	5.38
Mississippi	7.00	0.29	44(19,50)	-9.11	Wyoming	8.33	0.75	06(01,50)	8.15
Missouri	7.05	0.19	42(27,50)	-8.50					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p < .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.

<sup>d</sup> Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.