

Table 22.4
Cancer of the Pancreas (Invasive)

Delay-adjusted SEER Incidence^a Rates by Year, Race and Sex

Year of Diagnosis:	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
1975	11.84	15.63	9.12	11.70	15.66	8.89	15.90	18.08	13.91
1976	11.99	14.82	9.90	11.76	14.50	9.74	17.61	22.68	13.97
1977	11.90	14.81	9.83	11.62	14.57	9.54	17.12	19.44	15.25
1978	11.07	14.06	8.84	10.83	13.90	8.60	15.17	20.08	11.53
1979	11.52	14.15	9.65	11.21	13.84	9.39	16.08	19.14	13.68
1980	11.52	14.20	9.61	11.07	13.70	9.18	17.94	21.76	15.53
1981	11.68	13.54	10.17	11.39	13.26	9.91	17.18	20.86	14.10
1982	11.72	13.81	10.18	11.41	13.46	9.95	16.86	20.99	13.96
1983	12.19	14.66	10.44	11.75	14.07	10.12	18.13	23.05	14.90
1984	12.22	13.94	10.93	12.03	13.87	10.64	17.19	18.36	16.34
1985	12.10	14.35	10.50	11.70	13.60	10.34	18.80	25.28	14.55
1986	11.68	13.78	10.21	11.37	13.54	9.82	17.32	20.16	15.59
1987	11.60	13.81	10.09	11.09	13.38	9.54	18.32	18.98	17.78
1988	11.65	13.70	10.20	11.11	13.35	9.57	18.48	19.55	17.32
1989	11.33	12.88	10.07	10.98	12.66	9.63	15.02	16.12	14.00
1990	11.28	12.81	10.06	11.02	12.46	9.84	15.20	18.82	12.63
1991	11.39	12.90	10.22	11.04	12.61	9.79	17.01	18.20	16.16
1992	11.62	13.22	10.34	11.33	12.90	10.02	17.34	19.77	15.75
1993	10.97	12.54	9.70	10.51	11.98	9.30	16.97	18.90	15.33
1994	11.38	13.26	9.91	10.90	12.35	9.74	17.45	21.53	14.49
1995	11.13	12.72	9.96	10.71	12.31	9.57	18.11	20.31	16.31
1996	11.30	12.92	9.93	10.91	12.48	9.57	17.06	19.68	14.88
1997	11.48	13.07	10.18	11.02	12.56	9.72	17.92	19.47	16.55
1998	11.48	13.12	10.23	11.24	12.99	9.85	14.81	16.36	13.89
1999	11.10	12.58	9.83	10.71	12.23	9.35	16.70	19.54	14.70
2000	11.42	13.34	9.84	11.07	12.94	9.55	15.77	18.18	13.39
2001	11.34	13.18	9.99	11.06	13.07	9.53	14.31	15.55	13.34
2002	11.77	13.07	10.58	11.59	13.29	10.08	16.13	14.94	16.25
2003	11.66	12.91	10.61	11.44	12.70	10.35	16.16	17.13	15.43
2004	12.01	13.96	10.38	11.76	13.61	10.20	16.13	19.59	13.48
2005	12.08	13.80	10.69	11.76	13.46	10.34	17.15	18.61	16.40
2006	12.62	14.34	11.20	12.50	14.52	10.83	16.88	18.38	15.54
2007	12.50	14.44	10.92	12.34	14.44	10.61	15.60	16.42	14.87
2008	12.80	14.62	11.31	12.44	14.27	10.91	17.36	19.45	15.87
2009	13.18	14.81	11.83	12.87	14.53	11.50	17.57	20.61	15.23
2010	12.70	14.02	11.56	12.30	13.67	11.11	17.23	19.31	15.61
2011	12.62	14.73	10.90	12.38	14.46	10.66	16.25	18.12	14.84
2012	13.42	15.01	12.10	13.29	15.39	11.56	17.46	16.60	17.63

Additional information on the model used to delay-adjust SEER Incidence rates can be found at [\(http://surveillance.cancer.gov/delay/\)](http://surveillance.cancer.gov/delay/).

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Delay-adjusted rate is not shown for observed rates based on less than 16 cases for the time interval.