

Table 22.21  
Cancer of the Pancreas (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2012  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <20 <sup>e</sup>	0 to <37 <sup>e</sup>	>=37 <sup>g</sup>	Complete <sup>h</sup>	
Race	Sex											
All Races <sup>b</sup>	Both Sexes	34,042	5,941	2,492	1,380	789	430	44,240	45,432	270	45,702	
	Males	17,628	2,644	1,241	479	395	107	22,190	22,666	117	22,783	
	Females	16,414	3,297	1,251	901	394	323	22,050	22,766	153	22,919	
White <sup>b</sup>	Both Sexes	27,956	5,052	1,946	1,019	597	332	36,320	37,195	220	37,415	
	Males	14,674	2,268	1,083	345	336	74	18,561	18,920	104	19,024	
	Females	13,282	2,784	863	674	261	258	17,759	18,275	116	18,391	
Black <sup>b</sup>	Both Sexes	4,327	581	345	199	153	60	5,476	5,722	127	5,849	
	Males	2,084	261	93	42	52	26	2,481	2,581	53	2,634	
	Females	2,243	320	252	157	101	34	2,995	3,141	74	3,215	
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	1,335	239	121	+	+	+	1,755	+	+	+	
	Males	614	109	41	+	+	+	789	+	+	+	
	Females	721	130	80	+	+	+	966	+	+	+	
Hispanic <sup>d</sup>	Both Sexes	2,386	473	203	+	+	+	3,216	+	+	+	
	Males	1,247	155	83	+	+	+	1,518	+	+	+	
	Females	1,139	318	120	+	+	+	1,698	+	+	+	

Estimated prevalence percent<sup>a</sup> on January 1, 2012, of the SEER<sup>c</sup> population diagnosed in the previous 20 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)										Age-Adjusted <sup>f</sup>
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
Race	Sex											
All Races <sup>c</sup>	Both Sexes	0.0129%	-	0.0001%	0.0008%	0.0020%	0.0070%	0.0187%	0.0404%	0.0624%	0.0617%	0.0123%
	Males	0.0130%	-	-	0.0005%	0.0016%	0.0073%	0.0209%	0.0449%	0.0690%	0.0661%	0.0133%
	Females	0.0129%	-	-	0.0011%	0.0025%	0.0068%	0.0167%	0.0363%	0.0570%	0.0591%	0.0115%
White <sup>c</sup>	Both Sexes	0.0134%	-	-	0.0007%	0.0018%	0.0070%	0.0180%	0.0407%	0.0623%	0.0601%	0.0122%
	Males	0.0137%	-	-	0.0005%	0.0015%	0.0073%	0.0204%	0.0457%	0.0702%	0.0658%	0.0134%
	Females	0.0130%	-	-	0.0009%	0.0022%	0.0066%	0.0156%	0.0361%	0.0557%	0.0567%	0.0111%
Black <sup>c</sup>	Both Sexes	0.0124%	-	-	0.0010%	0.0030%	0.0089%	0.0261%	0.0466%	0.0661%	0.0699%	0.0146%
	Males	0.0114%	-	-	-	0.0019%	0.0092%	0.0282%	0.0514%	0.0642%	0.0554%	0.0143%
	Females	0.0133%	-	-	0.0013%	0.0040%	0.0087%	0.0243%	0.0428%	0.0675%	0.0770%	0.0147%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0115%	-	-	0.0009%	0.0020%	0.0061%	0.0169%	0.0331%	0.0595%	0.0659%	0.0114%
	Males	0.0107%	-	-	-	0.0019%	0.0054%	0.0173%	0.0345%	0.0607%	0.0726%	0.0117%
	Females	0.0121%	-	-	0.0017%	0.0021%	0.0067%	0.0166%	0.0320%	0.0586%	0.0618%	0.0113%
Hispanic <sup>d</sup>	Both Sexes	0.0063%	-	-	0.0008%	0.0013%	0.0056%	0.0143%	0.0356%	0.0451%	0.0675%	0.0104%
	Males	0.0058%	-	-	0.0007%	0.0007%	0.0059%	0.0150%	0.0400%	0.0525%	0.0493%	0.0105%
	Females	0.0068%	-	-	0.0010%	0.0020%	0.0053%	0.0137%	0.0318%	0.0396%	0.0784%	0.0101%

<sup>a</sup> US 2012 cancer prevalence counts are based on 2012 cancer prevalence proportions from the SEER registries and 1/1/2012 US population estimates based on the average of 2011 and 2012 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 37 years for 1975-2012 SEER 9 data; 20 years for 1992-2012 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 37 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <37 and >=37. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.