

Table 23.12
Cancer of the Prostate (Invasive)

Age-Adjusted SEER Incidence Rates^a
By Registry, Race and Age

	All Races			Whites			Blacks		
	All Ages	Ages <65	Ages 65+	All Ages	Ages <65	Ages 65+	All Ages	Ages <65	Ages 65+
<u>SEER Incidence Rates^a, 2008-2012</u>									
California	126.90	47.68	674.52	122.44	46.39	648.18	189.62	92.54	860.61
Greater Bay Area	137.07	53.04	717.91	143.10	55.81	746.50	195.82	95.59	888.67
San Francisco-Oakland	132.24	51.80	688.28	135.03	53.11	701.31	197.57	95.47	903.37
San Jose-Monterey	146.43	55.40	775.66	157.22	60.51	825.70	180.99	94.80	776.77
Los Angeles	121.98	45.46	650.92	112.62	42.09	600.12	189.70	92.50	861.60
Greater California	125.53	46.75	670.10	120.24	45.27	638.49	186.64	91.09	847.08
Connecticut	139.87	61.02	684.92	131.13	55.91	651.07	207.74	97.64	968.76
Detroit	178.92	74.67	899.59	159.21	62.85	825.25	246.56	116.73	1143.98
Georgia	150.05	63.21	750.35	128.42	51.00	663.57	233.79	100.47	1155.41
Atlanta	168.00	71.16	837.42	144.13	57.94	739.93	239.81	103.35	1183.06
Rural Georgia	144.04	58.08	738.20	119.08	43.12	644.16	202.51	85.61	1010.57
Greater Georgia	142.74	59.40	718.77	123.42	48.49	641.33	230.93	98.77	1144.45
Hawaii	104.96	33.11	601.62	107.14	39.16	577.05	173.17	107.62	626.30
Iowa	126.18	49.80	654.16	122.81	48.34	637.53	161.93	71.99	783.59
Kentucky	122.64	49.55	627.91	113.35	44.02	592.59	169.43	83.53	763.20
Louisiana	161.08	66.75	813.11	143.25	57.23	737.81	222.25	94.68	1104.03
New Jersey	157.35	65.41	792.82	150.33	61.50	764.39	223.34	98.20	1088.41
New Mexico	110.38	42.29	581.02	110.88	43.55	576.32	128.72	60.44	600.67
Seattle-Puget Sound	142.26	54.63	747.95	142.65	54.64	750.96	192.70	94.72	869.91
Utah	156.78	59.47	829.36	156.49	60.27	821.57	191.98	75.65	996.08
9 SEER Areas ^b	142.41	57.35	730.43	136.95	54.35	707.91	223.99	103.41	1057.47
11 SEER Areas ^b	138.21	54.64	715.81	133.19	52.20	693.04	214.39	100.85	999.18
13 SEER Areas ^b	138.08	54.58	715.26	133.14	52.17	692.81	214.25	100.67	999.34
18 SEER Areas ^b	137.92	54.66	713.49	130.36	50.97	679.12	214.52	97.64	1022.42

^a Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130)

^b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta.

The SEER 11 areas comprise the SEER 9 areas plus San Jose-Monterey and Los Angeles.

The SEER 13 areas comprise the SEER 11 areas plus the Alaska Native Registry and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG.

- Statistic not shown. Rate based on less than 16 cases for the time interval.