

Table 26.4  
Cancer of the Thyroid (Invasive)

Delay-adjusted SEER Incidence<sup>a</sup> Rates by Year, Race and Sex

Year of Diagnosis:	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
1975	4.85	3.12	6.45	4.65	2.77	6.37	2.97	1.56	4.21
1976	4.79	2.94	6.57	4.68	2.88	6.41	3.37	1.33	5.00
1977	5.45	3.50	7.32	5.34	3.44	7.20	3.84	1.75	5.59
1978	5.09	3.14	6.95	4.95	2.97	6.83	2.95	1.68	4.07
1979	4.48	2.68	6.17	4.45	2.59	6.21	3.24	1.93	4.30
1980	4.33	2.39	6.15	4.30	2.28	6.21	2.37	1.11	3.43
1981	4.42	2.52	6.23	4.38	2.48	6.26	2.39	1.04	3.44
1982	4.63	3.00	6.15	4.53	2.87	6.09	3.29	2.02	4.34
1983	4.71	2.80	6.50	4.74	2.61	6.73	3.02	2.15	3.77
1984	4.85	2.65	6.93	4.73	2.58	6.81	3.02	1.73	4.08
1985	5.13	3.09	7.09	5.08	3.09	7.04	2.86	1.89	3.66
1986	5.32	3.07	7.49	5.24	3.10	7.34	3.35	1.19	5.09
1987	5.04	2.81	7.12	5.02	2.86	7.04	2.82	0.97	4.30
1988	4.93	2.94	6.88	4.89	3.00	6.78	2.53	1.80	3.21
1989	5.37	2.98	7.66	5.47	3.21	7.71	2.57	0.62	4.03
1990	5.50	2.93	7.96	5.56	2.99	8.09	3.76	2.28	4.91
1991	5.49	3.19	7.69	5.68	3.31	8.00	3.22	2.07	4.20
1992	5.88	3.52	8.13	5.89	3.52	8.19	4.20	3.09	5.13
1993	5.64	3.58	7.64	5.74	3.68	7.77	3.38	2.07	4.56
1994	6.08	3.38	8.74	6.22	3.43	9.00	3.75	1.78	5.35
1995	6.22	3.36	9.00	6.40	3.57	9.22	3.09	1.28	4.55
1996	6.53	3.48	9.51	6.72	3.62	9.82	3.85	1.70	5.80
1997	6.78	3.67	9.79	7.06	3.79	10.33	3.98	2.44	5.24
1998	6.98	3.70	10.13	7.21	3.95	10.43	3.85	1.50	5.80
1999	7.34	3.87	10.68	7.57	4.08	11.01	4.59	1.60	7.03
2000	7.62	4.04	11.11	8.08	4.38	11.81	4.58	2.19	6.56
2001	8.32	4.34	12.20	8.70	4.63	12.80	4.95	2.53	6.90
2002	9.26	4.96	13.46	9.80	5.29	14.34	5.48	2.79	7.66
2003	9.68	4.73	14.50	10.42	5.08	15.78	5.33	2.83	7.55
2004	10.18	5.27	14.98	10.89	5.60	16.24	6.42	3.41	8.92
2005	11.03	5.85	16.06	12.02	6.54	17.52	5.94	2.78	8.70
2006	11.40	5.90	16.73	12.28	6.44	18.16	6.71	3.16	9.57
2007	12.48	6.15	18.64	13.39	6.61	20.23	7.36	3.20	10.76
2008	13.34	6.66	19.83	14.38	7.15	21.66	7.11	4.47	9.57
2009	14.58	7.10	21.78	15.69	7.79	23.57	8.77	3.31	13.48
2010	14.08	6.71	21.22	14.95	7.35	22.60	8.75	3.46	13.12
2011	14.99	7.50	22.25	16.00	8.08	23.95	9.11	3.84	13.45
2012	15.24	8.20	22.15	16.16	8.88	23.59	9.75	4.36	14.30

Additional information on the model used to delay-adjust SEER Incidence rates can be found at <http://surveillance.cancer.gov/delay/>.

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Delay-adjusted rate is not shown for observed rates based on less than 16 cases for the time interval.