

Table 26.12
Cancer of the Thyroid (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years,
Lifetime Risk of Being Diagnosed with Cancer Given Alive and Cancer-Free at Current Age, and
Lifetime Risk of Dying from Cancer Given Alive at Current Age
Both Sexes, 2010-2012 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer
		+10 yrs	+20 yrs	+30 yrs	Ever				+10 yrs	+20 yrs	+30 yrs	Ever	
All Races	0	0.00	0.02	0.10	1.15	0.06	Asian/ Pacific Islander	0	0.00	0.02	0.09	1.24	0.12
	10	0.02	0.10	0.26	1.17	0.06		10	0.02	0.09	0.25	1.25	0.12
	20	0.08	0.24	0.46	1.15	0.06		20	0.07	0.23	0.45	1.24	0.12
	30	0.17	0.38	0.61	1.08	0.06		30	0.17	0.38	0.61	1.17	0.12
	40	0.22	0.45	0.67	0.93	0.06		40	0.22	0.45	0.68	1.02	0.12
	50	0.24	0.47	0.65	0.73	0.06		50	0.24	0.48	0.67	0.81	0.12
	60	0.24	0.43	0.51	0.52	0.06		60	0.25	0.45	0.55	0.59	0.12
	70	0.21	0.30	-	0.31	0.06		70	0.22	0.33	-	0.37	0.11
80	0.11	-	-	0.13	0.05	80	0.13	-	-	0.18	0.09		
White	0	0.00	0.02	0.11	1.21	0.06	American Indian/ Alaska Native ^a	0	0.00	0.01	0.04	0.77	0.04
	10	0.02	0.11	0.29	1.22	0.06		10	0.01	0.04	0.14	0.78	0.04
	20	0.09	0.26	0.49	1.21	0.06		20	0.04	0.13	0.25	0.77	0.04
	30	0.18	0.40	0.65	1.13	0.06		30	0.09	0.22	0.34	0.75	0.04
	40	0.23	0.47	0.71	0.96	0.06		40	0.13	0.25	0.41	0.67	0.05
	50	0.25	0.49	0.67	0.75	0.06		50	0.13	0.30	0.50	0.57	0.05
	60	0.25	0.44	0.52	0.53	0.06		60	0.18	0.40	0.46	0.46	0.05
	70	0.22	0.30	-	0.32	0.06		70	0.25	0.32	-	0.32	0.05
80	0.12	-	-	0.13	0.05	80	0.10	-	-	0.10	0.04		
Black	0	0.00	0.01	0.03	0.66	0.05	Hispanic ^b	0	0.00	0.02	0.08	1.08	0.09
	10	0.01	0.04	0.12	0.68	0.05		10	0.02	0.08	0.21	1.09	0.09
	20	0.03	0.11	0.24	0.67	0.05		20	0.06	0.19	0.36	1.07	0.09
	30	0.09	0.21	0.35	0.65	0.05		30	0.13	0.30	0.50	1.01	0.09
	40	0.13	0.27	0.42	0.58	0.05		40	0.17	0.38	0.59	0.90	0.09
	50	0.15	0.31	0.42	0.47	0.05		50	0.21	0.43	0.62	0.74	0.09
	60	0.18	0.30	0.34	0.36	0.05		60	0.23	0.43	0.53	0.55	0.09
	70	0.15	0.20	-	0.21	0.05		70	0.22	0.33	-	0.36	0.08
80	0.07	-	-	0.10	0.04	80	0.14	-	-	0.17	0.06		

Devcan 6.8.0, August 2014, National Cancer Institute (<http://surveillance.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic could not be calculated.

A percent of 0.00 represents a value that is below 0.005.